

Mutations at *pipX* Suppress Lethality of P_{II}-Deficient Mutants of *Synechococcus elongatus* PCC 7942[∇]

Javier Espinosa, Miguel Angel Castells, Karim Boumediene Laichoubi, and Asunción Contreras*

División de Genética, Universidad de Alicante, Apartado 99, E-03080 Alicante, Spain

Received 27 April 2009/Accepted 26 May 2009

The P_{II} proteins are found in all three domains of life as key integrators of signals reflecting the balance of nitrogen and carbon. Genetic inactivation of P_{II} proteins is typically associated with severe growth defects or death. However, the molecular basis of these defects depends on the specific functions of the proteins with which P_{II} proteins interact to regulate nitrogen metabolism in different organisms. In *Synechococcus elongatus* PCC 7942, where P_{II} forms complexes with the NtcA coactivator PipX, attempts to engineer P_{II}-deficient strains failed in a wild-type background but were successful in *pipX* null mutants. Consistent with the idea that P_{II} is essential to counteract the activity of PipX, four different spontaneous mutations in the *pipX* gene were found in cultures in which *glnB* had been genetically inactivated.

Cyanobacteria are phototrophic organisms that perform oxygenic photosynthesis. Autotrophic growth requires the constant assimilation of ammonium via the coordinated action of glutamine synthetase (GS) and glutamate synthase, also known as the GS-GOGAT cycle, resulting in consumption of 2-oxoglutarate (34). Due to the lack of 2-oxoglutarate dehydrogenase in cyanobacteria, synthesis of 2-oxoglutarate represents the final step in the oxidative branch of the trichloroacetic acid cycle and directly links 2-oxoglutarate levels to nitrogen assimilation (35). Thus, 2-oxoglutarate accumulates during nitrogen starvation, making this metabolite an excellent indicator of the intracellular carbon-nitrogen balance (12, 25).

In cyanobacteria, multiple metabolic and developmental processes are induced by nitrogen starvation. NtcA, the global regulator for nitrogen control, activates genes involved in nitrogen assimilation, heterocyst differentiation, and acclimation to nitrogen starvation (20, 30, 41). 2-Oxoglutarate, the signal of nitrogen deficiency, stimulates binding of NtcA to target sites (45), transcription activation in vitro (44), and complex formation between the global nitrogen regulator NtcA and its coactivator factor PipX, a regulatory protein conserved in cyanobacteria (5, 9). The interaction between PipX and NtcA is known to be relevant for maximal activation of NtcA-dependent genes under nitrogen limitation (9, 10). PipX-deficient cultures of *Synechococcus elongatus* PCC 7942 showed reduced activity of nitrogen assimilation enzymes, retarded nitrogen induction, a slower rate of nitrate consumption, and when subjected to nitrogen starvation, retarded phycobilisome degradation and faster reduction of the chlorophyll content (10).

The homotrimeric P_{II} protein is one of the most conserved and widespread signal transduction proteins in nature and plays key roles in nitrogen assimilatory processes (28). P_{II} proteins contain three binding sites (one per subunit) for 2-oxoglutarate and ATP, and their primary function is to reg-

ulate, by direct protein-protein interactions, the activity of proteins implicated in nitrogen metabolism (reviewed in reference 13). In cyanobacteria, several proteins are known to form complexes with P_{II}. The first two P_{II} receptors were identified in *S. elongatus*: the enzyme *N*-acetyl-L-glutamate kinase (NAGK), a P_{II} target conserved across domains of life during the evolution of oxygenic photosynthetic organisms (5, 6, 42), and the regulatory factor PipX (5, 9). The nonconserved membrane protein PamA was identified as a P_{II} receptor in *Synechocystis* sp. strain PCC 6803 (37). Structural and functional details are only known for the P_{II}-NAGK complex (29). This complex consists of two polar P_{II} trimers sandwiching one ring-like hexameric NAGK, with the flexible T loop, a key element for regulatory interactions, adopting a novel compact shape. Other P_{II} functions for which direct protein-protein interactions have not been reported yet include the control of nitrate transport (23, 27), nitrate reductase (43), and the control of inorganic carbon transport (21).

P_{II} proteins bind 2-oxoglutarate and ATP synergistically. In *S. elongatus* and *Synechocystis* sp. strain PCC 6803, the T loop is phosphorylated at a seryl residue (S49) located at the apex of the solvent-exposed T loop (12). The phosphorylation status of P_{II} correlates with 2-oxoglutarate levels, both being maximal during nitrogen starvation. ATP in concert with elevated 2-oxoglutarate levels prevents complex formation of P_{II} with either NAGK or PipX (9, 31), suggesting that PipX-P_{II} complexes could also have a function under nitrogen-sufficient conditions. In this context, previous analyses indicated that PipX is not required for P_{II}-dependent functions like ammonium inhibition of nitrate transport (10) or stimulation of NAGK activity (8). However, we show here that P_{II} does, indeed, affect PipX functions. Our results indicate that P_{II} is essential under standard growth conditions in *S. elongatus* and that the *pipX* gene is a target of suppressor mutations in P_{II}-deficient cultures.

* Corresponding author. Mailing address: División de Genética, Facultad de Ciencias, Universidad de Alicante, E-03080 Alicante, Spain. Phone: 34 96 590 3957. Fax: 34 96 590 9569. E-mail: contrera@ua.es.

[∇] Published ahead of print on 29 May 2009.

MATERIALS AND METHODS

Strains and growth conditions. The strains and plasmids used in this work are listed in Table 1. Constructs and genomic mutations were analyzed by automated

TABLE 1. Strains and plasmids used in this work^a

Strain or plasmid	Genotype or relevant characteristic(s)	Source or reference
Strains		
<i>E. coli</i> DH5 α	F ⁻ ϕ 80dlacZ Δ M15 Δ (<i>lacZYA-argF</i>)U169 <i>endA1 recA1 hsdR17</i> (r _K ⁻ m _K ⁺)	16
<i>Synechococcus</i> sp. strain PCC 7942	Wild-type strain	Pasteur Culture Collection
<i>Synechococcus</i> sp. strain MP2	P _{II} ⁻ (<i>glnB</i> ::CK2) Km ^r	14
<i>Synechococcus</i> sp. strain MP2-A	<i>glnB</i> ::CK2 P _{II} ^{S49A} Sm ^r Km ^r	27
<i>Synechococcus</i> sp. strain MP2-E	<i>glnB</i> ::CK2 P _{II} ^{S49E} Sm ^r Km ^r	27
<i>Synechococcus</i> sp. strain GlnBK	<i>glnB/glnB</i> ::CK2 Km ^r	This work
<i>Synechococcus</i> sp. strain GlnBS(+)	<i>glnB/glnB</i> ::CS3(+) Sm ^r	This work
<i>Synechococcus</i> sp. strain GlnBS(-)	<i>glnB</i> ::CS3(-) Sm ^r	This work
<i>Synechococcus</i> sp. strain GlnBS(-) ^{T loop + 7}	<i>glnB</i> ::CS3(-)/ <i>glnB</i> 133_134insATAAAGCTTATCGATACCGTC Sm ^r	This work
<i>Synechococcus</i> sp. strain SA410	PipX(Con) [Φ(CK1- <i>pipX</i>)] Km ^r	9
<i>Synechococcus</i> sp. strain SA591	PipX ⁻ (<i>pipX</i> ::CK1) Km ^r	9
<i>Synechococcus</i> sp. strain SA410-GlnBS(-)	PipX(Con) <i>glnB/glnB</i> ::CS3(-) Sm ^r Km ^r	This work
<i>Synechococcus</i> sp. strain SA591-GlnBS(-)	PipX ⁻ <i>glnB</i> ::CS3(-) Sm ^r Km ^r	This work
<i>Synechococcus</i> sp. strain SA410-GlnBS(+)	PipX(Con) <i>glnB/glnB</i> ::CS3(+) Sm ^r Km ^r	This work
<i>Synechococcus</i> sp. strain SA591-GlnBS(+)	PipX ⁻ <i>glnB</i> ::CS3(+) Sm ^r Km ^r	This work
Plasmids		
pBluescriptII SK(+)	Cloning vector, Ap ^r	Stratagene
pPM128	<i>glnB</i> ::CK2, Km ^r	14
pUAGC59	pBluescript SK(+) with 1.8 kb from <i>pipX</i> region, Ap ^r	9
pUAGC453	pBluescriptII SK(+) with CS3, Ap ^r Sm ^r	40
pUAGC700	pBluescriptII SK(+) with 1.3-kb <i>glnB</i> genomic region, Ap ^r	This work
pUAGC701	pUAGC700 with CS3 into <i>glnB</i> [<i>glnB</i> ::CS3(-)], Ap ^r Sm ^r	This work
pUAGC702	Same as pUAGC701, with CS3 in opposite orientation [<i>glnB</i> ::CS3(+)], Ap ^r Sm ^r	This work

^a Note that some strains are heteroallelic for *glnB* and therefore unstable (see text for details).

dideoxy DNA sequencing. All cloning procedures were carried out with *Escherichia coli* DH5 α by using standard techniques.

S. elongatus strains were routinely grown photoautotrophically at 30°C while shaking under constant illumination (40 μ mol photons m⁻² s⁻¹) provided by cool white fluorescent lights. The media used were BG11 (BG11₀ plus 17.5 mM NaNO₃ and 10 mM HEPES/NaOH, pH 7.8) or BG11-NH₄ (BG11₀ plus 10 mM NH₄Cl and 10 mM HEPES/NaOH, pH 7.8). For growth on plates, the medium was solidified by addition of 1% (wt/vol) agar. Plates were incubated at 30°C under constant illumination. The transformation procedure was essentially as described previously (15). Cells were incubated for 48 h at 30°C under constant illumination on nitrocellulose filters (Millipore), and transformants were selected on kanamycin-, chloramphenicol-, or streptomycin-containing BG11 plates. The antibiotic concentrations used were 10 μ g ml⁻¹ (kanamycin), 5 μ g ml⁻¹ (chloramphenicol), and 2 μ g ml⁻¹ (streptomycin).

Construction of plasmids. To construct pUAGC700, a 1.3-kb genomic DNA fragment including *glnB* was PCR amplified with primers 5' GTCTAGAATTCCTTCTGTTGTGATG 3' and 5' GGACGGGATCCCTACCACCGCCTTT 3', cut with EcoRI and BamHI, and cloned into pBluescript SK(+). A HincII-EcoRV fragment containing the CS3 cassette from pUAGC453 was then cloned into the AfeI site of pUAGC700, rendering plasmids pUAGC701 and pUAGC702.

PCR verification of the correct integration of resistance cassettes in *S. elongatus* was carried out with primers 5' ACAAAACGGTTTACCAGCAT 3' (CS3-2F) or 5' CAACAAAGCTCTCATCAACC 3' (CK2-F) and 5' AACTGCAGTCGACGCTGACTTAGATTGCGTCG 3' (GlnB-1R). Detection of *glnB* alleles by PCR were carried out with primers 5' GGCTTAAGGAGAATTCCTTGAAGAAG 3' (GlnB-1F) and GlnB-1R.

RT-PCR analysis. For analysis of *pipX* mRNA abundance under nitrate growth conditions, cells were grown under standard conditions until they reached an optical density at 750 nm of 0.5. Fifty-milliliter aliquots were removed from the cultures for RNA extraction. The samples were rapidly chilled on ice and centrifuged, and pellets were stored at -80°C. Total RNA was isolated by the hot phenol method. Reverse transcription (RT)-PCR analysis of *pipX* mRNA was performed with 0.5 μ g of total *S. elongatus* RNA that was retrotranscribed in a total volume of 30 μ l with RevertAid H Minus Moloney murine leukemia virus reverse transcriptase (Fermentas) and primers 5' CTGCTCTGAATTCCTAG

CTGGCTACAG 3' (for *pipX*) and 5' GGGGGTACCTTGATTACAGAC 3' (for *sipA*). Ten microliters of the retrotranscription reaction mixture was subjected to 30 PCR cycles with primers 5' GAGAATTCGCTTCCGAGAACTACC 3' and 5' CTGCTCTGAATTCCTAGCTGGCTACAG 3' and NETZYME DNA polymerase. Twenty-four PCR cycles with primers 5' ACCCGGATCCCTCTATGGATTTTG 3' and 5' GGGGGTACCTTGATTACAGAC 3' were used for *sipA*, which was used as a loading control. For each pair of primers, a parallel reaction was carried out without reverse transcriptase as a control for DNA contamination of RNA preparations.

RESULTS

The *glnB* gene is essential in *S. elongatus*. Inactivation of the *glnB* gene of *S. elongatus* was attempted by routinely used procedures of allelic replacement with two different cassettes providing kanamycin (CK2) or streptomycin (CS3) resistance. The three constructs used to inactivate *glnB* are schematically represented in Fig. 1A. Plasmid pPM128, previously used to generate *glnB* null strain MP2 (14), was used as a source of the *glnB*::CK2 allele. Two other constructs, each one carrying the CS3 cassette in a different orientation in the AfeI site of the *glnB* gene, were also produced, *glnB*::CS3(+) and *glnB*::CS3(-). In all three cases, viable antibiotic-resistant transformants were obtained and subsequent PCR amplifications produced bands matching the expected sizes of cassettes CK2 (307 bp), CS3(-) (265 bp), and CS3(+) (254 bp), thus confirming their presence in the appropriated location within the *S. elongatus glnB* gene (Fig. 1B, lanes K, S+, and S-).

Due to the presence of multiple chromosome copies in cyanobacteria (3), gene inactivation in *S. elongatus* requires verification that allelic replacement has been completed. In this




FIG. 1. Genetic inactivation of *glnB*. (A) Schematic representation of *glnB* null alleles. Relevant restriction sites in *glnB* (open arrows) are indicated. Positions of primers used to verify allele replacement are indicated by black arrows. (B) PCR verification of the correct integration of resistance cassettes with primers CS3-2F (S2F) or CK2-F (K2F) and *GlnB*-1R (1R) in a representative clone after transformation of *S. elongatus* with *glnB::CK2* (lane K), *glnB::CS3(-)* (lane S-), and *glnB::CS3(+)* (lane S+). Lane L, size marker (GeneRuler 100-bp Plus DNA Ladder from Fermentas). PCR products corresponding to specific alleles are indicated at the right, and relevant marker sizes in base pairs are shown at the left. (C) Detection of *glnB* alleles by PCR with primers *GlnB*-1F (1F) and *GlnB*-1R (1R) in three independent clones after four consecutive transfers onto selective media. Other details are the same as in panel B.

context, the persistence of wild-type alleles after transfers to new media under selective conditions indicates that the targeted gene is essential. That was the case with the three types of clones generated here, since after several consecutive transfers onto selective plates they remained heteroallelic for *glnB* or gave rise to gene rearrangements (see below). As shown in Fig. 1C, wild-type *glnB* alleles were clearly observed by PCR in all of the independent *glnB::CK2* and *glnB::CS3(+)* transformants analyzed (lanes K and S+ and data not shown). These two types of clones retained wild-type *glnB* alleles after at least six transfers of colonies onto selective media. In contrast, some of the *glnB::CS3(-)* transformants apparently lost wild-type *glnB* alleles after the fourth transfer onto streptomycin-containing media (Fig. 1C, lanes S-, and data not shown). The viability of these apparently segregated *GlnB*(-) clones was compromised, since they failed to grow when transferred from solid to liquid media (data not shown). However, additional subculturing of these *glnB::CS3(-)* transformants on plates was possible and allowed the detection of a longer PCR product (Fig. 2A). Sequence analysis of this unexpected PCR product revealed the presence of a 21-bp insertion into the AfeI site of the *glnB* gene, corresponding to the ends of the CS3 cassette (allele *glnB133_134insATAAAGCTTATCGATACCGTC*, [hereafter, *glnB133_134ins*]). The rearrangement was probably generated by recombination between homologous sequences at the two junctions of the CS3 cassette with the *glnB* gene and resulted in seven extra amino acids between residues E44 and R45 (Fig. 2B), located at the flexible T loop of P_{II}. In contrast to their predecessor *GlnB*(-) clones, derivatives in which the *glnB133_134ins* allele was detected (strain *GlnB*(-)^{T loop + 7}) could be cultivated in liquid media. Detection of this spontaneously generated rearrangement of the *glnB::CS3(-)* allele suggested that the protein product, designated P_{II}^{T loop + 7},

conferred a selective advantage on *S. elongatus* cultures deficient in P_{II}.

In summary, attempts to generate *S. elongatus* P_{II}-deficient cells with three different null alleles, *glnB::CK2*, *glnB::CS3(+)*, and *glnB::CS3(-)*, failed. The alleles or allelic pairs detected by PCR in particular cultures from transformant clones were *glnB/glnB::CK2*, *glnB/glnB::CS3(+)*, *glnB/glnB::CS3(-)*, *glnB::CS3(-)*, and *glnB::CS3(-)/glnB133_134ins*. Since cultures in which *glnB::CS3(-)* was apparently in homozygosis were invariably classified as *glnB::CS3(-)/glnB133_134ins* in the next PCR test, it seems likely that viable cells in those cultures had already acquired the *glnB133_134ins* allele. Taken together, the results indicated that *glnB* is essential for culture growth, at least under our laboratory conditions.

***glnB* null mutants carry mutations in *pipX*.** On several occasions, we noticed that our P_{II}-deficient cultures (MP2 strains) carried secondary mutations affecting the *pipX* coding sequence. A point mutation (allele *pipX160C>T*, which encodes the substitution R54C) was found fortuitously while cloning *glnB* sequences for plasmid construction. Since PCR amplification of genomic DNA, followed by direct sequencing, confirmed that the *pipX160C>T* mutation was present in the MP2 strain used for DNA amplification, we obtained a "new" culture of the MP2 strain (a gift from K. Forchhammer) to perform further analysis. Although single-PCR amplification and subsequent sequencing demonstrated the presence of the wild-type *pipX* allele in this strain, after performing RT-PCR assays, we became aware that this strain also carried sequences with a 22-bp deletion in *pipX* (allele *pipX25_46del*) causing a frameshift after position N7. Subsequent PCR analysis confirmed that both alleles, *pipX25_46del* and wild-type *pipX*, were present in this MP2 strain that we had received from K. Forchhammer's laboratory. Furthermore, a strain with the same or-




FIG. 2. The spontaneously generated *glnB* rearrangement. (A) Schematic representation of the *glnB133_134ins* allele and detection of *glnB* and *glnB133_134ins* alleles with primers GlnB-1R (1R) and GlnB-1F (1F). Wild type, lane 2; GlnBS(-)^{T loop +7}, lane 1. Relevant alleles are indicated at the left. Other details are as in Fig. 1. (B) Alignment of DNA sequences at the *glnB*-CS3 and CS3-*glnB* junctions from the *glnB*::CS3(-) allele and sequences present after the rearrangement. Predicted amino acids are shown in one-letter code. Nucleotides and predicted amino acids derived from the CS3 cassette are in bold.

igin that had already been genetically manipulated in another laboratory (a gift from F. Fernández-Piñas) was also found to be heteroallelic (*pipX/pipX25_46del*) after PCR analysis.

Sequence analysis of *pipX* from strains MP2-E and MP2-A, which both carry the same *glnB*::CK2 insertion, revealed a distinct *pipX* point mutation in each strain, allele *pipX194T>A*, which encodes substitution L65Q (in strain MP2-E), and a single-base deletion upstream of *pipX* (*pipX-92delT*) (in strain MP2-A). Since several independent PCR amplifications and subsequent sequence analyses gave identical results, these strains were apparently homoallelic at the time of the analysis. The presence of two distinct mutations in strains with the same origin suggested that the parental strain (MP2) might be heteroallelic for *pipX*. Further evidence for the genetic heterogeneity of *pipX* alleles in P_{II}-deficient strains was also obtained retrospectively. In particular, three different *pipX* alleles (*pipX*, *pipX194T>A*, and *pipX25_46del*) were detected in strains derived from the same MP2 culture in our laboratory after selection for different genetic traits. Table 2 summarizes the mutant alleles selected in the different MP2 derivatives analyzed.

Figure 3 illustrates the information on the locations and

TABLE 2. Spontaneous mutations found at *pipX* in strains carrying *glnB*::CK2 alleles in homozygosis

Mutation ^a	Strain origin	P _{II} protein expressed
<i>pipX25_46del</i>	MP2	None
<i>pipX160C>T</i>	MP2	None
<i>pipX194T>A</i>	MP2	None
<i>pipX-92delT</i>	MP2-A	P _{II} ^{S49A}
<i>pipX194T>A</i>	MP2-E	P _{II} ^{S49E}

^a The mutation position is the distance, in nucleotides, from the start codon to the mutation site.

predicted impacts on the amino acid sequence of the spontaneous mutations found at the *pipX* open reading frame or the *pipX* promoter region. The location of *pipX-92delT*, just upstream of the *pipX* open reading frame, suggested that it could be a promoter-down mutation. To obtain additional evidence of the impact of the *pipX-92delT* mutation on *pipX* expression, we compared *pipX* mRNA levels in strains carrying the wild-type or mutant allele by RT-PCR. As shown in Fig. 4, *pipX* mRNA levels were significantly reduced in the *pipX-92delT* (represented as X^{P↓}) mutant, while transcripts from unrelated controls were not altered, supporting the idea that the *pipX-92delT* mutation diminished the transcription of *pipX*. The effects of the R54C and L65Q mutations on PipX interactions will be analyzed elsewhere. Importantly, none of the *pipX* changes were found in wild-type *S. elongatus* strains, suggesting that the four mutations detected were bona fide suppressor mutations and that loss-of-function mutations at *pipX* alleviate the P_{II} deficiency phenotype.

Inactivation by *pipX* suppresses the lethality associated with P_{II} deficiency. The finding of different putative suppressor mutations at *pipX* suggested that the lethality phenotype associated with P_{II} deficiency was due to an excess of active PipX protein, harmful to *S. elongatus* cells unless counteracted by P_{II}. If that were the case, inactivation of *glnB* alleles would be facilitated by elimination of PipX. To test this idea, we separately introduced the *glnB*::CS3(+) and *glnB*::CS3(-) alleles into a PipX⁻ strain (SA591) in which the CK1 cassette replaced part of the *pipX* coding sequence. As an additional control, the *glnB*::CS3(+) and *glnB*::CS3(-) alleles were introduced in parallel into kanamycin-resistant strain SA410 [PipX-(Con)], where the *pipX* gene is transcribed from a constitutive promoter present in the CK1 cassette. The only phenotypic features noticed in the PipX(Con) strain were a small increase in the nitrogen induction of NtcA-dependent promoters (10)




FIG. 3. Spontaneous mutations at *pipX*. (A) Comparison of the predicted amino acid sequences at the relevant regions of PipX, shown in one-letter code. Amino acids that differ from the wild type are in bold and boxed. The asterisks and triangle on the *pipX* genomic map indicate, respectively, point mutations and a deletion at mutant alleles. The vertical arrow indicates the site of the *pipX-92delT* mutation.

and a level of PipX protein twice that of the wild type (data not shown).

PCR analyses confirmed the correct integration of the CS3(+) and CS3(-) cassettes within *glnB* for each of the two strains used (Fig. 5A). After several transfers onto selective media, wild-type *glnB* alleles were observed in the PCR analysis of transformants from PipX(Con) (Fig. 5B, lanes X^{CS-} and X^{CS+}). Thus, as previously observed with the wild-type strain (Fig. 1C), the presence of active *pipX* alleles prevented homozygosity of the *glnB*::CS3(+) and *glnB*::CS3(-) alleles. In contrast, homozygosity of both the *glnB*::CS3(+) and *glnB*::CS3(-) alleles was easily achieved in the PipX⁻ background, as shown by the absence of *glnB* alleles in these PipX⁻ recombinants (Fig. 5B, lanes X^{-S-} and X^{-S+}). These results confirmed that the presence of functional *pipX* genes interfered with the inactivation of *glnB*, implying that an intact *glnB* gene is required for viability only in the presence of an active *pipX* gene.

DISCUSSION

Genetic inactivation of the genes that encode P_{II} proteins in many microorganisms leads to severe growth defects or death (2, 7, 32, 33, 48). In cyanobacteria, growth defects and death associated with P_{II} deficiency have also been reported, but depending on the model system, particular laboratory strains, or culture conditions, the *glnB* gene was considered essential or not essential. In *S. elongatus* PCC 7942 and *Synechocystis* sp. strain PCC 6803, *glnB* null mutants have been reported (14, 21, 24). In *Nostoc punctiforme* PCC 73102, *glnB* null mutants could not be obtained and the *glnB* gene was reported to be essential (17). In *Anabaena* sp. strain PCC 7120, previous attempts to inactivate *glnB* were unsuccessful (26). However, that report was followed by two others in which the obtention of *glnB* null mutants was described (38, 49). In the most recent of these, the




FIG. 4. Effect of *pipX-92delT* mutation on *pipX* transcript levels. RT-PCR amplification of *pipX* and *sipA* (used as a loading control) in the wild type (WT) and MP2-A (X^P). A representative experiment from one of two independent RNA extractions is shown. The values on the left are molecular sizes in base pairs.

authors showed that successful inactivation of *glnB* could only be achieved when the expression pattern of downstream genes of unknown function was altered (38). In the present work, we show the importance of P_{II} proteins for viability in the model system of *S. elongatus*, a cyanobacterium where successful inactivation of *glnB* was previously reported. Attention is called to the occurrence of suppressor mutations at the *pipX* gene of P_{II}-deficient strains.

The metabolic basis of the defect of *glnB* null mutants in the studied microorganisms appears to be diverse. In *E. coli*, *glnB* null mutants show elevated activity of the nitrogen response regulator NtrC that results in increased activity of the Nac protein which, in turn, represses *serA* to levels insufficient for normal growth (4). This regulatory cascade is not even conserved in other enterobacteria (36). In *Rhodospirillum rubrum*, the other model system for which the metabolic defect is known, the poor-growth phenotype is due to an excess of GS activity (47). In *S. elongatus*, the gene that encodes GS (*glnA*) is activated by NtcA and GS activity is associated with P_{II} deficiency in *S. elongatus*. However, none of the MP2 strains analyzed here carried mutations in the *glnA* or *ntcA* gene. Since mutations inactivating the *ntcA* gene have been reported (46), the impaired phenotype of *S. elongatus glnB* null mutants appears to be related to an unknown PipX activity, thus raising questions of whether *glnB* null mutants obtained in other cyanobacteria contain compensatory changes at *pipX* and whether previous inactivation of *pipX* would facilitate recovery of *glnB* null mutants in species that have been so far recalcitrant to *glnB* inactivation.

Previous studies with P_{II}-deficient strains carrying the *glnB*::CK2 allele (strains MP2) suggested that P_{II} is required to stimulate NtcA activity under conditions of nitrogen deprivation (11, 39) and has an inhibitory role with nitrate as a nitrogen source (1, 11). Structural determination of the NAGK-P_{II} complexes (29) provided a rationale for the negative impact that mutations at Ser49, the phosphorylatable residue of P_{II}, have in interactions with NAGK (5, 18, 29). Although no differences among P_{II}, P_{II}^{S49A}, and P_{II}^{S49E} were found regarding interactions with PipX (5) or in vitro interactions with the effectors 2-oxoglutarate and ATP (27), Ser49 has nevertheless been implicated in the regulation of NtcA activity. Two studies carried out with MP2 strains claimed that substitution of P_{II}^{S49A}, but not P_{II}^{S49E}, for P_{II} prevented the induction of NtcA-dependent genes (27, 39). Since the authors assumed that mutations S49A and S49E may mimic, respectively, the nonphosphorylated and phosphorylated states of P_{II}, the inference was that P_{II} phosphorylation was important for NtcA function (27, 39). Since we cannot safely exclude the possibility




FIG. 5. Genetic inactivation of *glnB* in *pipX* mutants. (A) PCR verification of the correct integration of resistance cassettes in a representative clone after transformation of *pipX* (lanes X⁻) and *pipX*(Con) (lanes X^c) mutant strains with *glnB*::CS3(-) (lanes S⁻) and *glnB*::CS3(+) (lanes S⁺). (B) Segregation of *glnB* alleles verified by PCR in three independent clones of *pipX* and *pipX*(Con) mutant strains transformed with *glnB*::CS3(-) and *glnB*::CS3(+) after four consecutive transfers onto selective media. See Fig. 1B for other details and lane labeling.

that the *pipX* mutations detected in strains MP2-A and MP2-E (Table 2) were already present in the particular cultures used then, inferences about NtcA-dependent activities in MP2 derivatives should be made with caution. Isogenic strains are needed to reevaluate the effect of P_{II} point mutations on NtcA function.

The effect of 2-oxoglutarate on PipX-P_{II} and PipX-NtcA complexes suggests that these protein associations are physiologically relevant at low and high C/N ratios, respectively (9), but little is known about the physiological role of PipX-P_{II} complexes. As a starting point to get insights into the function of PipX-P_{II} complexes, we contemplated three simple scenarios that are not necessarily incompatible: (i) PipX regulates P_{II} functions, (ii) P_{II} regulates PipX functions, and (iii) the complex itself has its own functions. The possibility that PipX-P_{II} complexes bind to additional proteins, fed by recent reports of ternary complexes in which P_{II} proteins are involved (19, 22), cannot be excluded at present. On the other hand, we did explore whether PipX is required for the P_{II}-dependent inhibition of nitrate transport by ammonium (10), stimulation of NAGK activity (8), or P_{II} phosphorylation (data not shown) and obtained negative responses in all of the cases examined. However, the results presented here strongly support a role for P_{II} in counteracting unknown but toxic functions of PipX in *S. elongatus*.

The features and occurrence of the spontaneous mutations at *pipX* suggest that P_{II} deficiency is best suppressed by partial rather than complete loss-of-function mutations at *pipX*, and some gene dosage (or specific activity) of *pipX* may improve the fitness of MP2 strains. This would explain the genetic heterogeneity noticed in *pipX* alleles. The coexistence of wild-type and mutant *pipX* alleles in several P_{II}-deficient cultures, and particularly the apparent stability of the allelic combination *pipX/pipX25_46del*, may be an adaptive mechanism for P_{II}-deficient cultures of *S. elongatus*, raising questions about the extent of genetic heterogeneity in cyanobacteria.

ACKNOWLEDGMENTS

This work was supported by grants BFU2006-12424 and ACOMP06/083. M.A.C. is the recipient of a predoctoral fellowship from the Universidad de Alicante.

We thank R. Dixon, K. Forchhammer, and V. Rubio for very helpful discussions and N. Tandeau de Marsac, K. Forchhammer, and F. Fernández-Piñas for plasmids and/or strains.

REFERENCES

- Aldehni, M. F., and K. Forchhammer. 2006. Analysis of a non-canonical NtcA-dependent promoter in *Synechococcus elongatus* and its regulation by NtcA and P_{II}. *Arch. Microbiol.* **184**:378–386.
- Atkinson, M. R., and A. J. Ninfa. 1998. Role of the GlnK signal transduction protein in the regulation of nitrogen assimilation in *Escherichia coli*. *Mol. Microbiol.* **29**:431–447.
- Binder, B. J., and S. W. Chisholm. 1990. Relationship between DNA cycle and growth rate in *Synechococcus* sp. strain PCC 6301. *J. Bacteriol.* **172**:2313–2319.
- Blauwkamp, T. A., and A. J. Ninfa. 2002. Nac-mediated repression of the *serA* promoter of *Escherichia coli*. *Mol. Microbiol.* **45**:351–363.
- Burillo, S., I. Luque, I. Fuentes, and A. Contreras. 2004. Interactions between the nitrogen signal transduction protein P_{II} and N-acetyl glutamate kinase in organisms that perform oxygenic photosynthesis. *J. Bacteriol.* **186**:3346–3354.
- Chen, Y. M., T. S. Ferrar, E. Lohmeir-Vogel, N. Morrice, Y. Mizuno, B. Berenger, K. K. Ng, D. G. Muench, and G. B. Moorhead. 2006. The P_{II} signal transduction protein of *Arabidopsis thaliana* forms an arginine-regulated complex with plastid N-acetyl glutamate kinase. *J. Biol. Chem.* **281**:5726–5733.
- de Zamaroczy, M. 1998. Structural homologues P_{II} and P_Z of *Azospirillum brasilense* provide intracellular signalling for selective regulation of various nitrogen-dependent functions. *Mol. Microbiol.* **29**:449–463.
- Espinosa, J., M. A. Castells, K. Forchhammer, and A. Contreras. 2008. Regulatory interactions involved in nitrogen signalling in cyanobacteria: partner swapping and role of PipX-P_{II} complexes in *Synechococcus* sp. PCC 7942, p. 361–362. In S. Netherlands (ed.), *Biological nitrogen fixation: towards poverty alleviation through sustainable agriculture*, vol. 42. Springer, New York, NY.
- Espinosa, J., K. Forchhammer, S. Burillo, and A. Contreras. 2006. Interaction network in cyanobacterial nitrogen regulation: PipX, a protein that interacts in a 2-oxoglutarate dependent manner with P_{II} and NtcA. *Mol. Microbiol.* **61**:457–469.
- Espinosa, J., K. Forchhammer, and A. Contreras. 2007. Role of the *Synechococcus* PCC 7942 nitrogen regulator protein PipX in NtcA-controlled processes. *Microbiology* **153**:711–718.
- Fadi Aldehni, M., J. Sauer, C. Spielhauer, R. Schmid, and K. Forchhammer. 2003. Signal transduction protein P_{II} is required for NtcA-regulated gene expression during nitrogen deprivation in the cyanobacterium *Synechococcus elongatus* strain PCC 7942. *J. Bacteriol.* **185**:2582–2591.
- Forchhammer, K. 2004. Global carbon/nitrogen control by P_{II} signal transduction in cyanobacteria: from signals to targets. *FEMS Microbiol. Rev.* **28**:319–333.
- Forchhammer, K. 2008. P_{II} signal transducers: novel functional and structural insights. *Trends Microbiol.* **16**:65–72.
- Forchhammer, K., and N. Tandeau de Marsac. 1995. Functional analysis of the phosphoprotein P_{II} (*glnB* gene product) in the cyanobacterium *Synechococcus* sp. strain PCC 7942. *J. Bacteriol.* **177**:2033–2040.
- Golden, S. S., and L. A. Sherman. 1984. Optimal conditions for genetic transformation of the cyanobacterium *Anacystis nidulans* R2. *J. Bacteriol.* **158**:36–42.
- Hanahan, D. 1985. Techniques for transformation of *Escherichia coli*, p. 109–135. In D. M. Glover (ed.), *DNA cloning*. IRL Press, Oxford, United Kingdom.
- Hanson, T. E., K. Forchhammer, N. Tandeau de Marsac, and J. C. Meeks. 1998. Characterization of the *glnB* gene product of *Nostoc punctiforme* strain ATCC 29133: *glnB* or the P_{II} protein may be essential. *Microbiology* **144**(Pt. 6):1537–1547.
- Heinrich, A., M. Maheswaran, U. Ruppert, and K. Forchhammer. 2004. The

- Synechococcus elongatus* P signal transduction protein controls arginine synthesis by complex formation with *N*-acetyl-L-glutamate kinase. *Mol. Microbiol.* **52**:1303–1314.
19. Heinrich, A., K. Woyda, K. Brauburger, G. Meiss, C. Detsch, J. Stulke, and K. Forchhammer. 2006. Interaction of the membrane-bound GlnK-AmtB complex with the master regulator of nitrogen metabolism TnrA in *Bacillus subtilis*. *J. Biol. Chem.* **281**:34909–34917.
 20. Herrero, A., A. M. Muro-Pastor, and E. Flores. 2001. Nitrogen control in cyanobacteria. *J. Bacteriol.* **183**:411–425.
 21. Hisbergues, M., R. Jeanjean, F. Josef, N. Tandeau de Marsac, and S. Bedu. 1999. Protein P_{II} regulates both inorganic carbon and nitrate uptake and is modified by a redox signal in *Synechocystis* PCC 6803. *FEBS Lett.* **463**:216–220.
 22. Huergo, L. F., M. Merrick, F. O. Pedrosa, L. S. Chubatsu, L. M. Araujo, and E. M. Souza. 2007. Ternary complex formation between AmtB, GlnZ and the nitrogenase regulatory enzyme DraG reveals a novel facet of nitrogen regulation in bacteria. *Mol. Microbiol.* **66**:1523–1535.
 23. Kloft, N., and K. Forchhammer. 2005. Signal transduction protein P_{II} phosphatase PphA is required for light-dependent control of nitrate utilization in *Synechocystis* sp. strain PCC 6803. *J. Bacteriol.* **187**:6683–6690.
 24. Kobayashi, M., N. Takatani, M. Tanigawa, and T. Omata. 2005. Posttranslational regulation of nitrate assimilation in the cyanobacterium *Synechocystis* sp. strain PCC 6803. *J. Bacteriol.* **187**:498–506.
 25. Laurent, S., H. Chen, S. Bedu, F. Ziarelli, L. Peng, and C. C. Zhang. 2005. Nonmetabolizable analogue of 2-oxoglutarate elicits heterocyst differentiation under repressive conditions in *Anabaena* sp. PCC 7120. *Proc. Natl. Acad. Sci. USA* **102**:9907–9912.
 26. Laurent, S., K. Forchhammer, L. Gonzalez, T. Heulin, C. C. Zhang, and S. Bedu. 2004. Cell-type specific modification of P_{II} is involved in the regulation of nitrogen metabolism in the cyanobacterium *Anabaena* PCC 7120. *FEBS Lett.* **576**:261–265.
 27. Lee, H. M., E. Flores, K. Forchhammer, A. Herrero, and N. Tandeau De Marsac. 2000. Phosphorylation of the signal transducer P_{II} protein and an additional effector are required for the P_{II}-mediated regulation of nitrate and nitrite uptake in the Cyanobacterium *Synechococcus* sp. PCC 7942. *Eur. J. Biochem.* **267**:591–600.
 28. Leigh, J. A., and J. A. Dodsworth. 2007. Nitrogen regulation in bacteria and archaea. *Annu. Rev. Microbiol.* **61**:349–377.
 29. Llácer, J. L., A. Contreras, K. Forchhammer, C. Marco-Marin, F. Gil-Ortiz, R. Maldonado, I. Fita, and V. Rubio. 2007. The crystal structure of the complex of P_{II} and acetylglutamate kinase reveals how P_{II} controls the storage of nitrogen as arginine. *Proc. Natl. Acad. Sci. USA* **104**:17644–17649.
 30. Luque, I., G. Zabulon, A. Contreras, and J. Houmard. 2001. Convergence of two global transcriptional regulators on nitrogen induction of the stress-acclimation gene nblA in the cyanobacterium *Synechococcus* sp. PCC 7942. *Mol. Microbiol.* **41**:937–947.
 31. Maheswaran, M., C. Urbanke, and K. Forchhammer. 2004. Complex formation and catalytic activation by the P_{II} signaling protein of *N*-acetyl-L-glutamate kinase from *Synechococcus elongatus* strain PCC 7942. *J. Biol. Chem.* **279**:55202–55210.
 32. Meletzus, D., P. Rudnick, N. Doetsch, A. Green, and C. Kennedy. 1998. Characterization of the *glnK-amtB* operon of *Azotobacter vinelandii*. *J. Bacteriol.* **180**:3260–3264.
 33. Michel-Reydellet, N., and P. A. Kaminski. 1999. *Azorhizobium caulinodans* P_{II} and GlnK proteins control nitrogen fixation and ammonia assimilation. *J. Bacteriol.* **181**:2655–2658.
 34. Muro-Pastor, M. L., J. C. Reyes, and F. J. Florencio. 2005. Ammonium assimilation in cyanobacteria. *Photosynth. Res.* **83**:135–150.
 35. Muro-Pastor, M. L., J. C. Reyes, and F. J. Florencio. 2001. Cyanobacteria perceive nitrogen status by sensing intracellular 2-oxoglutarate levels. *J. Biol. Chem.* **276**:38320–38328.
 36. Muse, W. B., and R. A. Bender. 1998. The *nac* (nitrogen assimilation control) gene from *Escherichia coli*. *J. Bacteriol.* **180**:1166–1173.
 37. Osanai, T., S. Sato, S. Tabata, and K. Tanaka. 2005. Identification of PamA as a P_{II}-binding membrane protein important in nitrogen-related and sugar-catabolic gene expression in *Synechocystis* sp. PCC 6803. *J. Biol. Chem.* **280**:34684–34690.
 38. Paz-Yepes, J., E. Flores, and A. Herrero. 2009. Expression and mutational analysis of the *glnB* genomic region in the heterocyst-forming cyanobacterium *Anabaena* sp. strain PCC 7120. *J. Bacteriol.* **191**:2353–2361.
 39. Paz-Yepes, J., E. Flores, and A. Herrero. 2003. Transcriptional effects of the signal transduction protein P_{II} (*glnB* gene product) on NtcA-dependent genes in *Synechococcus* sp. PCC 7942. *FEBS Lett.* **543**:42–46.
 40. Ruiz, D., P. Salinas, M. L. Lopez-Redondo, M. L. Cayuela, A. Marina, and A. Contreras. 2008. Phosphorylation-independent activation of the atypical response regulator NblR. *Microbiology* **154**:3002–3015.
 41. Sauer, J., U. Dirmeier, and K. Forchhammer. 2000. The *Synechococcus* strain PCC 7942 *glnN* product (glutamine synthetase III) helps recovery from prolonged nitrogen chlorosis. *J. Bacteriol.* **182**:5615–5619.
 42. Sugiyama, K., T. Hayakawa, T. Kudo, T. Ito, and T. Yamaya. 2004. Interaction of *N*-acetylglutamate kinase with a P_{II}-like protein in rice. *Plant Cell Physiol.* **45**:1768–1778.
 43. Takatani, N., M. Kobayashi, S. Maeda, and T. Omata. 2006. Regulation of nitrate reductase by non-modifiable derivatives of P_{II} in the cells of *Synechococcus elongatus* strain PCC 7942. *Plant Cell Physiol.* **47**:1182–1186.
 44. Tanigawa, R., M. Shirokane, S. Maeda Si, T. Omata, K. Tanaka, and H. Takahashi. 2002. Transcriptional activation of NtcA-dependent promoters of *Synechococcus* sp. PCC 7942 by 2-oxoglutarate in vitro. *Proc. Natl. Acad. Sci. USA* **99**:4251–4255.
 45. Vázquez-Bermúdez, M. F., A. Herrero, and E. Flores. 2002. 2-Oxoglutarate increases the binding affinity of the NtcA (nitrogen control) transcription factor for the *Synechococcus glnA* promoter. *FEBS Lett.* **512**:71–74.
 46. Vega-Palas, M. A., E. Flores, and A. Herrero. 1992. NtcA, a global nitrogen regulator from the cyanobacterium *Synechococcus* that belongs to the Crp family of bacterial regulators. *Mol. Microbiol.* **6**:1853–1859.
 47. Zhang, Y., E. L. Pohlmann, M. C. Conrad, and G. P. Roberts. 2006. The poor growth of *Rhodospirillum rubrum* mutants lacking P_{II} proteins is due to an excess of glutamine synthetase activity. *Mol. Microbiol.* **61**:497–510.
 48. Zhang, Y., E. L. Pohlmann, and G. P. Roberts. 2005. GlnD is essential for NifA activation, NtrB/NtrC-regulated gene expression, and posttranslational regulation of nitrogenase activity in the photosynthetic, nitrogen-fixing bacterium *Rhodospirillum rubrum*. *J. Bacteriol.* **187**:1254–1265.
 49. Zhang, Y., H. Pu, Q. Wang, S. Cheng, W. Zhao, and J. Zhao. 2007. P_{II} is important in regulation of nitrogen metabolism but not required for heterocyst formation in the cyanobacterium *Anabaena* sp. PCC 7120. *J. Biol. Chem.* **282**:33641–33648.