



CONTRIBUTIONS OF THE ELASTICITY TO THE PRECESSION OF A TWO-LAYER EARTH MODEL

T. BAENAS¹, J. M. FERRÁNDIZ¹, A. ESCAPA^{1,2}, J. GETINO³, AND J. F. NAVARRO¹

¹ Department of Applied Mathematics, University of Alicante, PO Box 99, E-03080 Alicante, Spain; Tomas.Baenas@ua.es

² Department of Aerospace Engineering, University of León, E-24071 León, Spain

³ Department of Applied Mathematics, University of Valladolid, E-47011 Valladolid, Spain

Received 2016 July 28; revised 2016 December 21; accepted 2016 December 22; published 2017 January 25

ABSTRACT

We focus on the updating of a specific contribution to the precession of the equator in longitude, usually named as “second order.” It stems from the crossing of certain terms of the lunisolar gravitational potential. The IAU2006 precession theory assigns it the value of -46.8 mas/cy that was derived for a rigid Earth model. Instead of that model, we consider a two-layer Earth composed of an elastic mantle and a liquid core, working out the problem within the Hamiltonian framework developed by Getino and Ferrándiz. The targeted effect is obtained without further simplifying assumptions through Hori’s canonical perturbation method applied up to the second order of perturbation. On account of using a more realistic Earth model, the revised value of the second-order contribution is significantly changed and reaches -55.29 mas/cy. That variation of the second-order contribution is larger than other contributions included in IAU2006. It must be compensated with an increase of -8.51 mas/cy in the value of the lunisolar first-order component p'_A of the precession of the equator rate, which is derived from the total rate by subtracting the remaining contributions accounted for in IAU2006 precession. The updating of the second-order contribution implies that the p'_A parameter has to be changed, from 5040684.593 to 5040693.104 mas/cy in absence of potential revisions of other contributions. It entails a proportional variation of Earth’s dynamical ellipticity H_d , for which the estimation associated with IAU2006, 0.00327379448 , should be updated to 0.00327380001 , about 1.7 ppm larger.

Key words: celestial mechanics – Earth – methods: analytical – reference systems

1. INTRODUCTION

The main constituent of the precession of Earth’s equator longitude is a linear variation due to the first-order effect of the lunisolar torque. Besides, there are additional contributions to the longitude rate coming from other sources. Most of the contributions considered in the International Astronomical Union (IAU) precession theories IAU1976 and IAU2006 are computed for a rigid Earth model. However, Earth’s internal structure also originates contributions of non-negligible magnitude that cannot be ignored henceforward given the current accuracy of the observations and the stringent accuracy and stability demanded for the reference frames.

The theoretical modeling of the precession has advanced significantly over the past decades following the increasingly accurate monitoring of Earth’s rotation by the very long baseline interferometry (VLBI) technique. In the comprehensive revision of the precession theory carried out by Fukushima (2003) for developing a new set of precession formulae, the speed of the general precession in longitude (p_A) was estimated as $p = 5028.7958 \pm 0.0003$ /cy (where cy stands for Julian century). That uncertainty of $300 \mu\text{as}/\text{cy}$, equivalent to a relative error of about 60 ppb (parts per billion), is of the same order of magnitude as the targets of accuracy and stability of the reference frames established by the Global Geodetic Observing System (GGOS) of the International Association of Geodesy (IAG), namely, 1 mm in position and 1 mm yr⁻¹ in stability (Plag & Pearlman 2009). Those goals, which correspond to geocentric angles and angular velocities near $30 \mu\text{as}$ and $300 \mu\text{as}/\text{cy}$, were adopted by the IAU/IAG Joint Working Group on Theory of Earth Rotation in 2013 (see, e.g., Ferrándiz & Gross 2015).

In 2006 (Hilton et al. 2006) the General Assembly of the IAU approved Resolution 1, which endorsed the P03 model by

Capitaine et al. (2003, 2005) as the new precession theory IAU2006 that superseded Lieske’s (1977) IAU1976. IAU2006 uses fifth-degree polynomials to model the precession variables of the equator and the ecliptic. This kind of polynomial expansion is the most used in precession theories (see, however, Vondrák et al. 2011). The rate of the longitude of the node of the equator adopted in IAU2006 is 5038.481507 /cy. It consists of several components. By far, the largest of them, denoted here⁴ as p'_A , is due to the *first-order* effect of the lunisolar torque acting on the oblate Earth, close to 5040 /cy—namely, the linear approximation of the solution for MacCullagh’s term of the lunisolar potential. It is given by a simple formula depending on Earth’s angular velocity ω_E and dynamical ellipticity

$$H_d = \frac{2C - A - B}{2C}, \quad (1)$$

the masses of the Sun, Moon, and Earth, and some orbital quantities (Kinoshita & Souchay 1990, Equation (8.7))

$$p'_A = 3H_d \left[\left(\frac{m_M}{m_M + m_E} \right) \left(\frac{n_M^2}{\omega_E} \right) M_0 + \left(\frac{m_S}{m_S + m_E + m_M} \right) \left(\frac{n_S^2}{\omega_E} \right) S_0 \right] \cos \epsilon_A. \quad (2)$$

Each theory arrives at different determinations of H_d (usually without specifying estimated errors) depending on the precession rate inferred from observations and on the remaining

⁴ We follow the notation introduced by Souchay & Kinoshita (1996). Notice that this parameter is the same as that denoted $(r_0)_1$ by Capitaine et al. (2003), while, e.g., Fukushima (2003) uses P' .

Table 1
Different Determinations of the Dynamical Ellipticity

H_d Value	Uncertainty	Source
0.0032737 634	...	Williams (1994)
0.0032737 548	...	REN2000, Souchay et al. (1999)
0.0032737 674	...	RDAN97, Roosbeek & Dehant (1998)
0.0032737 949	12×10^{-10}	MHB2000, Mathews et al. (2002)
0.0032737 804	3×10^{-10}	Fukushima (2003)
0.0032737 9448	...	P03, Capitaine et al. (2003)
0.0032737 96	...	FNEG04, Ferrándiz et al. (2004)
0.0032738 0001	...	This work

Note. Ellipsis points means that the uncertainty is not provided in the source along with the value. The H_d value in MHB2000 is 12 ppm larger than the corresponding parameter in its underlying rigid Earth theory REN2000.

theoretical contributions accounted for. A selection of values and their sources are displayed in Table 1.

Let us recall that the dynamical ellipticity is a main parameter that also factorizes all the first-order nutation terms. Therefore, any change of the adopted H_d requires correcting the nutation amplitudes by computing the so-called *indirect effects* (Escapa et al. 2016).

Besides the main lunisolar first-order rate of the equator longitude, the IAU2006 precession theory included many effects of noticeably smaller magnitude than p'_A , which are listed in Table 3 of Capitaine et al. (2003)—with an update of a single component in Capitaine et al. (2005). Most of them were derived before the adoption of the MHB2000 theory (Mathews et al. 2002) as the IAU2000 nutation theory, and their values are practically the same as given by, e.g., Kinoshita & Souchay (1990), Williams (1994, 1995), or Souchay & Kinoshita (1996), despite the simplifying assumptions made in some of the former theoretical models. The contributions included in IAU2006 (Capitaine et al. 2003, 2005) are listed in Table 2. All of them but the last one are derived for a rigid Earth model. The largest contribution is 400 parts per million (ppm) of p'_A , due to the geodesic precession.

However, new contributions to the precession rate have been found since the P03 derivation (e.g., Liu & Capitaine 2015). For this paper’s sake, we emphasize the generalization of the IAU2006 (REN2000) *lunisolar second-order* effects to a *non-rigid Earth* (Ferrándiz et al. 2004). The contribution to the longitude rate that IAU2006 names as *lunisolar second-order effect* (split in one part for the Moon and another for the Sun, labeled “a” and “b” respectively) comes from the main (zonal) terms of the second order of perturbation solution of the rigid Earth Hamiltonian (quadratic in H_d); it was derived by Kinoshita & Souchay (1990) and remained unchanged in REN2000. Ferrándiz et al. (2004) showed that, unlike commonly believed, the presence of a fluid core affects the precessional motion of Earth by a non-negligible amount, reaching -5 ppm of the value of the precession constant when a Poincaré-like Earth model (rigid mantle with fluid core) is assumed. An explanation can be found in the amplification of the Opolzer terms caused by the resonance associated with the core.

This research consists in a further extension of those lunisolar second-order effects for a two-layer Earth model (Ferrándiz et al. 2004) by including the contributions of the elasticity to the precession in longitude. Some partial and preliminary computations of those contributions were derived by Ferrándiz et al. (2007), but only the numerical results were presented, due to the nature of that concise proceedings report.

Table 2
Contributions to Precession Rate Included in IAU2006

Contribution	Value (mas/cy)	Source
First-order lunisolar (p'_A) ^a	5040684.593	Capitaine et al. (2005)
Second-order lunisolar	-46.780	Williams (1994)
Lunisolar J_4	+2.600	Williams (1994)
First-order planetary	+31.367	Capitaine et al. (2003)
J_2 -tilt	-269.430	Williams (1994)
Geodesic precession	-1919.883	Capitaine et al. (2003)
MHB2000’s nonlinear effect	-0.960	Mathews (unpublished) ^b

Notes.

^a Not provided explicitly, but obtained by subtraction.

^b Recomputation of the Mathews et al. (2002) value cited in Capitaine et al. (2005).

Note that these terms are of a different nature from those related with the redistribution tidal potential, and currently they cannot be provided by any theory but the Hamiltonian one presented in this work.⁵ Specifically, we evaluate the second-order contributions to the precession in longitude through Hori’s canonical method for a two-layer Earth model composed of an elastic mantle and a fluid core. They arise as a consequence of the crossing of first-order nutation terms due to the (2, 0) harmonic of the gravitational potential and the tidal rotational kinetic energy. We perform a rigorous computation and a detailed analysis of those contributions. They involve the indirect terms and also the direct ones, which must be incorporated when fully taking into account the influence of the elasticity of the model and nowadays accuracy requirements. The indirect terms reflect the effect of the core through the nearly diurnal free wobble (NDFW), i.e., they can be associated with the free core nutation (FCN), whereas the direct ones stem from an additional contribution to the kinetic energy of the system entirely due to the fluid core.

The structure of the paper is as follows. Section 2 contains an outline of Hori’s canonical method that is applied to the Hamiltonian of the non-rigid Earth model considered in this work. It provides analytical expressions for the second-order contributions to the precession rate. Those expressions are split according to their different origins, which is possible thanks to the analytical approach used in their derivation. Section 3 contains a numerical evaluation of new contributions, comparing them with other relevant second-order corrections appearing in the literature. Finally, in Section 4, we draw the main conclusions of this investigation. The paper is completed with two appendices including some auxiliary material.

2. DERIVATION OF THE NEW SECOND-ORDER CONTRIBUTION TO PRECESSION

The solution that contains the new contribution to the precession, due to the previously mentioned non-rigid Earth structure, is worked out using the Hamiltonian approach developed by Getino & Ferrándiz (1991, 1997, 2001) for the non-rigid Earth. Some of its main features are as follows: (1) it allows a consistent treatment, since the first- and second-order lunisolar perturbations are derived from a sole Hamiltonian, with only a set of basic Earth parameters (BEPs) and a sole

⁵ The nonlinear effects presented in Mathews et al. (2002) are related to some part of the redistribution tidal potential (Lambert & Mathews 2006), i.e., to the effects of the variation of the geopotential due to tidal deformations.

tide-generating expansion; (2) the crossing effects are rigorously obtained by Hori's (1966) perturbation method; and (3) the procedure is systematic and can be performed by computer algebra; therefore, no group of terms must be neglected in advance to simplify the analytical derivations (unlike what happens usually in less elaborated approaches dealing with the equations of motion).

2.1. Outline of the Method

Next, the procedure followed in the derivation is sketched. The method is very close to that used in Ferrándiz et al. (2004) to derive an analogous contribution to the precession in the case of rigid mantle, although the consideration of the elasticity introduces noticeable complexities and differences. The original Hamiltonian, \mathcal{H} , is divided into perturbation orders and then transformed into a new Hamiltonian, \mathcal{K} , easier to solve, by calculating the generating function, \mathcal{W} , of a certain canonical transformation close to the identity. Finally, the precession rate perturbation is computed by taking a certain partial derivative of the generating function.

More precisely, we apply a perturbation method based on Lie series up to the second order, which is equivalent to the Hori (1966) canonical method, combined with an averaging requirement. The starting ("old") and transformed ("new") Hamiltonians are each divided into an unperturbed or zeroth-order term and perturbation terms of first and second orders, namely,

$$\begin{aligned}\mathcal{H} &= \mathcal{H}_0 + \mathcal{H}_1 + \mathcal{H}_2, \\ \mathcal{K} &= \mathcal{K}_0 + \mathcal{K}_1 + \mathcal{K}_2,\end{aligned}\quad (3)$$

with

$$\mathcal{K}_0 = \mathcal{H}_0, \mathcal{K}_1 = \mathcal{H}_{1,\text{sec}}, \mathcal{K}_2 = \frac{1}{2} \{\mathcal{H}_1 + \mathcal{K}_1, \mathcal{W}_1\}_{\text{sec}} + \mathcal{H}_{2,\text{sec}}.\quad (4)$$

Here $\{--, --\}$ denotes the Poisson bracket and subscript "sec" the secular part (or time average) of a function. The components of the generating function $\mathcal{W} = \mathcal{W}_1 + \mathcal{W}_2$ satisfy the equations of the method

$$\{\mathcal{H}_0, \mathcal{W}_1\} = \mathcal{K}_1 - \mathcal{H}_1 \equiv \mathcal{A}_1, \quad (5)$$

$$\{\mathcal{H}_0, \mathcal{W}_2\} = \mathcal{K}_2 - \mathcal{H}_2 - \frac{1}{2} \{\mathcal{H}_1 + \mathcal{K}_1, \mathcal{W}_1\} \equiv \mathcal{A}_2. \quad (6)$$

Their solutions \mathcal{W}_i can be expressed as

$$\mathcal{W}_i = - \int_{\text{UP}} \mathcal{A}_i dt, \quad (7)$$

where subscript UP means that the integral is computed along the trajectories of the unperturbed problem (\mathcal{H}_0). It is understood that functions on the right-hand side of Equations (4)–(7) are expressed in the *new* canonical variables (q, p) on which \mathcal{K} depends by literally substituting the old by the new ones (Ferraz-Mello 2007).

Notice that all the terms \mathcal{K}_i of the new Hamiltonian are defined in order to be free from periodic components, whereas functions \mathcal{A}_i and \mathcal{W}_i average to zero. The method provides an asymptotic approximate solution to the original variables as the sum of periodic perturbations superposed to a secular evolution. The periodic perturbations are derived from the so-called perturbation equations and are not needed for this

paper's goals. The secular motion is the solution of the transformed Hamiltonian, which is free of periodic terms up to the second order of perturbation. Therefore, the equations derived from \mathcal{K} give rise to the secular motion of the canonical variables, which is the relevant one to study precession. More precisely, if the secular solution is assumed to be known up to the first order of perturbation, the additive contribution to the rate of a canonical coordinate q_j conjugate to p_j (e.g., the incremental precessional rate in our problem) is obtained by means of

$$\delta \left(\frac{dq_j}{dt} \right) = \{q_j, \mathcal{K}_2\} = \frac{\partial \mathcal{K}_2}{\partial p_j}. \quad (8)$$

2.2. Arrangement of the Hamiltonian of the Two-layer Earth

Let us consider a two-layer Earth composed of a mantle and a fluid core and subject to elastic deformations. A Hamiltonian formulation of that system was used by Getino & Ferrándiz (2001) in their GF2000 nutation solution, by means of an Andoyer-like set of canonical variables. The first step is casting the Hamiltonian into the form given by Equation (3). Term \mathcal{H}_2 gathers some small perturbations, e.g., the lunisolar potential accounting for the rigid Earth's triaxiality (Kinoshita 1977) or corresponding to its higher-degree harmonics (Folgueira et al. 1998, 1999, 2001), the planetary rigid Earth potential (Kinoshita & Souchay 1990), and non-rigid Earth terms like the additional potential due to the deformations induced by the rotational potential or lunisolar attraction (Getino & Ferrándiz 1995). However, the \mathcal{H}_2 contribution to precession, independently of its magnitude, only depends on its secular part \mathcal{K}_2 ; as we are concerned here only with the effect arising from the interaction among \mathcal{H}_1 , \mathcal{K}_1 , and \mathcal{W}_1 , according to Equations (4) and (8), the expression of \mathcal{H}_2 is irrelevant for our purpose and can be ignored without altering the results.

We can start from the simplified Hamiltonian

$$\mathcal{H} = \mathcal{H}_0 + \mathcal{H}_1, \quad (9)$$

which replaces Equation (3), where

$$\mathcal{H}_0 = T_0, \quad \mathcal{H}_1 = E + V_{\text{sec}} + V_{\text{per}} + T_r. \quad (10)$$

In these expressions, T_0 is the rotational kinetic energy of Earth in a certain unperturbed state prior to deformation considered in the definition of the canonical variables, E is the term introduced by Kinoshita (1977) for referring the problem to the ecliptic of date, and V_{sec} and V_{per} are, respectively, the secular and periodic part of the (2, 0) harmonic (the J_2 term) of the gravitational potential. Finally, T_r is the increment of the rotational kinetic energy arising from the change of the inertia tensor caused by Earth's tidal deformation. This term is responsible of the contributions to the precession in longitude worked out in this investigation, and it is absent in Poincaré's Earth model (Ferrándiz et al. 2004).

With this decomposition of \mathcal{H} , the secular perturbation terms of Hamiltonian \mathcal{K} read as

$$\mathcal{K}_1 = E + V_{\text{sec}}, \quad \mathcal{K}_2 = \frac{1}{2} \{\mathcal{H}_1 + \mathcal{K}_1, \mathcal{W}_1\}_{\text{sec}}, \quad (11)$$

where

$$\mathcal{W}_1 = \int_{\text{UP}} (V_{\text{per}} + T_r) dt. \quad (12)$$

where subscripts P and t refer to V_{per} and T_t parts in Equation (12), respectively, and its components are

$$\begin{aligned} \mathcal{W}_P &= \sum_q \sum_{j \neq 0} k'_q \frac{1}{2} (3 \cos^2 \sigma - 1) \frac{B_{j,q}(I)}{n_j} \sin \Theta_j \\ &\quad - \frac{1}{2} \sin 2\sigma \sum_q \sum_{j,\rho=\pm 1} k'_p C_{j,\rho,q}(I) F_{j,\rho}^{1a} \sin(\mu - \rho \Theta_j) \\ &\quad - \sin \sigma_c \frac{M_c}{M} \sum_q \sum_{j,\rho=\pm 1} k'_p C_{j,\rho,q}(I) F_{j,\rho}^{2a} \sin(h_{j,\rho} + \nu_c), \\ \mathcal{W}_t &= M \sin \sigma \sum_q \sum_{j,\rho=\pm 1} C_{j,\rho,q}(\tilde{I}) D_{j,\rho,q}^{1a} \sin(\tilde{h}_{j,\rho} - \nu) \\ &\quad + M_c \sin \sigma_c \sum_q \sum_{j,\rho=\pm 1} C_{j,\rho,q}(\tilde{I}) D_{j,\rho,q}^{2a} \sin(\tilde{h}_{j,\rho} + \nu_c). \end{aligned} \quad (20)$$

In the former equations $n_j = d\Theta_j/dt$ are the mean motions of the long-period nutation arguments, and functions $F_{j,\rho}^{ka}$, $G_{j,\rho}^{ka}$ and $D_{j,\rho,q}^{ka} = k_q^m (F_{j,\rho}^{ka} + G_{j,\rho}^{ka}) - k_q^c G_{j,\rho}^{ka}$ ($k = 1, 2$) are the coefficients used in Getino & Ferrándiz (2001), which are displayed in Appendix A. It can be seen there that coefficients F and G depend on some dimensionless parameters (ellipticities and ratio of inertia moments of Earth and core) and some angular velocities, namely, ω_E , the frequencies of the free two-layer Earth wobbles CW and FCN (Chandler Wobble and FCN) and the frequencies of the diurnal band terms in those expansions, $n_{h_{j,\rho}} = dh_{j,\rho}/dt = \omega_E - \rho n_j$.

The secular second-order term \mathcal{K}_2 (Equation (16)) can be expressed as

$$\begin{aligned} \mathcal{K}_2 &= \frac{1}{2} \{V_{\text{per}} + T_t, \mathcal{W}_P + \mathcal{W}_t\}_{\text{sec}} \\ &= \frac{1}{2} \{V_{\text{per}}, \mathcal{W}_P\}_{\text{sec}} + \frac{1}{2} \{V_{\text{per}}, \mathcal{W}_t\}_{\text{sec}} \\ &\quad + \frac{1}{2} \{T_t, \mathcal{W}_P\}_{\text{sec}} + \frac{1}{2} \{T_t, \mathcal{W}_t\}_{\text{sec}}. \end{aligned} \quad (21)$$

Therefore, the first step is the calculation of the four Poisson brackets, and then of their secular parts. Notice that in the absence of elasticity, i.e., in Poincaré's Earth model treated in Ferrándiz et al. (2004), only the first bracket was necessary to obtain \mathcal{K}_2 . The main guidelines for computing these Poisson brackets are given in Appendix B. Neglecting the terms of first order in σ and σ_c and higher, we get after some algebra

$$\begin{aligned} \{V_{\text{per}}, \mathcal{W}_P\} &= -\frac{1}{M \sin I} \sum_{p,q,i,j \neq 0} \sum_{\tau,\rho=\pm 1} \frac{k'_p k'_q}{n_j} B_{i,p} \frac{dB_{j,q}}{dI} m_{5i} \\ &\quad \times \cos(\Theta_i - \Theta_j) \\ &\quad - \frac{1}{M} \sum_{p,q} \sum_{i,j} \sum_{\tau,\rho=\pm 1} k'_p k'_q C_{i,\tau,p} C_{j,\rho,q} F_{j,\rho}^{1a} \\ &\quad \times \cos(\tau \Theta_i - \rho \Theta_j). \end{aligned} \quad (22)$$

The first group of terms (coming from zonal–zonal crossing as the presence of the B_i coefficients reveals) is analytically the same for rigid one-layer or two-layer Earth models, either rigid (Poincaré's) or elastic. The second group, with sectorial–sectorial crossing origin shown by coefficients C_i , is formally kept only inside the class of two-layer models. However, we can recover the rigid Earth expression by replacing F_i^{1a} functions with $1/(n_\mu - \tau n_i)$, where $n_\mu = \omega_E C/A$.

The remaining Poisson brackets in Equation (21), also truncated at $O(\sigma^0, \sigma_c^0)$, are

$$\begin{aligned} \{V_{\text{per}}, \mathcal{W}_t\} &= \sum_{p,q} \sum_{i,j} \sum_{\tau,\rho=\pm 1} k'_p D_{j,\rho,q}^{1a} C_{i,\tau,p} \tilde{C}_{j,\rho,q} \\ &\quad \times \cos(\tau \Theta_i - \rho \tilde{\Theta}_j), \\ \{T_t, \mathcal{W}_P\} &= \sum_{p,q} \sum_{i,j} \sum_{\tau,\rho=\pm 1} k_p^m k'_q F_{j,\rho}^{1a} \tilde{C}_{i,\tau,p} C_{j,\rho,q} \\ &\quad \times \cos(\tau \tilde{\Theta}_i - \rho \Theta_j) \\ &\quad + \frac{M_c}{M} \sum_{p,q} \sum_{i,j} \sum_{\tau,\rho=\pm 1} k'_p (k_p^c - k_p^m) F_{j,\rho}^{2a} \\ &\quad \times \tilde{C}_{i,\tau,p} C_{j,\rho,q} \cos(\tau \tilde{\Theta}_i - \rho \Theta_j), \\ \{T_t, \mathcal{W}_t\} &= -M \sum_{p,q} \sum_{i,j} \sum_{\tau,\rho=\pm 1} k_p^m D_{j,\rho,q}^{1a} \\ &\quad \times \tilde{C}_{i,\tau,p} \tilde{C}_{j,\rho,q} \cos(\tau \tilde{\Theta}_i - \rho \tilde{\Theta}_j) \\ &\quad - M_c \sum_{p,q} \sum_{i,j} \sum_{\tau,\rho=\pm 1} (k_p^c - k_p^m) D_{j,\rho,q}^{2a} \\ &\quad \times \tilde{C}_{i,\tau,p} \tilde{C}_{j,\rho,q} \cos(\tau \tilde{\Theta}_i - \rho \tilde{\Theta}_j). \end{aligned} \quad (23)$$

These expressions are exclusively due to the elastic yielding of the two-layer Earth model (T_t term), i.e., they are not present in Poincaré's Earth model (Ferrándiz et al. 2004). They have a double origin. On the one hand, some of them depend indirectly on the core, just through the functions $F_{j,\rho}^{ka}$ and $D_{j,\rho,q}^{ka}$, which are sensitive to the frequency of the NDFW. On the other hand, some terms depend directly on the modulus of the core angular momentum M_c .

2.4. Analytical Solution of the Second-order Precession

The secular part of the Poisson brackets that gives \mathcal{K}_2 can be obtained straightforwardly, by simply requiring that the angular arguments in Equations (22) and (23) vanish. Omitting the tildes for brevity, in the case of arguments $\Theta_i - \Theta_j = 0$ (zonal–zonal) there is only a possibility, which is satisfied when $i = j$; the contributions to precession of such origin are identified in the following with the superscript 00. In the most frequent case of having the argument $\tau \Theta_i - \rho \Theta_j = 0$, we have two possibilities: first, $i = j = 0$ and unrestricted τ, ρ ($= \pm 1$), identified here with the superscript 10; and second, $i = j \neq 0$ and $\tau = \rho$, noted with superscript 11.

The contributions to the precession rate in longitude derived from the resulting secular part of the first Poisson bracket given in Equation (22) are readily obtained from Equation (15) and

can be grouped in

$$\begin{aligned}
\delta p_p^{00} &= -\frac{1}{2} \sum_{p,q} \sum_{i \neq 0} \frac{1}{\sin^3 I} \frac{k_p k_q}{n_i} m_{Si} \times \left(\sin IB_{i,p} \frac{d^2 B_{i,q}}{dI^2} \right. \\
&\quad \left. + \sin I \frac{dB_{i,p}}{dI} \frac{dB_{i,q}}{dI} - \cos IB_{i,p} \frac{dB_{i,q}}{dI} \right), \\
\delta p_p^{10} &= -\frac{1}{2} \frac{1}{\sin I} \sum_{p,q,\tau,\rho=\pm 1} k_p k_q F_{0,\rho}^{1a} \\
&\quad \times \left(\frac{dC_{0,\tau,p}}{dI} C_{0,\rho,q} + C_{0,\tau,p} \frac{dC_{0,\rho,q}}{dI} \right), \\
\delta p_p^{11} &= -\frac{1}{2} \frac{1}{\sin I} \sum_{p,q} \sum_{i \neq 0} \sum_{\tau=\pm 1} k_p k_q F_{i,\tau}^{1a} \\
&\quad \times \left(\frac{dC_{i,\tau,p}}{dI} C_{i,\tau,q} + C_{i,\tau,p} \frac{dC_{i,\tau,q}}{dI} \right), \tag{24}
\end{aligned}$$

$k_p = k'_p/M = k'_p/(C\omega_E)$ being the usual Kinoshita's (1977) constants of Moon and Sun, proportional to the dynamical ellipticity H_d .

Let us remark that the δp_p^{00} term (arising from zonal–zonal crossing) corresponds to the second-order lunisolar effect first computed by Kinoshita & Souchay (1990), whose value has remained unchanged since then and has been adopted particularly in REN2000 (Souchay et al. 1999), Williams (1994), and Capitaine et al. (2003). Notice that the correspondence is not exact, because the original derivation of it was performed jointly with the J_2 -tilt effect and using a few terms obtained from an elliptical approximation of the Moon and Sun orbits (instead of their own accurate orbital coefficients $A_{i,p}^{(j)}$). These three terms were first calculated by Ferrándiz et al. (2004), who used the notation dp_0 , dp_{10} , dp_{11} .

The remaining three Poisson brackets given in Equation (23) provide the contributions steaming from the elasticity of the model and have been derived for the first time in this work, being one of its most important results. They can be arranged as

$$\begin{aligned}
\delta p_{t,I}^{10} &= \frac{1}{2} \frac{1}{\sin I} \sum_{p,q,\tau,\rho=\pm 1} k_p [k_q^m (F_{0,\rho}^{1a} + F_{0,\tau}^{1a}) \\
&\quad + (k_q^m - k_q^c) G_{0,\rho}^{1a}] \frac{dC_{0,\tau,p}}{dI} \tilde{C}_{0,\rho,q}, \\
\delta p_{t,I}^{11} &= \frac{1}{2} \frac{1}{\sin I} \sum_{p,q} \sum_{i \neq 0} \sum_{\tau=\pm 1} k_p [k_q^m (F_{i,\tau}^{1a} + F_{i,\tau}^{1a}) \\
&\quad + (k_q^m - k_q^c) G_{i,\tau}^{1a}] \frac{dC_{i,\tau,p}}{dI} \tilde{C}_{i,\tau,q}, \\
\delta p_{t,D}^{10} &= \frac{1}{2} \frac{1}{\sin I} \frac{M_c}{M} \sum_{p,q,\tau,\rho=\pm 1} k_q (k_p^c - k_p^m) \\
&\quad \times F_{0,\rho}^{2a} \frac{dC_{0,\rho,q}}{dI} \tilde{C}_{0,\tau,p}, \\
\delta p_{t,D}^{11} &= \frac{1}{2} \frac{1}{\sin I} \frac{M_c}{M} \sum_{p,q} \sum_{i \neq 0} \sum_{\tau=\pm 1} k_q (k_p^c - k_p^m) \\
&\quad \times F_{i,\tau}^{2a} \frac{dC_{i,\tau,q}}{dI} \tilde{C}_{i,\tau,p}. \tag{25}
\end{aligned}$$

Superscripts have the same meaning as before and, regarding subscripts, terms $\delta p_{t,I}$ gather the indirect effects of the fluid core, while $\delta p_{t,D}$ contain the direct ones. The latter come from

the terms of the generating function that depend on the canonical variables of the core, and are proportional to M_c/M . Note that the $\{T_i, \mathcal{W}_i\}$ bracket does not generate any contribution, since it does not depend on the canonical variable I but on the function of time \tilde{I} , so that all the direct effects of the core arise from the second term of $\{T_i, \mathcal{W}_p\}$. These effects have not been considered in the literature so far.

The direct and indirect contributions may be added up to a single analytical expression taking into account the following relationship between F_i and G_i functions:

$$\frac{M_c}{M} F_{i,\tau}^{2a} = -G_{i,\tau}^{1a}, \tag{26}$$

which can be obtained from the formulae included in Appendix A. The overall second-order contribution of the elasticity to the precession rate in longitude can thus be written as

$$\begin{aligned}
\delta p_t^{10} &= \frac{1}{2} \frac{1}{\sin I} \sum_{p,q,\tau,\rho=\pm 1} k_p [k_q^m (F_{0,\rho}^{1a} + F_{0,\tau}^{1a}) \\
&\quad + 2(k_q^m - k_q^c) G_{0,\rho}^{1a}] \frac{dC_{0,\tau,p}}{dI} \tilde{C}_{0,\rho,q} \\
\delta p_t^{11} &= \frac{1}{2} \frac{1}{\sin I} \sum_{p,q} \sum_{i \neq 0} \sum_{\tau=\pm 1} k_p [k_q^m (F_{i,\tau}^{1a} + F_{i,\tau}^{1a}) \\
&\quad + 2(k_q^m - k_q^c) G_{i,\tau}^{1a}] \frac{dC_{i,\tau,p}}{dI} \tilde{C}_{i,\tau,q}. \tag{27}
\end{aligned}$$

These expressions supersede Equation (25). Notice that the indirect effects can be isolated by changing the factor of 2 that multiplies $(k_q^m - k_q^c)$ in Equation (27) to a factor of 1.

2.4.1. Final Analytical Expression of the Second-order Contribution

In conclusion, the total effect $\delta p|_{\kappa_2}$ can be represented by means of a sum of three groups of terms in the form

$$\delta p \equiv \delta p|_{\kappa_2} = \delta p_p^{00} + (\delta p_p^{10} + \delta p_p^{11}) + (\delta p_t^{10} + \delta p_t^{11}). \tag{28}$$

Each group of terms is associated with certain improvement of the Earth model assumed in the derivation of the second-order effect. A (one-layer) rigid model gives rise to δp_p^{00} only; a two-layer model with rigid mantle (Poincaré's) adds the terms $\delta p_p^{10} + \delta p_p^{11}$; finally, $\delta p_t^{10} + \delta p_t^{11}$ is added in the elastic two-layer Earth case (including both direct and indirect core effects).

3. NUMERICAL RESULTS AND DISCUSSION

The analytical expressions that provide the different components of the lunisolar second-order effect have been evaluated numerically. The values assigned to the parameters are displayed in Table 3, together with the sources of each. Parameters specific to the Hamiltonian formulation have been taken from the Getino & Ferrándiz (2001) two-layer nutation series. Using the values resulting from the fit of a three-layer solution, or a different set of parameters, produces differences below the uncertainty threshold of the determination of precession, given the small magnitude of the effect and the small variations of parameters between both models. At any rate, the analytical nature of our results allows the evaluation of those contributions using different rheological models in a straightforward manner.

Table 3
Numerical Values of the Parameters Employed in the δp Evaluation

Parameter	Value	Uncertainty	Source
ω_E	$7.292115 \times 10^{-5} \text{ rad s}^{-1}$...	Luzum et al. (2011)
$I_0(\text{J2000})$	$-84381.406 \text{ arcsec}$	0.001 arcsec	Luzum et al. (2011)
$d\theta/dt$	$1.00273781191135448 \text{ rev UT1-day}^{-1}$...	Luzum et al. (2011)
H_d	3273795×10^{-9}	1×10^{-9}	IERS Conventions Petit & Luzum (2010)
k_{Moon}	$7567.870647 \text{ arcsec cy}^{-1}$...	Getino & Ferrándiz (2001)
k_{Sun}	$3474.613747 \text{ arcsec cy}^{-1}$...	Getino & Ferrándiz (2001)
k_{Moon}^m	$1906.852345 \text{ arcsec cy}^{-1}$...	Getino & Ferrándiz (2001)
k_{Sun}^m	$875.487397 \text{ arcsec cy}^{-1}$...	Getino & Ferrándiz (2001)
k_{Moon}^c	$4514.468392 \text{ arcsec cy}^{-1}$...	Getino & Ferrándiz (2001)
k_{Sun}^c	$2072.714329 \text{ arcsec cy}^{-1}$...	Getino & Ferrándiz (2001)
P_{CW}	430.9 msd	0.7 msd	Nastula & Gross (2015)
P_{FCN}	430.00 msd	0.01 msd	Krásná et al. (2013)
A_c/A_m	0.1284	...	Dziewonski & Anderson (1981)
Orbital coefficients	Getino et al. (2010), Table 8	...	Kinoshita & Souchay (1990)

Note. The ellipsis points signify that the uncertainty is not provided in the source along with the value. The H_d value and uncertainty in IERS Conventions (Petit & Luzum 2010) are consistent with the IAU2006/2000 precession-nutation model (Capitaine et al. 2003).

Table 4 contains the numerical values of the different individual components of the precession rate contribution derived in Section 2.4 for the considered elastic two-layer model. The three central rows correspond to the different kinds of groups identified with superscripts according to their origin, 00 (zonal) and 10, 11 (sectorial), whose constituents were distinguished with subscripts p , t . The last two groups arise from the components of the Hamiltonian and the generating function, which give rise to the *Oppolzer* nutation terms at first order. These kinds of terms were introduced in the precession theory by Ferrándiz et al. (2004). Their magnitude is strongly dependent on the Earth model features, since they are amplified by the resonance induced by the fluid core as happens with *Oppolzer* terms in nutations.

The first column of figures (column (2)) provides the first three terms, δp_p^{00} , δp_p^{10} , and δp_p^{11} , of the total contribution shown in Equation (28). Let us recall that their sum would contain the total second-order effect if we removed elasticity from the hypothesis; the column is labeled as *Poincaré-rigid mantle + fluid core* for that reason. Besides, the first term of column (2), δp_p^{00} , is common to rigid and non-rigid Earth solutions; it is obtained from the terms of the potential and of the first-order generating function that produce the *Poisson* nutation terms at first order (Kinoshita 1977). The next two columns display the additional contributions stemming from elasticity, named δp_t^{10} and δp_t^{11} in Equation (28), separating in each the indirect (column (3)) and direct (column (4)) effects of the core, defined in Equation (25)— $\delta p_{t,I}^{1x}$ and $\delta p_{t,D}^{1x}$, respectively. The last column sums the contributions of each kind, and the last row is the total by columns.

Let us remark that all the constituents of the total precession rate (except one, the direct core effect $\delta p_{t,D}^{11}$) exceed 0.3 mas/cy. We have taken here that value as proxy of the current accuracy needs because it is the GGOS stability goal and also the estimation of the speed of the general precession in longitude made by Fukushima (2003). It can also be seen that the direct effects of the core are smaller than but reach the same order of magnitude as the indirect ones. Looking at the last column, which gathers the total contribution of each group, we can consider that its first cell, -45.51 mas/cy , has been somehow accounted for in precession theory, since it corresponds to an

Table 4
Second-order Lunisolar Contributions to Precession Rate in Longitude (mas/cy) for the Elastic Two-layer Earth Model Considered in This Work

Additive contributions	Poincaré's Model		Elasticity		TOTAL
	Rigid Mantle + Fluid Core	Core Effects			
		Indirect	Direct		
δp^{00}	-45.51				-45.51
δp^{10}	-22.43	+9.45	+3.80		-9.17
δp^{11}	-1.26	+0.48	+0.17		-0.61
δp	-69.20	+9.94	+3.97		-55.29
			Uncertainty ^a :		±0.01

Note.

^a See the text for discussion.

effect (δp^{00} term) common to rigid and non-rigid Earth models, as noted in the previous paragraph. However, the contributions in the other two cells (which gather the total δp^{10} and δp^{11} effects of sectorial origin) have never been considered in IAU precession theories so far. These last effects amount to -9.78 mas/cy , about 20% of the zonal, rigid effect accounted for so far. Let us notice too that in the absence of elasticity the sectorial terms would produce a much larger effect that reaches -23.67 mas/cy ; therefore, elasticity reduces the impact of the NDFW resonance in the precession rate in the same way that it happens in nutation theory.

To ascertain the uncertainty of our evaluation to -55.29 mas/cy of the second-order effect is not simple, and any estimation of it has its own drawbacks. Anyway, as it follows from an approximate analytical solution of a given model, the discussion can be started by grouping the sources of inaccuracy into two classes: one relative to the goodness of the approximation provided by that mathematical solution, and the other associated with the inaccuracies of the values of the astronomical and geophysical parameters used in the numerical evaluation of the analytical solution. The first can be ignored since the error of our solution is of the third order

Table 5
Comparison of Second-Order Corrections (δp in mas/cy)

Solution	IAU2006	RDAN97	FNEG04	This Work
Earth model	Rigid	Rigid	Poincaré	Elastic two-layer
δp^{00}	-46.78	-45.40	-45.36	-45.51
δp^{10}			-22.59	-9.17
δp^{11}			-1.30	-0.61
Total δp	-46.78	-45.40	-69.25	-55.29
Deviation of δp from IAU2006 value	0	+1.38	-22.47	-8.51
Associated δH_d	0	-90×10^{-11}	1459×10^{-11}	553×10^{-11}
$H_{d,IAU2006} + \delta H_d$	0.00327379448	0.00327379358	0.00327380907	0.00327380001
Relative H_d change (ppm)	0	-0.3	+4.5	+1.7

of perturbation, and the dimensionless perturbation order is $O((V_{\text{sec}} + V_{\text{per}})/T_0) \approx 10^{-7}$ (see, e.g., Getino & Ferrándiz 1995).

The second kind of error is more intricate, since the uncertainties of many parameters are not provided in the original references and seem to be unavailable in the literature. It forces us to make complementary assumptions.

For instance, the first component δp^{00} is independent of the Earth model as noticed before; although the notation may not be revealing enough, it only depends on a few astronomical parameters, basically those present in Equation (2), all of which are known with a relative uncertainty better than 10^{-6} . Therefore, all four significant figures of its value displayed in Table 4 can be taken as exact if this reasoning is considered plausible.

The two remaining components, δp^{10} and δp^{11} , depend also on additional geophysical parameters, mainly the periods of the Chandler wobble and the retrograde FCN, P_{CW} and P_{FCN} —which have been widely studied—and the ratio $r_{\text{cm}} = A_c/A_m$. If we adopt for P_{CW} and P_{FCN} the values and uncertainties provided in Table 3, the resulting δp uncertainty evaluates to ± 0.002 mas/cy, and those parameters would have no effect on the displayed figures.

The r_{cm} ratio is the most critical parameter. Although there is a high degree of agreement among different determinations of it, we found no recent, reliable, explicit evaluation of its uncertainty and had to resort to indirect estimations. For instance, we could consider the recent determinations of the moments of inertia of Earth’s layers appearing in Table 9 of Chen et al. (2015), which assimilate information from the most accurate geopotential models. The r_{cm} uncertainty estimated from those data is ± 0.00005 and contributes with ± 0.006 mas/cy to the δp uncertainty, whose final value has been set to ± 0.01 mas/cy in Table 4. A more conservative estimation of the r_{cm} influence can be derived from the difference between their values derived from PREM and 1066A models (see, e.g., Table 1 of Mathews et al. 1991). The uncertainties of r_{cm} and δp are then estimated to be ± 0.0008 and ± 0.06 mas/cy, respectively.

Therefore, the r_{cm} ratio is the dominant source of uncertainty under both assumptions, and the respective values, 0.01 and 0.06 mas/cy, have the same order of magnitude. In conclusion, our correction can be considered more accurate than the determination of the precession rate cited in the introduction even under the less favorable assumption.

In Table 5 our solution is compared to previous determinations of this second-order effect made for simpler Earth models. Column (2), with the IAU2006 heading, shows the value -46.78 mas/cy given to this correction in the P03 theory, the sum of the second-order lunisolar effects (a) and (b) in Capitaine et al. (2003) (Table 3). That value was first derived by Kinoshita & Souchay (1990) and kept unchanged by Williams (1994), Souchay & Kinoshita (1996), and in the final release of REN2000 (Souchay et al. 1999) as well. The original derivation only took into account four zonal terms and used an elliptical approximation for the orbits of the Moon and Sun. It is remarkable that their value only deviates 1% from our complete computation in the rigid Earth case, which includes 198 terms with the same orbits used for the first-order nutations and arrives at -45.51 mas/cy. In fact, an evaluation with a reduced set of the largest 11 terms (Getino et al. 2010) produces no significant variations. The value obtained by Roosbeek & Dehant (1998) in their rigid Earth RDAN97 theory appears in column (3) and is very close to ours.

Column (4) displays the results published for a two-layer Poincaré Earth model FNEG04 (Ferrándiz et al. 2004) with a total amount of -69.25 mas/cy, close to the recomputation of this work (see Table 4, column (2)). The small difference is due to the different set of parameters used in both investigations, P_{CW} and P_{FCN} in particular. Although for the sake of simplicity they are not displayed in Table 5, the results derived in this work are also consistent with the preliminary computations of the indirect terms for a similar two-layer Earth model (Ferrándiz et al. 2007), also shown in Table 4. Regarding the rigid case, the first detailed derivation of the sectorial contribution to the second-order effect on the longitude rate appeared jointly with the second-order nutation solution by Getino et al. (2010), who obtained -0.69 mas/cy, very close to the -0.70 mas/cy here. The row below the total value of each solution provides the respective deviations with respect to the IAU2006 reference correction (i.e., the REN2000 value). It is readily seen that all the deviations exceed 0.3 mas/cy and are thus above the current goals of accuracy of Earth rotation theories cited in Section 1.

The first-order precession rate p'_A is estimated by subtracting the corrections recognized in the relevant theory from the observed rate, and then the dynamical ellipticity is determined from p'_A taking into account Equation (2) (maybe in the context of a wider parameter fitting). Therefore, any change δp of the second-order correction with respect to its reference value in

IAU2006 must be compensated by an equal and opposite change $\delta p'_A = -\delta p$ of the first-order rate p'_A to keep the total rate unchanged. In turn, that variation induces an associated variation δH_d of the dynamical ellipticity such that $\delta H_d = (H_d/p'_A)\delta p$ and requires a rescaling of the nutation series (Escapa et al. 2016).

The next row of Table 5 displays the variations in each case. The uncertainties of the implied H_d variations can be estimated similarly. Taking into account the pair of former possible values of the δp uncertainty, the $\delta H_d = 553 \times 10^{-11}$ induced by the δp deviation -8.51 mas/cy can be attributed to an uncertainty of $\pm 0.4 \times 10^{-11}$ (and below $\pm 3.9 \times 10^{-11}$ in any case). It is worth underlining that this value is more than four times larger than the uncertainty in the estimation of H_d made by MHB2000, which was 1.2×10^{-9} , equivalent to 0.4 ppm.

The results of applying the former corrections to the reference value $H_d = 0.00327379448$ adopted in IAU2006 (Capitaine et al. 2003, 2009) are shown in the next row. Note that the uncertainty of the δH_d corrections should not be extrapolated to the modified values of H_d since the main source of error of p'_A still comes from the determination of the observed rate. Finally, the relative variations of the modified H_d values with respect to the reference one are provided in ppm in the last row. This makes the estimation of the magnitude of the corrections to the main nutation terms easy, taking $17''$ and $9''$ as proxies of maximum longitude and obliquity amplitudes.

4. CONCLUSIONS

The contribution to the precession rate due to the second-order solution associated with the main lunisolar potential has been considered in precession theory for decades; it is the third in magnitude, after the geodesic precession and the $J_2 - \text{tilt}$ rates. The IAU2006 precession theory adopted for it the value -46.78 mas/cy, corresponding to a simplified rigid Earth model and derived by means of Hamiltonian perturbation methods. In this paper we have carried out the computation of that effect in the Hamiltonian framework, assuming a two-layer Earth model and linear elasticity, and have arrived at an analytical expression to evaluate those contributions. They provide a final value of -55.29 mas/cy. The difference between both values amounts to -8.51 mas/cy and is noticeably larger than the -0.3 mas/cy that can be taken as a target of precession theory considering the observational accuracy and the GGOS goals for reference frames. Therefore, the value of this correction must be updated and the dynamical ellipticity of Earth revised accordingly.

This research has been partially supported by the Spanish government MINECO projects AYA2010-22039-C02-02 and AYA2016-79775-P (AEI/FEDER, UE). The authors acknowledge the valuable advice of an anonymous referee that helped to improve the manuscript.

APPENDIX A FUNCTIONS APPEARING IN THE PRECESSION FORMULAE

The functions $B_{i,p}$, $C_{i,\tau,p}$, and $D_{i,\tau,p}$ are those defined in Kinoshita (1977) and depend on the orbital coefficients $A_{i,p}^{(j)}$ provided in Kinoshita & Souchay (1990) or Navarro (2001).

Their expressions are

$$\begin{aligned} B_{i,p}(I) &= -\frac{1}{6}(3 \cos^2 I - 1)A_{i,p}^{(0)} - \frac{1}{2} \sin 2IA_{i,p}^{(1)} \\ &\quad - \frac{1}{4} \sin^2 IA_{i,p}^{(2)}, \\ C_{i,\tau,p}(I) &= -\frac{1}{4} \sin 2IA_{i,p}^{(0)} + \frac{1}{2}(1 + \tau \cos I) \\ &\quad \times (-1 + 2\tau \cos I)A_{i,p}^{(1)} \\ &\quad + \frac{1}{4} \tau \sin I(1 + \tau \cos I)A_{i,p}^{(2)}, \\ D_{i,\tau,p}(I) &= -\frac{1}{2} \sin^2 IA_{i,p}^{(0)} + \tau \sin I(1 + \tau \cos I)A_{i,p}^{(1)} \\ &\quad - \frac{1}{4}(1 + \tau \cos I)^2 A_{i,p}^{(2)}. \end{aligned} \quad (29)$$

The two-layer Earth nutation solution by Getino & Ferrándiz (2001) is given in terms of a set of functions that allows determining the amplitudes of the nutation series. When particularizing them to the Earth model considered in this work ($\Gamma = 0$), we have

$$F_{i,\tau}^{1a} = \frac{r_4 + n_{h,i,\tau}}{f_{1;i,\tau} f_{2;i,\tau}}, \quad F_{i,\tau}^{2a} = \frac{r_2}{f_{1;i,\tau} f_{2;i,\tau}}, \quad G_{i,\tau}^{1a} = \frac{r_3}{f_{1;i,\tau} f_{2;i,\tau}}, \quad (30)$$

with

$$f_{1;i,\tau} = m_1 + \omega_E - \tau n_i, \quad f_{2;i,\tau} = m_2 + \omega_E - \tau n_i. \quad (31)$$

These expressions can be written (Getino & Ferrándiz 2001) in terms of the BEP of the model $\{P_{\text{CW}}, P_{\text{FCN}}, A_c/A_m\}$ as

$$\begin{aligned} r_2 &= -\omega_E \left(1 + \frac{A_c}{A_m} + \frac{1}{P_{\text{CW}}} \right), \\ r_3 &= \omega_E \frac{(A_c/A_m)}{1 + (A_c/A_m)} \left(1 + \frac{A_c}{A_m} + \frac{1}{P_{\text{FCN}}} \right), \\ r_4 &= -\omega_E \left(1 + \frac{A_c}{A_m} + \frac{1}{P_{\text{FCN}}} \right), \\ m_1 &= \frac{\omega_E}{P_{\text{CW}}}, \\ m_2 &= -\omega_E \left(1 + \frac{1}{P_{\text{FCN}}} \right), \end{aligned} \quad (32)$$

where P_{CW} is associated with the Chandler wobble, P_{FCN} is associated with the FCN, and A_c/A_m is the ratio between the equatorial moment of inertia of the core and the mantle.

APPENDIX B COMPUTATION OF POISSON BRACKETS

In the notation we follow, the Poisson bracket of two smooth functions f and g of the canonical set (q, p) , is defined by the bilinear operation

$$\{f(q, p), g(q, p)\} = \sum_{i=1}^n \left(\frac{\partial f}{\partial q_i} \frac{\partial g}{\partial p_i} - \frac{\partial f}{\partial p_i} \frac{\partial g}{\partial q_i} \right). \quad (33)$$

In terms of the Andoyer-like canonical set of a two-layer Earth, the Poisson bracket is explicitly written as

$$\begin{aligned} \{f, g\} = & \frac{\partial f}{\partial \mu} \frac{\partial g}{\partial M} - \frac{\partial f}{\partial M} \frac{\partial g}{\partial \mu} + \frac{\partial f}{\partial \nu} \frac{\partial g}{\partial N} - \frac{\partial f}{\partial N} \frac{\partial g}{\partial \nu} \\ & + \frac{\partial f}{\partial \lambda} \frac{\partial g}{\partial \Lambda} - \frac{\partial f}{\partial \Lambda} \frac{\partial g}{\partial \lambda} + \frac{\partial f}{\partial \mu_c} \frac{\partial g}{\partial M_c} - \frac{\partial f}{\partial M_c} \frac{\partial g}{\partial \mu_c} \\ & + \frac{\partial f}{\partial \nu_c} \frac{\partial g}{\partial N_c} - \frac{\partial f}{\partial N_c} \frac{\partial g}{\partial \nu_c} + \frac{\partial f}{\partial \lambda_c} \frac{\partial g}{\partial \Lambda_c} - \frac{\partial f}{\partial \Lambda_c} \frac{\partial g}{\partial \lambda_c}. \end{aligned} \quad (34)$$

The former expression can be split into two parts formally identical to two Poisson brackets that would depend only on the subset of variables corresponding to the whole Earth, $\mathcal{S}_E = (\lambda, \mu, \nu, \Lambda, M, N)$, and that of the core, $\mathcal{S}_c = (\lambda_c, \mu_c, \nu_c, \Lambda_c, M_c, N_c)$,

$$\{f, g\} = \{f, g\}_E + \{f, g\}_c, \quad (35)$$

denoted with E and c subscripts.

In our developments the functions f and g may depend on the auxiliary angles σ , σ_c , I , and I_c , defined through

$$\cos \sigma = \frac{N}{M}, \quad \cos I = \frac{\Lambda}{M}, \quad \cos \sigma_c = \frac{N_c}{M_c}, \quad \cos I_c = \frac{\Lambda_c}{M_c}. \quad (36)$$

Hence, the derivatives with respect to canonical momenta must be performed by means of the following expressions (chain rule):

$$\begin{aligned} \frac{\partial}{\partial M} &= \left(\frac{\partial}{\partial M} \right) + \frac{\cot \sigma}{M} \frac{\partial}{\partial \sigma} + \frac{\cot I}{M} \frac{\partial}{\partial I}, \\ \frac{\partial}{\partial N} &= \left(\frac{\partial}{\partial N} \right) - \frac{1}{M \sin \sigma} \frac{\partial}{\partial \sigma}, \\ \frac{\partial}{\partial \Lambda} &= \left(\frac{\partial}{\partial \Lambda} \right) - \frac{1}{M \sin I} \frac{\partial}{\partial I}, \end{aligned} \quad (37)$$

and the analogous ones for the core variables (c subscript). The derivatives in parentheses refer to partial derivatives in the case of explicit dependence of the related variables.

Computing the Poisson brackets with the assistance of a symbolic processor is a good option to avoid annoying calculations and decrease the risk of flaws. The brackets in Section 2.3 were derived in this way, with the aid of Maple software. If hand calculation is preferred or used for checking purposes, we can tackle the advantage of the particular functional dependence of V_{per} , T_b , \mathcal{W}_p , and \mathcal{W}_l —Equations (17) and (20)—functions to simplify the process.

First, σ and σ_c angles are of small magnitude, about 10^{-6} rad (see, e.g., Getino 1995). Therefore, once the Poisson brackets have been computed, we can ignore their terms beyond the zeroth order in σ and σ_c , since we only need to compute the partial derivative with respect to I according to Equation (15). Furthermore, most of the brackets in Equation (21) can be computed simply as reduced $\{-, -\}_E$ or $\{-, -\}_c$ brackets, since only the third term in \mathcal{W}_p given in Equation (20) depends on \mathcal{S}_c variables and also on a sole \mathcal{S}_E variable, I .

Finally, the functional dependence of all the terms involved in the computation (except the one just mentioned) allows them to be arranged in one of the following forms:

$$\begin{aligned} f_0 &= S_0 V_0(I, \lambda), \quad g_0 = S_0 \mathcal{W}_0(I, \lambda), \\ f_1 &= S_1 V_1(I, \lambda, \mu, \nu), \quad g_1 = S_1 \mathcal{W}_1(I, \lambda, \mu, \nu), \end{aligned} \quad (38)$$

with $S_0 = 3 \cos^2 \sigma - 1$, $S_1 = \sin \sigma \cos \sigma$, and similarly for the core (c subscript).

For instance, the computation of the bracket $\{V_{\text{per}}, \mathcal{W}_p\}$ of Equation (22) reduces to the form $\{f_0 + f_1, g_0 + g_1\}_E$, which produces few nonvanishing terms. Specifically, we have

$$\begin{aligned} \{f_0, g_0\}_E &= \frac{4}{M \sin I} \left[-\frac{\partial V_0}{\partial \lambda} \frac{\partial \mathcal{W}_0}{\partial I} + \frac{\partial V_0}{\partial I} \frac{\partial \mathcal{W}_0}{\partial \lambda} \right] + O(\sigma), \\ \{f_0, g_1\}_E &= \{f_1, g_0\}_E = 0 + O(\sigma), \\ \{f_1, g_1\}_E &= \frac{1}{M} \left[\left(\frac{\partial V_1}{\partial \mu} - \frac{\partial V_1}{\partial \nu} \right) \mathcal{W}_1 - \left(\frac{\partial \mathcal{W}_1}{\partial \mu} - \frac{\partial \mathcal{W}_1}{\partial \nu} \right) V_1 \right] \\ &+ O(\sigma). \end{aligned} \quad (39)$$

The remaining Poisson brackets of Section 2.3 can be derived in a similar way.

REFERENCES

- Capitaine, N., Mathews, P. M., Dehant, V., Wallace, P. T., & Lambert, S. B. 2009, *CeMDA*, 103, 179
- Capitaine, N., Wallace, P. T., & Chapront, J. 2003, *A&A*, 412, 567
- Capitaine, N., Wallace, P. T., & Chapront, J. 2005, *A&A*, 432, 355
- Chen, W., Li, J. C., Ray, J., Shen, W. B., & Huang, C. L. 2015, *JGeod*, 89, 179
- Dziewonski, A. M., & Anderson, D. L. 1981, *PEPI*, 25, 297
- Efroimsky, M., & Williams, J. G. 2009, *CeMDA*, 104, 257
- Escapa, A. 2011, *CeMDA*, 110, 99
- Escapa, A., Ferrándiz, J. M., Baenas, T., et al. 2016, *PapGe*, 173, 861
- Ferrándiz, J. M., & Gross, R. S. 2015, in The New IAU/IAG Joint Working Group on Theory of Earth Rotation, International Association of Geodesy Symposia Ser. Vol. 143 (New York: Springer), 543
- Ferrándiz, J. M., Navarro, J. F., Escapa, A., & Getino, J. 2004, *AJ*, 128, 1407
- Ferrándiz, J. M., Navarro, J. F., Escapa, A., Getino, J., & Baenas, T. 2007, Influence of the Core on the Earth Precession, Proc. Workshop Organized in Honour of Professor Jacques Henrard at the Occasion of His Retirement (Presses Universitaires of Namur), 9
- Ferraz-Mello, S. 2007, Canonical Perturbation Theories. Degenerate Systems and Resonance (Berlin: Springer)
- Folgueira, M., Bizouard, C. H., & Souchay, J. 2001, *CeMDA*, 81, 191
- Folgueira, M., Souchay, J., & Kinoshita, H. 1998, *CeMDA*, 69, 373
- Folgueira, M., Souchay, J., & Kinoshita, H. 1999, *CeMDA*, 70, 147
- Fukushima, T. A. 2003, *AJ*, 126, 494
- Getino, J. 1995, *GeoJI*, 122, 803
- Getino, J., Escapa, A., & Miguel, D. 2010, *AJ*, 139, 1916
- Getino, J., & Ferrándiz, J. M. 1991, *CeMDA*, 51, 35
- Getino, J., & Ferrándiz, J. M. 1995, *CeMDA*, 61, 117
- Getino, J., & Ferrándiz, J. M. 1997, *GeoJI*, 130, 326
- Getino, J., & Ferrándiz, J. M. 2001, *MNRAS*, 322, 785
- Hilton, J. L., Capitaine, N., Chapront, J., et al. 2006, *CeMDA*, 94, 351
- Hori, G. I. 1966, *PASJ*, 18, 287
- Kaula, W. M. 1964, *RvGSP*, 2, 661
- Kinoshita, H. 1977, *CeMDA*, 15, 277
- Kinoshita, H., & Souchay, J. 1990, *CeMDA*, 48, 187
- Krásná, H., Böhm, J., & Schuh, H. 2013, *A&A*, 555, A29
- Kubo, Y. 1991, *CeMDA*, 50, 165
- Lambert, S. B., & Mathews, P. M. 2006, *A&A*, 453, 363
- Lieske, J. H., Lederle, T., Fricke, W., & Morando, B. 1977, *A&A*, 58, 1
- Liu, J.-C., & Capitaine, N. 2015, in Proc. Journées 2014 Syst. Ref. Possible Improvements of the IAU 2006 Precession Based on Recent Progress, ed. Z. Malkin & N. Capitaine (St. Petersburg: Pulkovo Observatory), 155

- Luzum, B., Capitaine, N., Fienga, A., et al. 2011, [CeMDA](#), **110**, 293
- Mathews, P. M., Buffet, B. A., Herring, T. A., & Shapiro, I. 1991, [JGR](#), **96**, 8243
- Mathews, P. M., Herring, T. A., & Buffet, B. A. 2002, [JGR](#), **107**, 2068
- Nastula, J., & Gross, R. 2015, [JGRB](#), **120**, 4474
- Navarro, J. F. 2001, PhD thesis, Univ. Alicante
- Petit, G., & Luzum, B. (ed.) 2010, IERS Conventions (2010), IERS Technical Note 36 (Verlag des Bundesamtes für Kartographie und Geodäsie, Frankfurt am Main)
- Plag, H., & Pearlman, M. 2009, Global Geodetic Observing System (Berlin: Springer)
- Roosbeek, F., & Dehant, V. 1998, [CeMDA](#), **70**, 215
- Souchay, J., & Kinoshita, H. 1996, [A&A](#), **312**, 1017
- Souchay, J., Losley, B., Kinoshita, H., & Figueira, M. 1999, [A&AS](#), **135**, 111
- Vondrák, J., Capitaine, N., & Wallace, P. 2011, [A&A](#), **534**, 1
- Williams, J. G. 1994, [AJ](#), **108**, 711
- Williams, J. G. 1995, [AJ](#), **110**, 1420