

## VARIATIONAL ANALYSIS IN SEMI-INFINITE AND INFINITE PROGRAMMING, I: STABILITY OF LINEAR INEQUALITY SYSTEMS OF FEASIBLE SOLUTIONS\*

M. J. CÁNOVAS<sup>†</sup>, M. A. LÓPEZ<sup>‡</sup>, B. S. MORDUKHOVICH<sup>§</sup>, AND J. PARRA<sup>†</sup>

**Abstract.** This paper concerns applications of advanced techniques of variational analysis and generalized differentiation to parametric problems of semi-infinite and infinite programming, where decision variables run over finite-dimensional and infinite-dimensional spaces, respectively. Part I is primarily devoted to the study of robust Lipschitzian stability of feasible solutions maps for such problems described by parameterized systems of infinitely many linear inequalities in Banach spaces of decision variables indexed by an arbitrary set  $T$ . The parameter space of admissible perturbations under consideration is formed by all bounded functions on  $T$  equipped with the standard supremum norm. Unless the index set  $T$  is finite, this space is intrinsically infinite-dimensional (nonreflexive and nonseparable) of the  $l_\infty$  type. By using advanced tools of variational analysis and exploiting specific features of linear infinite systems, we establish complete characterizations of robust Lipschitzian stability entirely via their initial data with computing the exact bound of Lipschitzian moduli. A crucial part of our analysis addresses the precise computation of the coderivative of the feasible set mapping and its norm. The results obtained are new in both semi-infinite and infinite frameworks.

**Key words.** semi-infinite and infinite programming, variational analysis, linear infinite inequality systems, robust stability, generalized differentiation, coderivatives

**AMS subject classifications.** 90C34, 90C05, 49J52, 49J53, 65F22

**DOI.** 10.1137/090765948

**1. Introduction.** This paper mainly deals with parameterized *infinite* systems of linear inequalities

$$(1.1) \quad \mathcal{F}(p) := \{x \in X \mid \langle a_t^*, x \rangle \leq b_t + p_t, t \in T\}, \quad p = (p_t)_{t \in T},$$

with an *arbitrary index set*  $T$ , where  $x \in X$  is a *decision* variable belonging to an arbitrary Banach space  $X$  (which may be finite-dimensional), and where  $p = (p_t)_{t \in T} \in P$  is a functional *parameter* taking values in the prescribed Banach space  $P$  of perturbations specified below. Infinite inequality systems of this type are important in various areas of mathematics and applications, while our primary interest in them is driven by applications to problems of *semi-infinite* and *infinite programming* corresponding to finite-dimensional and infinite-dimensional decision spaces  $X$ , respectively; see, e.g., the books [1, 5, 19, 37] and the references therein. Some applications of the results obtained in this paper to necessary optimality conditions for semi-infinite and infinite programs can be found in [11].

The data of (1.1) are given as follows:

---

\*Received by the editors July 22, 2009; accepted for publication (in revised form) September 23, 2009; published electronically December 16, 2009. This research was partially supported by grants MTM2005-08572-C03 (01-02) from MEC (Spain) and FEDER (EU), MTM2008-06695-C03 (01-02) from MICINN (Spain), and ACOMP/2009/047&133 from Generalitat Valenciana (Spain).  
<http://www.siam.org/journals/siopt/20-3/76594.html>

<sup>†</sup>Center of Operations Research, Miguel Hernández University of Elche, 03202 Elche (Alicante), Spain (canovas@umh.es, parra@umh.es).

<sup>‡</sup>Department of Statistics and Operations Research, University of Alicante, 03080 Alicante, Spain (marco.antonio@ua.es).

<sup>§</sup>Department of Mathematics, Wayne State University, Detroit, MI 48202 (boris@math.wayne.edu). The research of this author was partially supported by the National Science Foundation (USA) under grant DMS-0603846.

- $a_t^* \in X^*$  for all  $t \in T$ , where the space  $X^*$  is topologically dual to  $X$  with the canonical pairing  $\langle \cdot, \cdot \rangle$  between  $X$  and  $X^*$ . If no confusion arises, we use the *same notation*  $\| \cdot \|$  for the given norm in  $X$  and the corresponding dual norm in  $X^*$  defined by

$$\|x^*\| := \sup \{ \langle x^*, x \rangle \mid \|x\| \leq 1 \}, \quad x^* \in X^*.$$

We always assume that  $a_t^*$  are *fixed* and *arbitrary* in  $X^*$  for all  $t \in T$ .

- $b_t \in \mathbb{R}$  for all  $t \in T$ . We identify the collection  $\{b_t \mid t \in T\}$  with the real-valued function  $b: T \rightarrow \mathbb{R}$  and assume that it is *fixed* and *arbitrary*.

- $p_t = p(t) \in \mathbb{R}$  for all  $t \in T$ . These functional parameters  $p: T \rightarrow \mathbb{R}$  are our varying *perturbations*, which are taken from the Banach *parameter space*  $P := l_\infty(T)$  of all bounded functions on  $T$  with the supremum norm

$$(1.2) \quad \|p\|_\infty := \sup_{t \in T} |p_t| = \sup \{ |p(t)| \mid t \in T \}$$

(see, e.g., [17]), where the subscript “ $\infty$ ” is omitted if no confusion arises. When the index set  $T$  is compact (which is not assumed in this paper) and the perturbations  $p(\cdot)$  are restricted to be continuous on  $T$ , the maximum is realized in (1.2), and thus the parameter space  $l_\infty(T)$  reduces to the classical space  $\mathcal{C}(T)$  of continuous functions over a compact set.

The primary goal of this paper is to obtain comprehensive characterizations of *robust Lipschitzian stability* of infinite inequality systems (1.1) expressed entirely in terms of their initial data. By robust Lipschitzian stability we understand the fulfillment of the so-called *Lipschitz-like* (known also as Aubin) property of the mapping  $\mathcal{F}(p)$  in (1.1) *around* the reference point. This property is stable with respect to small perturbations of parameters and is crucial for both qualitative and quantitative/numerical aspects of optimization theory and applications; see, e.g., [31, 32, 35] and section 2 for more details and references.

To establish constructive characterizations of robust Lipschitzian stability in this paper and to derive efficient optimality conditions for semi-infinite and infinite programs in [11], we develop an advanced approach of *variational analysis* based on *generalized differentiation*. To the best of our knowledge, this approach is new in the literature on semi-infinite and infinite programming despite many publications related to various stability properties and applications of linear infinite inequality systems, most of which concern the case of finite-dimensional spaces  $X$  of decision variables (i.e., in the semi-infinite programming framework); see, e.g., [1, 19] for comprehensive overviews on this field and also [5], which is confined to the parameter space of continuous perturbations  $P = \mathcal{C}(T)$  when the index set  $T$  is a compact Hausdorff space. We refer the reader to [12] for the study of *qualitative stability* (formalized through certain semicontinuity properties of feasible solution and optimal solution mappings) in the framework of  $X = \mathbb{R}^n$ , an arbitrary index set  $T$ , and arbitrary perturbations. In the same semi-infinite context, for a *quantitative perspective* (through Lipschitzian properties), the reader is directed to [7], and to [6] for the case of continuous perturbations. Let us mention the recent paper [14] addressing the case of infinite linear programming from the viewpoint of qualitative stability. We also refer the reader to, e.g., [8, 9, 13, 15, 18, 24, 25] for the study of convex semi-infinite and infinite programs and to [21, 22, 37] for their smooth nonlinear counterparts.

The approach of this paper is mainly based on *coderivative analysis* of the parametric linear infinite inequality systems  $\mathcal{F}$  in (1.1), which eventually leads us to *com-*

plete characterizations of robust Lipschitzian stability for the parametric sets of feasible solutions in infinite/semi-infinite programming expressed entirely via their initial data with precise computing the *exact bound* of Lipschitzian moduli.

Coderivatives of set-valued mappings introduced in [27] have been well recognized as a powerful tool of variational analysis and its numerous applications, particularly to problems of optimization and control; see, e.g., [3, 23, 30, 31, 32, 33, 35, 36] and the references therein. However, we are not familiar with any implementation of coderivatives in problems of infinite or semi-infinite programming as well as with their application to analyzing stability of linear infinite inequality systems of type (1.1) in finite or infinite dimensions.

The power of coderivatives in variational analysis and its applications comes, first, from the possibility of obtaining in their terms verifiable *pointwise characterizations* of *robust Lipschitzian* properties of set-valued mappings (as well as of the equivalent properties of *metric regularity* and *linear openness* for the inverse mappings) and of deriving *necessary optimality conditions* in rather general settings. These developments are strongly supported by comprehensive pointwise *coderivative calculus* based on *variational/extremal principles* of advanced variational analysis; see [31, 32] and the references therein. However, a number of results in this vein are limited in infinite dimensions. In particular, the available coderivative characterizations of the Lipschitz-like property of closed-graph mappings  $F: Z \rightrightarrows Y$  obtained in [31, Theorem 4.10] require that both spaces  $Z$  and  $Y$  are *Asplund* (i.e., every separable subspace of them has a separable dual) while the precise coderivative formula for computing the *exact Lipschitzian bound* is established therein via the coderivative norm under the *finite-dimensionality* assumption on  $Z$ . But this is *never the case* for our infinite inequality system  $\mathcal{F}: P \rightrightarrows X$  from (1.1), where the parameter space ( $Z =$ )  $P = l_\infty(T)$  is always *infinite-dimensional* and *not Asplund* unless the index set  $T$  is finite!

This paper contains new and fairly comprehensive results in the aforementioned directions for the infinite/semi-infinite systems under consideration, which essentially take into account underlying specific features of the infinite inequality constraints (1.1) largely related to the possibility of employing an appropriate extended version of the fundamental *Farkas lemma* for infinite systems of linear inequalities in *general Banach spaces*.

The rest of paper is organized as follows. Section 2 presents some preliminary material from convex and variational analysis widely used in formulations and proofs of the subsequent main results. In section 3 we provide *precise calculations* of the basic *coderivative*  $D^*\mathcal{F}$  and its *norm* at the reference/nominal point for the feasible solution map  $\mathcal{F}: l_\infty(T) \rightrightarrows X$  in (1.1) via the initial data of  $\mathcal{F}$  in the general case of an arbitrary index set  $T$  and an arbitrary Banach space  $X$  of decision variables.

Section 4 is devoted to deriving *coderivative characterizations* of *robust stability* for the feasible solution system (1.1) of infinite inequalities with an arbitrary index set  $T$ , which are explicitly expressed in terms of the initial data  $\{a_t^*, b_t, t \in T\}$ . We establish verifiable *criteria* (i.e., necessary and sufficient conditions) for the fulfillment of the Lipschitz-like (and hence the classical local Lipschitzian) property of  $\mathcal{F}$  around the reference points and derive furthermore the *precise formulas* for computing the *exact bounds* of Lipschitzian moduli in the case of general Banach spaces  $X$ . It is worth mentioning that the criteria and exact bound formulas obtained in this section are represented in the conventional *coderivative form* of variational analysis as in [31, Theorem 4.10] for the case of abstract set-valued mappings, but with *no* Asplund space and finite-dimensionality requirements imposed therein. In fact, the latter requirements are *never* satisfied for the infinite linear inequality systems (1.1)

under consideration in either the infinite programming or semi-infinite programming framework.

Our notation is basically standard and conventional in the areas of variational analysis and infinite/semi-infinite programming; see, e.g., [19, 31, 35]. Unless otherwise stated, all the spaces under consideration are *Banach* with the corresponding norm  $\|\cdot\|$ . Recall that  $w^*$  indicates the weak\* topology of a dual space, and we use the symbol  $w^*$ -lim for the weak\* *topological* limit, which generally means the weak\* convergence of *nets* denoted usually by  $\{x_\nu^*\}_{\nu \in \mathcal{N}}$ . In the case of sequences we use the standard notation  $:= \{1, 2, \dots\}$  for the collection of all natural numbers.

Given a subset  $\Omega \subset Z$  of a Banach space, the symbols  $\text{int } \Omega$ ,  $\text{cl } \Omega$ ,  $\text{co } \Omega$ , and  $\text{cone } \Omega$  stand, respectively, for the interior, closure, convex hull, and conic convex hull of  $\Omega$ ; the notation  $\text{cl}^* \Theta$  signifies the weak\* closure of a subset  $\Theta \subset Z^*$  in the dual space. Given a set-valued mapping  $F: Z \rightrightarrows Y$ , we denote its *domain*, *graph*, and *inverse* by, respectively,

$$\text{dom } F = \{z \in Z \mid F(z) \neq \emptyset\}, \quad \text{gph } F := \{(z, y) \in Z \times Y \mid y \in F(z)\},$$

and  $F^{-1}(y) := \{z \in Z \mid (z, y) \in \text{gph } F\}$ . Considering finally an arbitrary index set  $T$ , let  $\mathbb{R}^T$  be the *product space* of  $\lambda = (\lambda_t \mid t \in T)$  with  $\lambda_t \in \mathbb{R}$  for all  $t \in T$ , let  $\mathbb{R}^{(T)}$  be the collection of  $\lambda \in \mathbb{R}^T$  such that  $\lambda_t \neq 0$  for *finitely many*  $t \in T$ , and let  $\mathbb{R}_+^{(T)}$  be the *positive cone* in  $\mathbb{R}^{(T)}$  defined by

$$\mathbb{R}_+^{(T)} := \{\lambda \in \mathbb{R}^{(T)} \mid \lambda_t \geq 0 \text{ for all } t \in T\}.$$

**2. Basic definitions and preliminaries.** In this section we discuss the basic notions and tools needed for our subsequent analysis and results. As mentioned in section 1, a major focus of this paper is *robust Lipschitzian stability* of the feasible solution map given by (1.1). By such a robust stability we understand Lipschitzian behavior *around* (i.e., in a neighborhood) of the reference point. The most natural formalization of this behavior widely recognized in variational (as well as in general nonlinear) analysis is known as the Lipschitz-like or Aubin property, which can be viewed as a graphical localization (in the set-valued case) of the classical local Lipschitzian property of single-valued and set-valued mappings.

Given a set-valued mapping  $F: Z \rightrightarrows Y$  between Banach spaces, we say that  $F$  is *Lipschitz-like* around  $(\bar{z}, \bar{y}) \in \text{gph } F$  with modulus  $\ell \geq 0$  if there are neighborhoods  $U$  of  $\bar{z}$  and  $V$  of  $\bar{y}$  such that

$$(2.1) \quad F(z) \cap V \subset F(u) + \ell \|z - u\| \mathbb{B} \text{ for any } z, u \in U,$$

where  $\mathbb{B}$  stands for the closed unit ball in the space in question. The infimum of moduli  $\{\ell\}$  over all the combinations of  $\{\ell, U, V\}$  satisfying (2.1) is called the *exact Lipschitzian bound* of  $F$  around  $(\bar{z}, \bar{y})$  and is labeled as  $\text{lip } F(\bar{z}, \bar{y})$ . If  $V = Y$  in (2.1), this relationship signifies the classical (Hausdorff) *local Lipschitzian* property of  $F$  around  $\bar{z}$  with the exact Lipschitzian bound denoted by  $\text{lip } F(\bar{z})$  in this case.

It is worth mentioning that the Lipschitz-like property of an arbitrary mapping  $F: Z \rightrightarrows Y$  between Banach spaces is equivalent to two other fundamental properties in nonlinear analysis but for the *inverse* mapping  $F^{-1}: Y \rightrightarrows Z$ , namely, to the *metric regularity* of  $F^{-1}$  and to the *linear openness* of  $F^{-1}$  around  $(\bar{y}, \bar{z})$ , with the corresponding relationships between their exact bounds (see, e.g., [20, 31, 35]).

It is well known that the exact Lipschitzian bound of  $F$  around  $(\bar{z}, \bar{y})$  admits the following limiting representation via the *distance function* to a set:

$$(2.2) \quad \text{lip } F(\bar{z}, \bar{y}) = \limsup_{(z, y) \rightarrow (\bar{z}, \bar{y})} \frac{\text{dist}(y; F(z))}{\text{dist}(z; F^{-1}(y))},$$

where  $\inf \emptyset = \infty$  (and hence  $\text{dist}(x; \emptyset) = \infty$ ) as usual and where  $0/0 := 0$ . We have accordingly that  $\text{lip } F(\bar{z}, \bar{y}) = \infty$  if  $F$  is not Lipschitz-like around  $(\bar{z}, \bar{y})$ .

A remarkable fact consists of the possibility of *characterizing* pointwisely the (derivative-free) Lipschitz-like property of  $F$  around  $(\bar{z}, \bar{y})$ —and hence its local Lipschitzian, metric regularity, and linear openness counterparts—in terms of a *dual-space* construction of generalized differentiation called the *coderivative* of  $F$  at  $(\bar{z}, \bar{y}) \in \text{gph } F$ . The latter is a positively homogeneous multifunction  $D^*F(\bar{z}, \bar{y}): Y^* \rightrightarrows Z^*$  defined by

$$(2.3) \quad D^*F(\bar{z}, \bar{y})(y^*) := \{z^* \in Z^* \mid (z^*, -y^*) \in N((\bar{z}, \bar{y}); \text{gph } F)\}, \quad y^* \in Y^*,$$

where  $N(\cdot; \Omega)$  stands for the collection of generalized normals to a set at a given point known as the basic, or limiting, or Mordukhovich, normal cone; see, e.g., [26, 31, 35, 36] and the references therein. When both  $Z$  and  $Y$  are *finite-dimensional*, it is proved in [28] (cf. also [35, Theorem 9.40]) that a closed-graph mapping  $F: Z \rightrightarrows Y$  is Lipschitz-like around  $(\bar{z}, \bar{y}) \in \text{gph } F$  *if and only if*

$$(2.4) \quad D^*F(\bar{z}, \bar{y})(0) = \{0\},$$

and that the exact Lipschitzian bound of moduli  $\{\ell\}$  in (2.1) is computed by

$$(2.5) \quad \text{lip } F(\bar{z}, \bar{y}) = \|D^*F(\bar{z}, \bar{y})\| := \sup \{\|z^*\| \mid z^* \in D^*F(\bar{z}, \bar{y})(y^*), \|y^*\| \leq 1\}.$$

The situation is significantly more involved in infinite dimensions. It is proved in [29] (see also [31, Theorem 4.10]) that a closed-graph mapping  $F: Z \rightrightarrows Y$  is Lipschitz-like around  $(\bar{z}, \bar{y}) \in \text{gph } F$  *if and only if* the coderivative condition (2.4) holds in terms of the so-called *mixed coderivative* (which reduces to (2.3) in finite dimensions and the setting considered in this paper) together with a certain “partial sequential normal compactness” condition (which automatically holds in finite dimensions and in the setting of this paper) provided that both spaces  $Z$  and  $Y$  are *Asplund*. The latter property is defined in section 1; we also refer the reader to [17, 31, 34] for more details and various characterizations of this remarkable and well-investigated subclass of Banach spaces that includes, in particular, all reflexive spaces, while it does not include, e.g., the classical spaces  $\mathcal{C}$ ,  $l_1$ ,  $L_1$ ,  $l_\infty$ , and  $L_\infty$ .

The situation is even more complicated with infinite-dimensional extensions of the exact bound formula in (2.5). The aforementioned results of [29, 31] give merely upper and lower estimates for  $\text{lip } F(\bar{z}, \bar{y})$ , which ensure the precise equality in (2.5) in our setting here provided that  $Y$  is *Asplund* while  $Z$  is *finite-dimensional*.

The set-valued mapping  $F: Z \rightrightarrows Y$  considered in this paper is  $\mathcal{F}: l_\infty(T) \rightrightarrows X$  defined by the infinite system of linear inequalities (1.1); in what follows we always assume that the index set  $T$  is *infinite*, which is a characteristic feature of infinite and semi-infinite programs. In this setting the domain/parameter space  $Z = l_\infty(T)$  is an *infinite-dimensional* Banach that is *never Asplund*. Also, we do not suppose in this paper that our decision space  $X$  is anything but *arbitrary Banach*.

In this general setting for (1.1) we show that the coderivative condition (2.4) is *necessary and sufficient* for the Lipschitz-like property of  $F = \mathcal{F}$  around the reference/

nominal solution  $(\bar{p}, \bar{x}) \in \text{gph } \mathcal{F}$  and (2.5) is a *precise formula* for computing the exact Lipschitzian bound  $\text{lip } \mathcal{F}(\bar{p}, \bar{x})$ . This is *exactly what we have in finite dimensions*, while it is far removed from being a part of the infinite-dimensional variational theory in [31]. Moreover, we express the relationships in (2.4) and (2.5) *explicitly* in terms of the *initial data* of (1.1).

To proceed further, observe that the graph

$$(2.6) \quad \text{gph } \mathcal{F} = \{(p, x) \in l_\infty(T) \times X \mid \langle a_t^*, x \rangle \leq b_t + p_t \text{ for all } t \in T\}$$

of the mapping  $\mathcal{F}: l_\infty(T) \rightrightarrows X$  in (1.1) is *convex*. Hence the basic normal cone to  $\text{gph } \mathcal{F}$  at  $(\bar{p}, \bar{x}) \in \text{gph } \mathcal{F}$  reduces to

$$N((\bar{p}, \bar{x}); \text{gph } \mathcal{F}) = \{(p^*, x^*) \in l_\infty(T)^* \times X^* \mid \langle (p^*, x^*), (p, x) - (\bar{p}, \bar{x}) \rangle \leq 0 \text{ for } (p, x) \in \text{gph } \mathcal{F}\}$$

and the coderivative (2.3) of  $\mathcal{F}$  admits the representation

$$(2.7) \quad D^* \mathcal{F}(\bar{p}, \bar{x})(x^*) = \{p^* \in l_\infty(T)^* \mid \langle p^*, \bar{p} \rangle - \langle x^*, \bar{x} \rangle = \max_{(p,x) \in \text{gph } \mathcal{F}} [\langle p^*, p \rangle - \langle x^*, x \rangle]\}.$$

Let us now present two preliminary results that play an important role in our subsequent analysis. The first one is taken from [13, Lemma 2.4] and can be viewed as an *extended Farkas lemma* for linear infinite inequality systems in Banach spaces. An alternative proof can be derived from [38, Theorem 4].

LEMMA 2.1 (extended Farkas lemma). *Let  $p \in \text{dom } \mathcal{F}$  for the infinite system (1.1) with a Banach decision space  $X$ , and let  $(x^*, \alpha) \in X^* \times \mathbb{R}$ . The following are equivalent:*

- (i) *We have  $\langle x^*, x \rangle \leq \alpha$  whenever  $x \in \mathcal{F}(p)$ , i.e.,*

$$[\langle a_t^*, x \rangle \leq b_t + p_t \text{ for all } t \in T] \implies [\langle x^*, x \rangle \leq \alpha].$$

- (ii) *The pair  $(x^*, \alpha)$  satisfies the inclusion*

$$(x^*, \alpha) \in \text{cl}^* \text{cone}[\{(a_t^*, b_t + p_t) \mid t \in T\} \cup \{(0, 1)\}] \text{ with } 0 \in X^*.$$

Throughout the paper we largely use the parametric *characteristic sets*

$$(2.8) \quad C(p) := \text{co}\{(a_t^*, b_t + p_t) \mid t \in T\}, \quad p \in l_\infty(T),$$

and suppose with no loss of generality that our *nominal parameter* is the zero function  $\bar{p} = 0$  in the parameter space  $l_\infty(T)$ .

Let us recall a well-recognized *qualification condition* for linear infinite inequalities, which is often used in problems of semi-infinite and infinite programming.

DEFINITION 2.2 (strong Slater condition). *We say that the infinite system (1.1) satisfies the strong Slater condition (SSC) at  $p = (p_t)_{t \in T}$  if there is  $\hat{x} \in X$  such that*

$$(2.9) \quad \sup_{t \in T} [\langle a_t^*, \hat{x} \rangle - b_t - p_t] < 0.$$

*Furthermore, every point  $\hat{x} \in X$  satisfying condition (2.9) is a strong Slater point for system (1.1) at  $p = (p_t)_{t \in T}$ .*

The next result contains several equivalent descriptions and interpretations of the strong Slater condition used in what follows; the most important is the equivalence

(i) $\iff$ (ii). Note that a similar equivalence can be found in [14] for more general convex systems in locally convex spaces with different spaces of associated parameters. We include here a simplified proof of this equivalence for the reader's convenience.

LEMMA 2.3 (equivalent descriptions of the strong Slater condition). *Let  $X$  be a Banach space, and let  $p \in \text{dom } \mathcal{F}$  for the linear infinite inequality system (1.1). Then the following properties are equivalent:*

- (i)  $\mathcal{F}$  satisfies the strong Slater condition at  $p$ .
- (ii)  $(0, 0) \notin \text{cl}^* C(p)$  via the characteristic set from (2.8).
- (iii)  $p \in \text{int}(\text{dom } \mathcal{F})$ .
- (iv)  $\mathcal{F}$  is Lipschitz-like around  $(p, x)$  for all  $x \in \mathcal{F}(p)$ .

Moreover, if the set  $\{a_t^* \mid t \in T\}$  is bounded in  $X^*$ , the conditions above are equivalent to

- (v) there exists  $\hat{x} \in X$  such that  $(p, \hat{x}) \in \text{int}(\text{gph } \mathcal{F})$ .

*Proof.* We begin with the proof of (i) $\implies$ (ii). Arguing by contradiction, assume that  $(0, 0) \in \text{cl}^* C(p)$ . Then there is a net  $\{\lambda_\nu\}_{\nu \in \mathcal{N}} \in \mathbb{R}_+^{(T)}$  satisfying  $\sum_{t \in T} \lambda_{t\nu} = 1$  for all  $\nu \in \mathcal{N}$  and the limiting condition

$$(2.10) \quad (0, 0) = w^*\text{-}\lim_{\nu} \sum_{t \in T} \lambda_{t\nu} (a_t^*, b_t + p_t).$$

If  $\hat{x}$  is a strong Slater point for system (1.1) at  $p$ , we find  $\vartheta > 0$  such that

$$\langle a_t^*, \hat{x} \rangle - b_t - p_t \leq -\vartheta \quad \text{for all } t \in T.$$

Then (2.10) leads to the following contradiction:

$$0 = \langle 0, \hat{x} \rangle + 0 \cdot (-1) = \lim_{\nu} \sum_{t \in T} \lambda_{t\nu} (\langle a_t^*, \hat{x} \rangle + (b_t + p_t) \cdot (-1)) \leq -\vartheta.$$

Let us next justify the converse implication (ii) $\implies$ (i). By [13, Theorem 3.1] we have

$$p \in \text{dom } \mathcal{F} \iff (0, -1) \notin \text{cl}^* \text{cone}\{(a_t^*, b_t + p_t) \mid t \in T\}.$$

Then the *strong separation theorem* ensures the existence of  $(0, 0) \neq (v, \alpha) \in X \times \mathbb{R}$  with

$$(2.11) \quad \langle a_t^*, v \rangle + \alpha(b_t + p_t) \leq 0 \quad \text{for all } t \in T,$$

$$\langle 0, v \rangle + (-1)\alpha = -\alpha > 0.$$

At the same time by (ii) we have  $(0, 0) \neq (z, \beta) \in X \times \mathbb{R}$  and  $\gamma \in \mathbb{R}$  for which

$$(2.12) \quad \langle a_t^*, z \rangle + \beta(b_t + p_t) \leq \gamma < 0 \quad \text{whenever } t \in T.$$

Consider further the combination

$$(u, \eta) := (z, \beta) + \lambda(v, \alpha),$$

and select  $\lambda > 0$  to be sufficiently large to ensure that  $\eta < 0$ . Defining now  $\hat{x} := -\eta^{-1}u$ , we observe from (2.11) and (2.12) that

$$\langle a_t^*, \hat{x} \rangle - b_t - p_t = -\eta^{-1}(\langle a_t^*, u \rangle + \eta(b_t + p_t)) \leq -\eta^{-1}\gamma < 0.$$

This allows us to conclude that  $\hat{x}$  is a strong Slater point for system (1.1) at  $p$ .

To prove implication (i) $\implies$ (iii), assume that  $\hat{x}$  is a strong Slater point for system (1.1) at  $p$ , and find  $\vartheta > 0$  such that

$$\langle a_t^*, \hat{x} \rangle - b_t - p_t \leq -\vartheta \text{ for all } t \in T.$$

Then it is obvious that for any  $q \in l_\infty(T)$  with  $\|q\| < \vartheta$  we have  $\hat{x} \in \mathcal{F}(p + q)$ . Therefore,  $p + q \in \text{dom } \mathcal{F}$ , and thus (iii) holds.

Let us further proceed with justifying implication (iii) $\implies$ (i). If  $p \in \text{int}(\text{dom } \mathcal{F})$ , then  $p + q \in \text{dom } \mathcal{F}$  provided that  $q_t = -\vartheta$  as  $t \in T$  and that  $\vartheta > 0$  is sufficiently small. Thus every  $\hat{x} \in \mathcal{F}(p + q)$  is a strong Slater point for the infinite system (1.1) at  $p$ .

The equivalence between (iii) and (iv) is a consequence of the classical Robinson–Ursescu closed-graph/metric regularity theorem; see, e.g., [20] and [31, Chapter 4] and references therein for more discussion.

It remains to consider condition (v). We can easily observe that (v) always implies (iii) and hence the other conditions of the lemma. Suppose now that the set  $\{a_t^* \mid t \in T\}$  is bounded in  $X^*$ , and show that in this case (i) implies (v). Select  $M \geq 0$  such that  $\|a_t^*\| \leq M$  for every  $t \in T$ , and take  $\hat{x} \in X$  satisfying (2.9). Denote

$$\gamma := -\sup_{t \in T} [\langle a_t^*, \hat{x} \rangle - b_t - p_t] > 0,$$

and consider any pair  $(p', u) \in l_\infty(T) \times X$  satisfying the relationships

$$\|u\| \leq \eta := \gamma / (M + 1) > 0 \text{ and } \|p'\| \leq \eta.$$

It is easy to see that for such  $(p', u)$  and every  $t \in T$  we have

$$\langle a_t^*, \hat{x} + u \rangle - b_t - p_t - p'_t \leq -\gamma + M \|u\| + \|p'\| \leq \eta(M + 1) - \gamma = 0,$$

and so  $(p + p', \hat{x} + u) \in \text{gph } \mathcal{F}$ . Thus  $(p, \hat{x}) \in \text{int}(\text{gph } \mathcal{F})$ , which gives (i) $\implies$ (v) and completes the proof of the lemma.  $\square$

*Remark 2.4* (relationships between interiority and boundedness conditions). Note that we have the nonempty *interiority condition*  $\text{int}(\text{gph } \mathcal{F}) \neq \emptyset$  provided that the set  $\{a_t^* \mid t \in T\}$  is *bounded* in  $X^*$  and that  $\text{gph } \mathcal{F} \neq \emptyset$ . Also we have  $\text{int}(\text{dom } \mathcal{F}) \neq \emptyset$  if  $\text{gph } \mathcal{F} \neq \emptyset$  with no boundedness assumption.

To justify the first statement, take  $(p, x) \in \text{gph } \mathcal{F}$  and  $\|a_t^*\| \leq M < \infty$  for every  $t \in T$ , and define  $e \in l_\infty(T)$  by  $e_t := 1$  for all  $t \in T$ . Then arguing similarly to the proof of the last implication in Lemma 2.1, we get  $(p + e + p', x + u) \in \text{gph } \mathcal{F}$  whenever  $\|p'\| \leq \eta := 1 / (M + 1)$  and  $\|u\| \leq \eta$ . This gives  $(p + e, x) \in \text{int}(\text{gph } \mathcal{F})$ . If, furthermore,  $(p, x) \in \text{gph } \mathcal{F}$ , we can easily check that  $x \in \mathcal{F}(p + e + p')$  provided that  $\|p'\| \leq 1$ , and therefore  $p + e \in \text{int}(\text{dom } \mathcal{F})$ .

The major space for our consideration in this paper is the parameter space  $l_\infty(T)$  of bounded functions  $p: T \rightarrow \mathbb{R}$  on  $T$  with the supremum norm (1.2). It is obviously a Banach space that is *never finite-dimensional* when the index set  $T$  is infinite, which is our standing assumption. Let us show that it is *never Asplund*.

**PROPOSITION 2.5** (parameter space is never Asplund). *The parameter space  $l_\infty(T)$  is Asplund if and only if the index set  $T$  is finite.*

*Proof.* If  $T$  is countable (i.e.,  $T = \mathbb{N}$  and the parameter space is the classical space of sequences  $l_\infty$ ), the proof can be found in [34, Example 1.21]; in fact, this space is



not even weak Asplund. The same arguments can be adapted for any infinite index set  $T$ .  $\square$

Finally, in this section, we recall a convenient description of the topological dual space  $l_\infty(T)^*$  to the parameter space  $l_\infty(T)$ . According to [16], there is an isometric isomorphism between  $l_\infty(T)^*$  and the space of *bounded and additive measures*

$$ba(T) = \{\mu: 2^T \rightarrow \mathbb{R} \mid \mu \text{ is bounded and additive}\}$$

satisfying the relationship

$$\langle \mu, p \rangle = \int_T p_t \mu(dt) \quad \text{with } p = (p_t)_{t \in T}.$$

The dual norm on  $ba(T)$  corresponding to (1.2) is the *total variation* of  $\mu \in ba(T)$  on the index set  $T$  defined by

$$\|\mu\| := \sup_{A \subset T} \mu(A) - \inf_{B \subset T} \mu(B).$$

In what follows we always identify the measure space  $ba(T)$  with the dual parameter space  $l_\infty(T)^*$  and use, for the notational unification,  $p^* \in l_\infty(T)^*$  instead of  $\mu \in ba(T)$ .

**3. Computing coderivatives and coderivative norms for linear infinite inequality systems.** In this section we establish a constructive representation of the coderivative  $D^*\mathcal{F}(0, \bar{x})$  for the feasible solution map  $\mathcal{F}$  at the nominal point  $(0, \bar{x})$  and compute its norm  $\|D^*\mathcal{F}(0, \bar{x})\|$  in terms of the initial data of the linear infinite inequality system (1.1). Let us first describe the *normal cone* to the convex graph (2.6) employing the *extended Farkas lemma* presented above. In what follows  $\delta_t$  denotes the classical *Dirac measure* at  $t \in T$  satisfying

$$\langle \delta_t, p \rangle = p_t \quad \text{as } t \in T \quad \text{for } p = (p_t)_{t \in T} \in l_\infty(T).$$

**PROPOSITION 3.1** (computing normals to the graphical set of feasible solutions). *Let  $(\bar{p}, \bar{x}) \in \text{gph } \mathcal{F}$  for the graphical set (2.6) with a Banach decision space  $X$ , and let  $(p^*, x^*) \in l_\infty(T)^* \times X^*$ . Then we have  $(p^*, x^*) \in N((\bar{p}, \bar{x}); \text{gph } \mathcal{F})$  if and only if*

$$(3.1) \quad (p^*, x^*, \langle p^*, \bar{p} \rangle + \langle x^*, \bar{x} \rangle) \in \text{cl}^* \text{cone}[\{(-\delta_t, a_t^*, b_t) \mid t \in T\} \cup \{(0, 0, 1)\}],$$

where  $0 \in l_\infty(T)^*$  and  $0 \in X^*$  stand for the first and second entry of the last triple in (3.1), respectively. Furthermore, the inclusion  $(p^*, x^*) \in N((\bar{p}, \bar{x}); \text{gph } \mathcal{F})$  implies that  $p^* \leq 0$  in the space  $ba(T)$ , i.e.,  $p^*(A) \leq 0$  for all  $A \subset T$ .

*Proof.* Observe from (2.6) and from the definition of the Dirac measure that the graph of  $\mathcal{F}$  admits the representation

$$\text{gph } \mathcal{F} = \{(p, x) \in l_\infty(T) \times X \mid \langle a_t^*, x \rangle - \langle \delta_t, p \rangle \leq b_t \text{ for all } t \in T\}.$$

Therefore, we have  $(p^*, x^*) \in N((\bar{p}, \bar{x}); \text{gph } \mathcal{F})$  if and only if

$$(3.2) \quad \langle p^*, p \rangle + \langle x^*, x \rangle \leq \langle p^*, \bar{p} \rangle + \langle x^*, \bar{x} \rangle$$

for every  $(p, x) \in \text{gph } \mathcal{F}$ . Employing now the equivalence between (i) and (ii) in Lemma 2.1, we conclude that  $(p^*, x^*) \in N((\bar{p}, \bar{x}); \text{gph } \mathcal{F})$  if and only if inclusion (3.1) holds.

To justify the last statement of the proposition, for every set  $A \subset T$  consider its characteristic function  $\chi_A: T \rightarrow \{0, 1\}$  defined by

$$\chi_A(t) := \begin{cases} 1 & \text{if } t \in A, \\ 0 & \text{if } t \notin A. \end{cases}$$

It is obvious that the inclusion  $(p, x) \in \text{gph } \mathcal{F}$  implies that  $(p + \lambda\chi_A, x) \in \text{gph } \mathcal{F}$  for each  $\lambda > 0$ . Replacing now in (3.2) the pair  $(p, x)$  by  $(p + \lambda\chi_A, x)$ , keeping  $p$  and  $x$  fixed, dividing both sides of the inequality by  $\lambda$ , and letting  $\lambda \rightarrow \infty$ , we get the relationships

$$\langle p^*, \chi_A \rangle = \int_T \chi_A(t) p^*(dt) = p^*(A) \leq 0,$$

which complete the proof of the proposition.  $\square$

Based on the above proposition and the general coderivative definition, we now obtain a constructive representation of the coderivative  $D^*\mathcal{F}(0, \bar{x})$  in question.

**THEOREM 3.2** (coderivative of the feasible solution map). *Let  $\bar{x} \in \mathcal{F}(0)$  for the feasible solution map  $\mathcal{F}: l_\infty(T) \rightrightarrows X$  from (1.1) with a Banach decision space  $X$ . Then  $p^* \in D^*\mathcal{F}(0, \bar{x})(x^*)$  if and only if*

$$(3.3) \quad (p^*, -x^*, -\langle x^*, \bar{x} \rangle) \in \text{cl}^* \text{cone}\{(-\delta_t, a_t^*, b_t) \mid t \in T\}.$$

*Proof.* By the coderivative construction (2.3) applied to  $\mathcal{F}$  and by the normal cone formula from Proposition 3.1 as  $\bar{p} = 0$ , we get that  $p^* \in D^*\mathcal{F}(0, \bar{x})(x^*)$  if and only if

$$(3.4) \quad (p^*, -x^*, -\langle x^*, \bar{x} \rangle) \in \text{cl}^* \text{cone}\left[\{(-\delta_t, a_t^*, b_t) \mid t \in T\} \cup \{(0, 0, 1)\}\right].$$

To justify the coderivative representation claimed in the theorem, we need to show that inclusion (3.4) implies, in fact, the “smaller” one in (3.3). Assuming indeed that (3.4) holds, we find by the structure of the right-hand side on (3.4) some nets  $\{\lambda_\nu\}_{\nu \in \mathcal{N}} \subset \mathbb{R}_+^{(T)}$  and  $\{\gamma_\nu\}_{\nu \in \mathcal{N}} \subset \mathbb{R}_+$  satisfying the limiting relationship

$$(3.5) \quad (p^*, -x^*, -\langle x^*, \bar{x} \rangle) = w^*\text{-}\lim_\nu \left( \sum_{t \in T} \lambda_{t\nu} (-\delta_t, a_t^*, b_t) + \gamma_\nu (0, 0, 1) \right),$$

where  $\lambda_{t\nu}$  stands for the  $t$ -entry of  $\lambda_\nu = (\lambda_{t\nu})_{t \in T}$  as  $\nu \in \mathcal{N}$ . It follows directly from the component structure in (3.5) that

$$(3.6) \quad 0 = \langle p^*, 0 \rangle + \langle -x^*, \bar{x} \rangle + (-\langle x^*, \bar{x} \rangle)(-1) = \lim_\nu \left( \sum_{t \in T} \lambda_{t\nu} (\langle a_t^*, \bar{x} \rangle - b_t) - \gamma_\nu \right).$$

Finally, taking into account the definition of the positive cone  $\mathbb{R}_+^{(T)}$  and the fact that the pair  $(0, \bar{x})$  satisfies the infinite inequality system in (1.1), we conclude from (3.6) that  $\lim_\nu \gamma_\nu = 0$ . This justifies (3.3) and thus completes the proof of the theorem.  $\square$

The next consequence of Theorem 3.2 is useful in what follows.

**COROLLARY 3.3** (limiting descriptions of coderivatives). *If  $p^* \in D^*\mathcal{F}(0, \bar{x})(x^*)$  in the framework of Theorem 3.2, then there is a net  $\{\lambda_\nu\}_{\nu \in \mathcal{N}} \subset \mathbb{R}_+^{(T)}$  such that*

$$\begin{aligned} \sum_{t \in T} \lambda_{t\nu} &\rightarrow \|p^*\| = -\langle p^*, e \rangle, \\ \sum_{t \in T} \lambda_{t\nu} a_t^* &\xrightarrow{w^*} -x^*, \quad \text{and} \quad \sum_{t \in T} \lambda_{t\nu} b_t \rightarrow -\langle x^*, \bar{x} \rangle. \end{aligned}$$

*Proof.* It follows from Theorem 3.2 that there is a net  $\{\lambda_\nu\}_{\nu \in \mathcal{N}} \subset \mathbb{R}_+^{(T)}$  such that

$$\sum_{t \in T} \lambda_{t\nu} \delta_t \xrightarrow{w^*} -p^*, \quad \sum_{t \in T} \lambda_{t\nu} a_t^* \xrightarrow{w^*} -x^*, \quad \text{and} \quad \sum_{t \in T} \lambda_{t\nu} b_t \rightarrow -\langle x^*, \bar{x} \rangle,$$

which readily imply the relationships

$$\left\langle \sum_{t \in T} \lambda_{t\nu} \delta_t, e \right\rangle = \sum_{t \in T} \lambda_{t\nu} \rightarrow \langle p^*, -e \rangle =: \lambda \in [0, \infty).$$

Since the dual norm on  $X^*$  is  $w^*$ -lower semicontinuous, we have

$$\|p^*\| \leq \liminf_{\nu \in \mathcal{N}} \left\| \sum_{t \in T} \lambda_{t\nu} \delta_t \right\| \leq \liminf_{\nu \in \mathcal{N}} \sum_{t \in T} \lambda_{t\nu} = \lambda.$$

It follows at the same time from the norm definition that

$$\|p^*\| = \sup_{\|p\| \leq 1} \langle p^*, p \rangle \geq \langle p^*, -e \rangle = \lambda,$$

which finally yields  $\|p^*\| = -\langle p^*, e \rangle$  and completes the proof of the corollary.  $\square$

Our further intention is to provide the *exact calculation* of the *coderivative norm*

$$(3.7) \quad \|D^* \mathcal{F}(0, \bar{x})\| := \sup \{ \|p^*\| \mid p^* \in D^* \mathcal{F}(0, \bar{x})(x^*), \|x^*\| \leq 1 \}$$

in terms of the initial data of the linear infinite inequality system (1.1). A part of our analysis in this direction is the following lemma on properties of the characteristic set (2.8) at  $p = 0$ , which is also used in section 4 to compute the exact Lipschitzian bound  $\text{lip } \mathcal{F}(0, \bar{x})$ .

LEMMA 3.4 (properties of the characteristic set). *Let  $X$  be an arbitrary Banach space. The following assertions hold:*

(i) *Assume that  $\bar{x} \in \mathcal{F}(0)$  is not a strong Slater point for the infinite system (1.1) at  $p = 0$  and that the collection  $\{a_t^* \mid t \in T\}$  is bounded in  $X^*$ . Then the set*

$$(3.8) \quad S := \{x^* \in X^* \mid (x^*, \langle x^*, \bar{x} \rangle) \in \text{cl}^* C(0)\}$$

*built upon the characteristic set  $C(0)$  in (2.8) is nonempty and  $w^*$ -compact at  $X^*$ .*

(ii) *Assume that  $\bar{x} \in \mathcal{F}(0)$  is a strong Slater point of (1.1) at  $p = 0$ . Then the set  $S$  in (3.8) is empty.*

*Proof.* To justify (i), let  $\bar{x}$  be not a strong Slater point for the infinite system (1.1) at  $p = 0$ . Then there is a sequence  $\{t_k\}_{k \in \mathbb{N}} \subset T$  such that  $\lim_k (\langle a_{t_k}^*, \bar{x} \rangle - b_{t_k}) = 0$ . The boundedness of  $\{a_t^* \mid t \in T\}$  implies by the classical Alaoglu–Bourbaki theorem that this set is relatively  $w^*$ -compact in  $X^*$ ; i.e., there is a *subnet*  $\{a_{t_\nu}^*\}_{\nu \in \mathcal{N}}$  of the latter sequence that  $w^*$ -converges to some element  $u^* \in \text{cl}^* \{a_t^* \mid t \in T\}$ . This gives  $\lim_\nu b_{t_\nu} = \langle u^*, \bar{x} \rangle$ , and therefore

$$(u^*, \langle u^*, \bar{x} \rangle) = w^*\text{-}\lim_\nu (a_{t_\nu}^*, b_{t_\nu}) \in \text{cl}^* C(0),$$

which justifies the *nonemptiness* of the set  $S$  in (3.8). Next we prove that  $S$  is  $w^*$ -compact.

Indeed, by our assumption, the set  $A := \{a_t^* \mid t \in T\}$  is bounded in  $X^*$ , and so is  $\text{cl}^* \text{co } A$ ; the latter is actually  $w^*$ -compact due to its automatic  $w^*$ -closedness. Observe

further that the set  $S$  in (3.8) is a *preimage* of  $\text{cl}^*C(0)$  under the  $w^*$ -continuous mapping  $x^* \mapsto (x^*, \langle x^*, \bar{x} \rangle)$ , and thus it is  $w^*$ -closed in  $X^*$ . Since  $S$  is a subset of  $\text{cl}^* \text{co} A$ , it is also bounded and hence  $w^*$ -compact in  $X^*$ . This completes the proof of assertion (i).

To prove assertion (ii), let  $\bar{x}$  be a strong Slater point of system (1.1) at  $p = 0$ , and let

$$\gamma := - \sup_{t \in T} \{ \langle a_t^*, \bar{x} \rangle - b_t \}.$$

Then we have the inequality

$$\langle x^*, \bar{x} \rangle \leq \beta - \gamma \quad \text{whenever} \quad (x^*, \beta) \in \text{cl}^*C(0),$$

which justifies (ii) and thus completes the proof of the lemma.  $\square$

Now we are ready to compute the coderivative norm  $\|D^*\mathcal{F}(0, \bar{x})\|$  at the reference point.

**THEOREM 3.5** (computing the coderivative norm). *Let  $\bar{x} \in \text{dom } \mathcal{F}$  for the infinite system (1.1) with an arbitrary Banach space  $X$  of decision variables. Assume that  $\mathcal{F}$  satisfies the strong Slater condition at  $p = 0$  and that the coefficient set  $\{a_t^* \mid t \in T\}$  is bounded in  $X^*$ . The following assertions hold:*

(i) *If  $\bar{x}$  is a strong Slater point for  $\mathcal{F}$  at  $p = 0$ , then  $\|D^*\mathcal{F}(0, \bar{x})\| = 0$ .*

(ii) *If  $\bar{x}$  is not a strong Slater point for  $\mathcal{F}$  at  $p = 0$ , then the coderivative norm (3.7) is positive and is computed by*

$$(3.9) \quad \|D^*\mathcal{F}(0, \bar{x})\| = \max \left\{ \|x^*\|^{-1} \mid (x^*, \langle x^*, \bar{x} \rangle) \in \text{cl}^*C(0) \right\}$$

via the  $w^*$ -closure of the characteristic set (2.8) at  $p = 0$ .

*Proof.* To justify assertion (i), assume that  $\bar{x}$  is a strong Slater point for the system  $\mathcal{F}$  at  $p = 0$ . It follows from the proof of implication (i) $\implies$ (v) in Lemma 2.3 that we have  $(0, \bar{x}) \in \text{int}(\text{gph } \mathcal{F})$ . It gives

$$N((0, \bar{x}); \text{gph } \mathcal{F}) = \{(0, 0)\},$$

and the conclusion in (i) follows from (2.3) and (3.7).

To prove assertion (ii), take  $x^* \in X^*$  such that  $(x^*, \langle x^*, \bar{x} \rangle) \in \text{cl}^*C(0)$ ; the latter set is nonempty according to Lemma 3.4. Then there exists a net  $\{\lambda_\nu\}_{\nu \in \mathcal{N}} \subset \mathbb{R}_+^{(T)}$  with  $\sum_{t \in T} \lambda_{t\nu} = 1$  for all  $\nu \in \mathcal{N}$  such that

$$\sum_{t \in T} \lambda_{t\nu} a_t^* \xrightarrow{w^*} x^* \quad \text{and} \quad \sum_{t \in T} \lambda_{t\nu} b_t \rightarrow \langle x^*, \bar{x} \rangle.$$

Form further net elements  $p_\nu^* \in l_\infty(T)^*$  by

$$p_\nu^* := - \sum_{t \in T} \lambda_{t\nu} \delta_t, \quad \nu \in \mathcal{N}.$$

Since  $\|p_\nu^*\| = \langle p_\nu^*, -e \rangle = 1$ , the Alaoglu–Bourbaki theorem allows us to select a subnet (without relabeling) such that  $p_\nu^* \xrightarrow{w^*} p^*$  with  $\|p^*\| \leq 1$ . Following the same reasoning as in the proof of Corollary 3.3, we conclude that

$$(3.10) \quad 1 = \lim_{\nu \in \mathcal{N}} \sum_{t \in T} \lambda_{t\nu} = \|p^*\| = \langle p^*, -e \rangle.$$

Moreover, we also obtain the inclusion

$$(p^*, x^*, \langle x^*, \bar{x} \rangle) \in \text{cl}^* \text{co} \{(-\delta_t, a_t^*, b_t) \mid t \in T\},$$

which implies by Theorem 3.2 that

$$(3.11) \quad p^* \in D^* \mathcal{F}(0, \bar{x})(-x^*).$$

Suppose now that  $x^* = 0$  in (3.11). Since  $p^* \neq 0$  by (3.10), we get from (3.11) that

$$(3.12) \quad D^* \mathcal{F}(0, \bar{x})(0) \neq \{0\},$$

which yields by [31, Theorem 1.44] that  $\mathcal{F}$  is *not Lipschitz-like* around  $(0, \bar{x})$ , and therefore it cannot satisfy the strong Slater condition by implication (i)  $\implies$  (iv) in Lemma 2.3. This contradicts the assumption of the theorem.

Thus  $x^* \neq 0$  in (3.11), and we derive from the latter relationship that

$$\|x^*\|^{-1} p^* \in D^* \mathcal{F}(0, \bar{x}) \left( -\|x^*\|^{-1} x^* \right),$$

which gives in turn that

$$\|D^* \mathcal{F}(0, \bar{x})\| \geq \left\| \|x^*\|^{-1} p^* \right\| = \|x^*\|^{-1}$$

and hence justifies the inequality “ $\geq$ ” in (3.9).

It remains to prove the opposite inequality in (3.9). For the nonempty and  $w^*$ -compact set  $S$  in (3.8) we have  $0 \notin S$  by Lemma 2.3, which ensures the  $w^*$ -upper semicontinuity of the function  $x^* \mapsto \|x^*\|^{-1}$  on  $S$ . Thus the supremum in the right-hand side of (3.9) is attained and belongs to  $(0, \infty)$ . Then condition (v) in Lemma 2.3 implies that  $(0, \hat{x}) \in \text{int}(\text{gph } \mathcal{F})$  for some  $\hat{x} \in X$ , and so  $0 \in \text{int}(\text{dom } \mathcal{F})$ . Moreover, we have that  $p^* \in D^* \mathcal{F}(0, \bar{x})(-x^*)$  if and only if  $(p^*, x^*) \in N((0, \bar{x}); \text{gph } \mathcal{F})$ , which is equivalent to

$$(3.13) \quad \langle p^*, p \rangle + \langle x^*, x \rangle \leq \langle x^*, \bar{x} \rangle \quad \text{for all } (p, x) \in \text{gph } \mathcal{F}.$$

This allows us, by taking into account that  $0 \in \text{int}(\text{dom } \mathcal{F})$ , to arrive at the equivalences

$$(3.14) \quad p^* \in D^* \mathcal{F}(0, \bar{x})(0) \iff \langle p^*, p \rangle \leq 0 \text{ for all } p \in \text{dom } \mathcal{F} \iff p^* = 0.$$

Observe, furthermore, that, since  $\bar{x}$  is not a strong Slater point for  $\mathcal{F}$  at  $p = 0$ , we have  $(0, \bar{x}) \notin \text{int}(\text{gph } \mathcal{F})$  and thus conclude by the classical separation theorem that there is a pair  $(p^*, x^*) \neq (0, 0)$  for which relation (3.13) holds. Employing (3.14), we have  $x^* \neq 0$  and  $p^* \in D^* \mathcal{F}(0, \bar{x})(-x^*)$ .

Take now  $p^* \in D^* \mathcal{F}(0, \bar{x})(-x^*)$  with  $\|x^*\| \leq 1$  and suppose that  $x^* \neq 0$ ; the arguments of the previous paragraph ensure the existence of such an element. By Corollary 3.3 there is a net  $\{\lambda_\nu\}_{\nu \in \mathcal{N}} \subset \mathbb{R}_+^{(T)}$  for which

$$\begin{aligned} \gamma_\nu &:= \sum_{t \in T} \lambda_{t\nu} \rightarrow \|p^*\| = -\langle p^*, e \rangle, \\ x_\nu^* &:= \sum_{t \in T} \lambda_{t\nu} a_t^* \xrightarrow{w^*} x^*, \quad \text{and} \quad \sum_{t \in T} \lambda_{t\nu} b_t \rightarrow \langle x^*, \bar{x} \rangle. \end{aligned}$$

Taking  $M \geq \|a_t^*\|$  for every  $t \in T$ , we get the estimate

$$\|x_\nu^*\| \leq M\gamma_\nu \quad \text{whenever } \nu \in \mathcal{N}$$

and also the limiting relationships

$$0 < \|x^*\| \leq \liminf_{\nu \in \mathcal{N}} \|x_\nu^*\| \leq M \liminf_{\nu \in \mathcal{N}} \gamma_\nu = M \|p^*\|,$$

which ensure that  $p^* \neq 0$ . It follows, furthermore, that

$$\|p^*\|^{-1} (x^*, \langle x^*, \bar{x} \rangle) \in \text{cl}^* C(0).$$

Remembering, finally, that  $0 < \|x^*\| \leq 1$ , we arrive at the estimates

$$\|p^*\| \leq \left\| \|p^*\|^{-1} x^* \right\|^{-1} \leq \max \left\{ \|u^*\|^{-1} \mid (u^*, \langle u^*, \bar{x} \rangle) \in \text{cl}^* C(0) \right\},$$

which justify the inequality “ $\leq$ ” in (3.9) and complete the proof of the theorem.  $\square$

**4. Characterizations of robust Lipschitzian stability for feasible solution maps.** In this section we employ the above coderivative analysis combined with appropriate techniques developed in linear semi-infinite/infinite programming to establish a *coderivative characterization* of robust Lipschitzian stability, in the sense discussed in section 2, for the infinite inequality system  $\mathcal{F}$  in (1.1) at the reference point  $(0, \bar{x})$  with computing the *exact Lipschitzian bound*  $\text{lip } \mathcal{F}(0, \bar{x})$ .

The first result of this section establishes the coderivative *necessary and sufficient* condition in form (2.4) for the Lipschitz-like property of  $\mathcal{F}$  around  $(0, \bar{x}) \in \text{gph } \mathcal{F}$  in the general setting under consideration.

**THEOREM 4.1** (coderivative criterion for robust Lipschitzian stability of linear infinite inequalities). *Let  $\bar{x} \in \mathcal{F}(0)$  for the infinite inequality system (1.1) with a Banach space  $X$  of decision variables. Then  $\mathcal{F}$  is Lipschitz-like around  $(0, \bar{x})$  if and only if*

$$(4.1) \quad D^* \mathcal{F}(0, \bar{x})(0) = \{0\}.$$

*Proof.* The “only if” part follows from [31, Theorem 1.44] specified for the mapping  $\mathcal{F}: l_\infty(T) \rightrightarrows X$  under consideration. Let us now prove the “if” part of the theorem.

Arguing by contradiction, suppose that  $D^* \mathcal{F}(0, \bar{x})(0) = \{0\}$  while the mapping  $\mathcal{F}$  is *not* Lipschitz-like around  $(0, \bar{x})$ . Then, by the equivalence between properties (ii) and (iv) of Lemma 2.3, we get the inclusion

$$(0, 0) \in \text{cl}^* \text{co} \{ (a_t^*, b_t) \in X^* \times \mathbb{R} \mid t \in T \},$$

which means that there is a net  $\{\lambda_\nu\}_{\nu \in \mathcal{N}} \in \mathbb{R}_+^{(T)}$  such that  $\sum_{t \in T} \lambda_{t\nu} = 1$  for all  $\nu \in \mathcal{N}$  and

$$(4.2) \quad w^* \text{-} \lim_{\nu} \sum_{t \in T} \lambda_{t\nu} (a_t^*, b_t) = (0, 0).$$

Since the net  $\{\sum_{t \in T} \lambda_{t\nu} (-\delta_t)\}_{\nu \in \mathcal{N}}$  is obviously bounded in  $l_\infty(T)^*$ , the Alaoglu–Bourbaki theorem ensures the existence of its subnet (with no relabeling) that  $w^*$ -converges to some element  $p^* \in l_\infty(T)^*$ , i.e.,

$$(4.3) \quad p^* = w^* \text{-} \lim_{\nu} \sum_{t \in T} \lambda_{t\nu} (-\delta_t).$$

It follows from (4.3) by the Dirac function definition that

$$\langle p^*, -e \rangle = \lim_{\nu \in \mathcal{N}} \sum_{t \in T} \lambda_{t\nu} = 1, \text{ where } e = (e_t)_{t \in T} \text{ with } e_t = 1 \text{ for all } t \in T,$$

which implies that  $p^* \neq 0$ . It follows from (4.2) and (4.3) that

$$(p^*, 0, 0) = w^* \text{-}\lim_{\nu} \sum_{t \in T} \lambda_{t\nu} (-\delta_t, a_t^*, b_t) \text{ with } p^* \neq 0,$$

and therefore, by the explicit coderivative description of Theorem 3.2, we get the inclusion  $p^* \in D^* \mathcal{F}(0, \bar{x})(0) \setminus \{0\}$ , which contradicts the assumed condition (4.1). This justifies the sufficiency of the coderivative condition (4.1) for the Lipschitz-like property of  $\mathcal{F}$  around  $(0, \bar{x})$  and thus completes the proof of the theorem.  $\square$

Our further goal is to compute the *exact Lipschitzian bound*  $\text{lip } \mathcal{F}(0, \bar{x})$  of  $\mathcal{F}$  around  $(0, \bar{x})$ . We are going to do it according to the *distance representation* (2.2) for the exact Lipschitzian bound. To proceed, denote

$$(4.4) \quad H(x^*, \alpha) := \{x \in X \mid \langle x^*, x \rangle \leq \alpha\} \text{ for } (x^*, \alpha) \in X^* \times \mathbb{R},$$

and observe the following representation (known as the *Ascoli formula*; see, e.g., [2]):

$$(4.5) \quad \text{dist}(x; H(x^*, \alpha)) = \frac{[\langle x^*, x \rangle - \alpha]_+}{\|x^*\|},$$

where  $[\gamma]_+ := \max\{\gamma, 0\}$  for  $\gamma \in \mathbb{R}$ . Recall that we are under the convention  $0/0 := 0$ .

The next proposition and the subsequent lemma, which are certainly of their own interest, provide a significant extension of the Ascoli formula (4.4) to the case of *infinite* systems of linear inequalities instead of the single one as in (4.4). These results are essentially employed in what follows for computing the exact Lipschitzian bound  $\text{lip } \mathcal{F}(0, \bar{x})$ . We refer the reader to [6, Lemma 2.3] and [7, Lemma 1] for related results in the framework of semi-infinite programming and observe that in infinite dimensions we use the *w\*-closure* of the characteristic sets  $C(p)$  from (2.8); see also Example 4.4 below for more discussion.

From now on, given an extended-real-valued function  $f: X \rightarrow \overline{\mathbb{R}} := (-\infty, \infty]$  assumed to be *proper* (i.e., not identically equal to  $\infty$ ), consider its *Fenchel conjugate*  $f^*: X^* \rightarrow \overline{\mathbb{R}}$  defined by

$$f^*(x^*) := \sup \{ \langle x^*, x \rangle - f(x) \mid x \in X \} = \sup \{ \langle x^*, x \rangle - f(x) \mid x \in \text{dom } f \},$$

where  $\text{dom } f := \{x \in X \mid f(x) < \infty\}$  is the *effective domain* of  $f$ . In what follows we also use the standard notation  $\text{epi } f$  for the *epigraph* of  $f$  given by

$$\text{epi } f := \{(x, \gamma) \in X \times \mathbb{R} \mid x \in \text{dom } f, f(x) \leq \gamma\}.$$

The following result gives an *exact formula* for computing the distance to a set defined by a convex inequality via the corresponding conjugate function.

**PROPOSITION 4.2** (distance function for solutions of convex inequalities in Banach spaces). *Let  $g: X \rightarrow \overline{\mathbb{R}}$  be a proper convex function on a Banach space  $X$ , and let*

$$(4.6) \quad Q := \{y \in X \mid g(y) \leq 0\}.$$

Assume the fulfillment of the classical Slater condition: there is  $\hat{x} \in X$  such that  $g(\hat{x}) < 0$ . Then the distance function to the set  $Q$  in (4.6) is computed by

$$(4.7) \quad \text{dist}(x; Q) = \max_{(x^*, \alpha) \in \text{epi } g^*} \frac{[\langle x^*, x \rangle - \alpha]_+}{\|x^*\|}.$$

*Proof.* Observe that the nonemptiness of  $Q$  in (4.6) yields that  $\alpha \geq 0$  whenever  $(0, \alpha) \in \text{epi } g^*$  and that the possibility of  $x^* = 0$  is not an obstacle in (4.7) under our convention  $0/0 = 0$ . Obviously the distance function  $\text{dist}(x; Q)$  is nothing else but the optimal value function in the parametric *convex optimization* problem

$$(4.8) \quad \text{minimize } \|y - x\| \quad \text{subject to } g(y) \leq 0.$$

Since the classical Slater condition holds for problem (4.8) by our assumption, we have the *strong Lagrange duality* in (4.8) by, e.g., [39, Theorem 2.9.3], which gives

$$\begin{aligned} \text{dist}(x; Q) &= \max_{\lambda \geq 0} \inf_{y \in X} \{ \|y - x\| + \lambda g(y) \} \\ &= \max \left\{ \max_{\lambda > 0} \inf_{y \in X} \{ \|y - x\| + \lambda g(y) \}, \inf_{y \in X} \|y - x\| \right\} \\ &= \max \left\{ \max_{\lambda > 0} \inf_{y \in X} \{ \|y - x\| + \lambda g(y) \}, 0 \right\}. \end{aligned}$$

Applying the classical *Fenchel duality theorem* to the inner infimum problem above for a fixed  $\lambda > 0$  (observing to this end that Rockafellar’s regularity condition needed for Fenchel duality is satisfied for this problem due to the continuity of the norm function), we get

$$(4.9) \quad \inf_{y \in X} \{ \|y - x\| + \lambda g(y) \} = \max_{y^* \in X^*} \{ -\|\cdot - x\|^*(-y^*) - (\lambda g)^*(y^*) \}.$$

It is well known in convex analysis that

$$\|\cdot - x\|^*(-y^*) = \begin{cases} \langle -y^*, x \rangle & \text{if } \|y^*\| \leq 1, \\ \infty & \text{otherwise.} \end{cases}$$

Implementing this into formula (4.9) and employing elementary transformations, we obtain

$$\begin{aligned} \inf_{y \in X} \{ \|y - x\| + \lambda g(y) \} &= \max_{\|y^*\| \leq 1} \{ \langle y^*, x \rangle - (\lambda g)^*(y^*) \} \\ &= \max_{\|y^*\| \leq 1, (\lambda g)^*(y^*) \leq \eta} \{ \langle y^*, x \rangle - \eta \} \\ &= \max_{\|y^*\| \leq 1, \lambda g^*(y^*/\lambda) \leq \eta} \{ \langle y^*, x \rangle - \eta \} \\ &= \max_{\|y^*\| \leq 1, (1/\lambda)(y^*, \eta) \in \text{epi } g^*} \{ \langle y^*, x \rangle - \eta \}. \end{aligned}$$

The latter representation yields, by denoting  $x^* := (1/\lambda)y^*$  and  $\alpha := (1/\lambda)\eta$ , that

$$\inf_{y \in X} \{ \|y - x\| + \lambda g(y) \} = \max_{(x^*, \alpha) \in \text{epi } g^*, \|x^*\| \leq 1/\lambda} \lambda \{ \langle x^*, x \rangle - \alpha \}.$$

Combining this with the formulas above, we arrive at

$$(4.10) \quad \begin{aligned} \text{dist}(x; Q) &= \max \left\{ \max_{(x^*, \alpha) \in \text{epi } g^*, \|x^*\| \leq 1/\lambda} \lambda \{ \langle x^*, x \rangle - \alpha \}, 0 \right\} \\ &= \max_{(x^*, \alpha) \in \text{epi } g^*, \|x^*\| \leq 1/\lambda} \left\{ \lambda [\langle x^*, x \rangle - \alpha]_+ \right\}. \end{aligned}$$



It is easy to observe the following relationships held for any  $\lambda > 0$ :

$$\begin{aligned} \max_{(0,\alpha) \in \text{epi } g^*} \lambda \{ \langle 0, x \rangle - \alpha \} &= \max_{g^*(0) \leq \alpha} \lambda (\langle 0, x \rangle - \alpha) \\ &= \lambda (-g^*(0)) \leq \lambda \inf_{x \in X} g(x) \leq \lambda g(\hat{x}) < 0. \end{aligned}$$

Taking this into account, we have from (4.10) that

$$\begin{aligned} \text{dist}(x; Q) &= \max_{(x^*, \alpha) \in \text{epi } g^*, \|x^*\| \leq 1/\lambda} \left\{ \lambda [\langle x^*, x \rangle - \alpha]_+ \right\} \\ &= \max_{(x^*, \alpha) \in \text{epi } g^*} \max_{\|x^*\| \leq 1/\lambda} \left\{ \lambda [\langle x^*, x \rangle - \alpha]_+ \right\} \\ &= \max_{(x^*, \alpha) \in \text{epi } g^*} \frac{[\langle x^*, x \rangle - \alpha]_+}{\|x^*\|}, \end{aligned}$$

which gives (4.7) and thus completes the proof of the proposition.  $\square$

Now we are ready we establish the required extension of the Ascoli formula (4.5) to the case of the infinite inequality systems under consideration.

**LEMMA 4.3** (distance to infinite linear inequalities in Banach spaces). *Assume that the infinite system  $\mathcal{F}(p)$  in (1.1) satisfies the strong Slater condition at  $p = (p_t)_{t \in T}$ . Then for any  $x \in X$  and any  $p \in l_\infty(T)$  we have the extended Ascoli distance formula*

$$(4.11) \quad \text{dist}(x; \mathcal{F}(p)) = \max_{(x^*, \alpha) \in \text{cl}^* C(p)} \frac{[\langle x^*, x \rangle - \alpha]_+}{\|x^*\|}.$$

If, furthermore,  $X$  is reflexive, then the distance formula (4.11) can be simplified as follows:

$$(4.12) \quad \text{dist}(x; \mathcal{F}(p)) = \sup_{(x^*, \alpha) \in C(p)} \frac{[\langle x^*, x \rangle - \alpha]_+}{\|x^*\|}.$$

*Proof.* It is obvious to observe that the infinite system (1.1) is represented in

$$(4.13) \quad \mathcal{F}(p) = \{x \in X \mid g(x) \leq 0\},$$

where the convex function  $g: X \rightarrow \overline{\mathbb{R}}$  is given in the supremum form

$$(4.14) \quad g(x) := \sup_{t \in T} (f_t(x) - p_t) \quad \text{with} \quad f_t(x) := \langle a_t^*, x \rangle - b_t.$$

The assumed strong Slater condition for  $\mathcal{F}(p)$  ensures the fulfillment of the classical Slater condition for  $g$  imposed in Proposition 4.2. To imply this proposition in the framework of (4.13), we need to compute the conjugate function to the supremum function in (4.14). The recent results in this direction [4, 18] yield that

$$\text{epi } g^* = \text{epi} \left\{ \sup_{t \in T} (f_t - p_t) \right\}^* = \text{cl}^* \text{co} \left( \bigcup_{t \in T} \text{epi} (f_t - p_t)^* \right) = \text{cl}^* C(p) + \mathbb{R}_+(0, 1)$$

with  $0 \in X^*$ , where the weak\* closedness of the set  $\text{cl}^* C(p) + \mathbb{R}_+(0, 1)$  is a consequence of the classical Dieudonné theorem; see, e.g., [39, Theorem 1.1.8]. Thus we get the distance formula (4.11) from Proposition 4.2 in general Banach spaces.

To prove the simplified distance formula (4.12) in the case of reflexive spaces, suppose, on the contrary, that it does not hold. Then there is a scalar  $\beta \in \mathbb{R}$  such that

$$(4.15) \quad \max_{(x^*, \alpha) \in \text{cl}^* C(p)} \frac{[\langle x^*, x \rangle - \alpha]_+}{\|x^*\|} > \beta > \sup_{(x^*, \alpha) \in C(p)} \frac{[\langle x^*, x \rangle - \alpha]_+}{\|x^*\|}.$$

This yields the existence of a pair  $(\bar{x}^*, \bar{\alpha}) \in \text{cl}^* C(p)$  with  $\bar{x}^* \in X^* \setminus \{0\}$  and  $\bar{\alpha} \in \mathbb{R}$  such that

$$\frac{[\langle \bar{x}^*, x \rangle - \bar{\alpha}]_+}{\|\bar{x}^*\|} > \beta.$$

Taking into account that the space  $X$  is reflexive and that the set  $C(p)$  is convex and employing the Mazur weak closure theorem, we can replace the weak\* closure of the  $C(p)$  above by its norm closure in  $X^*$ . This allows us to find a sequence  $(x_k^*, \alpha_k) \in C(p)$ ,  $k \in \mathbb{N}$ , converging in norm to  $(\bar{x}^*, \bar{\alpha})$  as  $k \rightarrow \infty$ . Thus we get

$$\lim_{k \rightarrow \infty} \frac{[\langle x_k^*, x \rangle - \alpha_k]_+}{\|x_k^*\|} = \frac{[\langle \bar{x}^*, x \rangle - \bar{\alpha}]_+}{\|\bar{x}^*\|} > \beta,$$

and therefore there is  $k_0 \in \mathbb{N}$  such that

$$\frac{[\langle x_{k_0}^*, x \rangle - \alpha_{k_0}]_+}{\|x_{k_0}^*\|} > \beta.$$

The latter surely contradicts (4.15), and this completes the proof of the lemma.  $\square$

The following example shows that the *reflexivity* of the decision space  $X$  is an *essential* requirement for the validity of the simplified distance formula (4.12), even in the framework of (nonreflexive) *Asplund* spaces.

*Example 4.4* (failure of simplified distance formula in nonreflexive Asplund spaces). Consider the classical space  $c_0$  of sequences of real numbers converging to zero endowed with the supremum norm. This space is well known to be *Asplund* while *not reflexive*; see, e.g., [17]. Let us show that the simplified distance formula (4.12) *fails* in  $X = c_0$  for a rather plain linear system of *countable* inequalities. Of course, we need to demonstrate that the inequality “ $\leq$ ” is generally violated in (4.12), since the opposite inequality holds in any Banach space. Form the infinite (countable) linear inequality system

$$(4.16) \quad \mathcal{F}(0) := \{x \in c_0 \mid \langle e_1^* + e_t^*, x \rangle \leq -1, t \in \mathbb{N}\},$$

where  $e_t^* \in l_1$  has 1 as its  $t$ th component while all the remaining components are 0. System (4.16) can be rewritten as

$$x \in \mathcal{F}(0) \iff x(1) + x(t) \leq -1 \text{ for all } t \in \mathbb{N}.$$

Observe that for the origin  $z = 0$  we have  $\text{dist}(0; \mathcal{F}(0)) = 1$ , and the distance is realized at, e.g.,  $u = (-1, 0, 0, \dots)$ . Indeed, passing to the limit in the inequality

$$x(1) + x(t) \leq -1 \text{ as } t \rightarrow \infty$$

and taking into account that  $x(t) \rightarrow 0$  as  $t \rightarrow \infty$ , by the structure of the space of  $c_0$ , we get  $x(1) \leq -1$ . Furthermore, it can be checked that

$$\begin{aligned} (e_1^*, -1) \in \text{cl}^*C(0), \quad \langle e_1^*, x - u \rangle \leq 0 \quad \text{for all } x \in \mathcal{F}(0), \\ \text{dist}(z; \mathcal{F}(0)) = \|z - u\| = \langle e_1^*, z - u \rangle = \frac{\langle e_1^*, z \rangle - (-1)}{\|e_1^*\|}. \end{aligned}$$

On the other hand, for the pair  $(x^*, \alpha) \in X^* \times \mathbb{R}$  given by

$$(x^*, \alpha) := \left( e_1^* + \sum_{t \in \mathbb{N}} \lambda_t e_t^*, -1 \right) \in C(0) \quad \text{with } \lambda \in \mathbb{R}_+^{(\mathbb{N})} \quad \text{and} \quad \sum_{t \in \mathbb{N}} \lambda_t = 1,$$

we can directly check that  $\|x^*\| = 2$  and hence

$$\frac{[\langle x^*, z \rangle - \alpha]_+}{\|x^*\|} = \frac{1}{2},$$

which shows that the equality in (4.12) is violated for the countable system (4.16) in the nonreflexive Asplund space  $X = c_0$  of decision variables.

Our next step is to derive a verifiable precise formula for computing the *exact Lipschitzian bound*  $\text{lip } \mathcal{F}(0, \bar{x})$  for the infinite system (1.1) in the general Banach space  $X$ . As a preliminary result we need the following technical lemma.

LEMMA 4.5 (closed-graph property of the characteristic set mapping). *The set-valued mapping  $l_\infty(T) \ni p \mapsto \text{cl}^*C(p) \subset X^* \times \mathbb{R}$  is closed-graph in the norm  $\times$  weak\* topology of  $l_\infty(T) \times (X^* \times \mathbb{R})$ ; i.e., for any nets*

$$\{p_\nu\}_{\nu \in \mathcal{N}} \subset l_\infty(T), \quad \{x_\nu^*\}_{\nu \in \mathcal{N}} \subset X^*, \quad \{\beta_\nu\}_{\nu \in \mathcal{N}} \subset \mathbb{R}$$

satisfying  $p_\nu \rightarrow p$ ,  $x_\nu^* \xrightarrow{w^*} x^*$ ,  $\beta_\nu \rightarrow \beta$ , and  $(x_\nu^*, \beta_\nu) \in \text{cl}^*C(p_\nu)$  for every  $\nu \in \mathcal{N}$  we have the inclusion  $(x^*, \beta) \in \text{cl}^*C(p)$ .

*Proof.* Arguing by contradiction, suppose that  $(x^*, \beta) \notin \text{cl}^*C(p)$ . Then the classical strict separation theorem allows us to find a pair  $(x, \alpha) \in X \times \mathbb{R}$  with  $(x, \alpha) \neq (0, 0)$  and real numbers  $\gamma$  and  $\gamma'$  satisfying

$$\langle x^*, x \rangle + \beta \alpha < \gamma' < \gamma \leq \langle a_t^*, x \rangle + (b_t + p_t) \alpha \quad \text{for all } t \in T.$$

Hence there exists a net index  $\nu_0 \in \mathcal{N}$  such that

$$\langle x_\nu^*, x \rangle + \beta_\nu \alpha < \gamma' \quad \text{and} \quad \|\alpha(p - p_\nu)\| \leq \gamma - \gamma' \quad \text{whenever } \nu \succeq \nu_0.$$

We have, therefore, that

$$\begin{aligned} \langle a_t^*, x \rangle + \alpha(b_t + p_{t\nu}) &= \langle a_t^*, x \rangle + \alpha(b_t + p_t) + \alpha(p_{t\nu} - p_t) \\ &\geq \gamma - \|\alpha(p_\nu - p)\| \geq \gamma' \quad \text{for all } t \in T. \end{aligned}$$

The latter implies that  $\gamma' \leq \langle z^*, x \rangle + \eta \alpha$  for all  $(z^*, \eta) \in \text{cl}^*C(p_\nu)$  whenever  $\nu \succeq \nu_0$ . Thus we arrive at the contradiction

$$\langle x_\nu^*, x \rangle + \beta_\nu \alpha < \gamma' \leq \langle x_\nu^*, x \rangle + \beta_\nu \alpha, \quad \nu \succeq \nu_0,$$

which completes the proof of the lemma.  $\square$

Now we are ready to establish the desired formula for computing the exact Lipschitzian bound of  $\mathcal{F}$  around  $(0, \bar{x})$ .

**THEOREM 4.6** (computing the exact Lipschitzian bound). *Let  $\bar{x} \in \mathcal{F}(0)$  for the linear infinite inequality system (1.1) with a Banach decision space  $X$ . Assume that  $\mathcal{F}$  satisfies the strong Slater condition at  $p = 0$  and that the coefficient set  $\{a_t^* \mid t \in T\}$  is bounded in  $X^*$ . The following assertions hold:*

(i) *If  $\bar{x}$  is a strong Slater point for  $\mathcal{F}$  at  $p = 0$ , then  $\text{lip } \mathcal{F}(0, \bar{x}) = 0$ .*

(ii) *If  $\bar{x}$  is not a strong Slater point for  $\mathcal{F}$  at  $p = 0$ , then the exact Lipschitzian bound of  $\mathcal{F}$  around  $(0, \bar{x})$  is computed by*

$$(4.17) \quad \text{lip } \mathcal{F}(0, \bar{x}) = \max \{ \|x^*\|^{-1} \mid (x^*, \langle x^*, \bar{x} \rangle) \in \text{cl}^* C(0) \} > 0$$

via the  $w^*$ -closure of the characteristic set (2.8) at  $p = 0$ .

*Proof.* Let us first justify (i). We have seen in the proof of Theorem 3.5(i) that our current assumptions imply that  $(0, \bar{x}) \in \text{int}(\text{gph } \mathcal{F})$ , which in turn yields by the definition of the exact Lipschitzian bound that  $\text{lip } \mathcal{F}(0, \bar{x}) = 0$  in this case.

Next we prove the more difficult assertion (ii) of the theorem, assuming that  $\bar{x}$  is not a strong Slater point for  $\mathcal{F}$  at  $p = 0$ . Observe that by Lemma 3.4 the set (3.8) under the maximum operation on the right-hand side in (4.17) is nonempty and  $w^*$ -compact in  $X^*$  and the maximum over this set is realized and hence it is finite. The inequality “ $\geq$ ” in (4.17) follows from the estimate

$$\text{lip } \mathcal{F}(0, \bar{x}) \geq \|D^* \mathcal{F}(0, \bar{x})\|$$

established for general mappings between Banach spaces in [31, Theorem 1.44] and from formula (3.9) for computing the coderivative norm of the inequality system  $\mathcal{F}$  in (1.1) derived above in Theorem 3.5. It remains to prove the opposite inequality “ $\leq$ ” in (4.17).

To proceed, let  $M := \sup_{t \in T} \|a_t^*\| < \infty$ , and observe that the inequality “ $\leq$ ” in (4.17) is obvious when  $L := \text{lip } \mathcal{F}(0, \bar{x}) = 0$ . Suppose now that  $L > 0$ , and consider any pair  $(p, x)$  sufficiently close to  $(0, \bar{x})$  in the limiting representation (2.2) of the exact Lipschitzian bound  $\text{lip } \mathcal{F}(0, \bar{x})$ . We can confine ourselves to the case of  $(p, x) \notin \text{gph } \mathcal{F}$  by  $L > 0$ . Furthermore, it follows from the structure of  $\mathcal{F}$  in (1.1) that

$$(4.18) \quad 0 < \text{dist}(p; \mathcal{F}^{-1}(x)) = \sup_{t \in T} [\langle a_t^*, x \rangle - b_t - p_t]_+.$$

Moreover, we have the relationships

$$\begin{aligned} \langle a_t^*, x \rangle - b_t - p_t &= \langle a_t^*, x - \bar{x} \rangle + \langle a_t^*, \bar{x} \rangle - b_t - p_t \\ &\leq M \|x - \bar{x}\| + \|p\| \quad \text{for all } t \in T, \end{aligned}$$

which allow us to conclude that

$$(4.19) \quad \begin{aligned} 0 &< \sup_{(x^*, \beta) \in \text{cl}^* C(p)} [\langle x^*, x \rangle - \beta]_+ = \sup_{(x^*, \beta) \in \text{cl}^* C(p)} \{ \langle x^*, x \rangle - \beta \} \\ &\leq M \|x - \bar{x}\| + \|p\|. \end{aligned}$$

Consider further the set

$$C_+(p, x) := \{ (x^*, \beta) \in \text{cl}^* C(p) \mid \langle x^*, x \rangle - \beta > 0 \},$$

which is obviously nonempty, and denote

$$M_{(p,x)} := \sup \{ \|x^*\|^{-1} \mid (x^*, \beta) \in C_+(p, x) \}.$$

In our setting we have  $0 \in \text{int}(\text{dom } \mathcal{F})$  (cf. Lemma 2.3 and Remark 2.4), and therefore  $p \in \text{dom } \mathcal{F}$  for all  $p \in l_\infty(T)$  sufficiently close to the origin. In this case the set  $C_+(p, x)$  cannot contain any element of the form  $(0, \beta)$ , since the contrary would yield by the definition of  $C_+(p, x)$  that  $\beta < 0$ , while Lemma 2.1 gives  $\beta \geq 0$ . Thus we get  $0 < \|x^*\| \leq M$  whenever  $(x^*, \beta) \in C_+(p, x)$  and, in particular,  $M_{(p,x)} \in (0, \infty]$ . It follows, furthermore, that

$$\frac{\sup_{(x^*, \beta) \in \text{cl}^* C(p)} \frac{[\langle x^*, x \rangle - \beta]_+}{\|x^*\|}}{\sup_{(x^*, \beta) \in \text{cl}^* C(p)} [\langle x^*, x \rangle - \beta]_+} = \frac{\sup_{(x^*, \beta) \in \text{cl}^* C(p)} \frac{\langle x^*, x \rangle - \beta}{\|x^*\|}}{\sup_{(x^*, \beta) \in \text{cl}^* C(p)} \{\langle x^*, x \rangle - \beta\}} \leq M_{(p,x)},$$

where the latter inequality implies the estimate

$$L \leq \limsup_{(p,x) \rightarrow (0,\bar{x}), x \notin \mathcal{F}(p) \neq \emptyset} M_{(p,x)} := K.$$

Considering next a sequence  $(p_k, x_k) \rightarrow (0, \bar{x})$  with  $x_k \notin \mathcal{F}(p_k) \neq \emptyset$  and

$$L \leq \lim_{k \rightarrow \infty} M_{(p_k, x_k)} = K,$$

we select a sequence  $\{\alpha_k\}_{k=1}^\infty \subset \mathbb{R}$  such that

$$\lim_{k \rightarrow \infty} \alpha_k = K \quad \text{and} \quad 0 < \alpha_k < M_{(p_k, x_k)} \quad \text{as } k \in \mathbb{N}.$$

Take now  $(x_k^*, \beta_k) \in C_+(p_k, x_k)$  with  $\alpha_k < \|x_k^*\|^{-1}$  for all  $k \in \mathbb{N}$ . Since the sequence  $\{x_k^*\}_{k \in \mathbb{N}} \subset X^*$  is bounded, it contains a subnet  $\{x_\nu^*\}_{\nu \in \mathcal{N}}$  that  $w^*$ -converges to some  $x^* \in X^*$ . Denoting by  $\{p_\nu\}$ ,  $\{x_\nu\}$ ,  $\{\beta_\nu\}$ , and  $\{\alpha_\nu\}$  the corresponding subnets of  $\{p_k\}$ ,  $\{x_k\}$ ,  $\{\beta_k\}$ , and  $\{\alpha_k\}$ , we get from (4.19) that

$$0 < \langle x_\nu^*, x_\nu \rangle - \beta_\nu \leq M \|x_\nu - \bar{x}\| + \|p_\nu\|.$$

Thus  $\langle x_\nu^*, x_\nu \rangle - \beta_\nu \rightarrow 0$ , which implies by the construction above that  $\beta_\nu \rightarrow \langle x^*, \bar{x} \rangle$ . By Lemma 4.5 we get that

$$(x^*, \langle x^*, \bar{x} \rangle) \in \text{cl}^* C(0),$$

and then Lemma 2.3 ensures that  $x^* \neq 0$ .

To conclude proving the inequality “ $\leq$ ” in (4.17), observe that

$$\|x^*\| \leq \liminf_{\nu \in \mathcal{N}} \|x_\nu^*\| \leq \lim_{\nu} \frac{1}{\alpha_\nu} = \frac{1}{K}$$

due to  $\|x_\nu^*\| \leq \alpha_\nu^{-1}$  and  $\lim_{\nu} \alpha_\nu = K$ , which gives

$$L \leq K \leq \frac{1}{\|x^*\|} \leq \max \{ \|z^*\|^{-1} \mid (z^*, \langle z^*, \bar{x} \rangle) \in \text{cl}^* C(0) \}.$$

Remembering the notation above, we complete the proof of the theorem.  $\square$

Comparing, finally, the results on computing the coderivative norm in Theorem 3.5 and the exact Lipschitzian bound in Theorem 4.6 allows us to arrive at the *unconditional* relationship between the coderivative norm and exact Lipschitzian bound of the infinite inequality system  $\mathcal{F}$  with arbitrary Banach decision spaces. This

was known before only for set-valued mappings between finite-dimensional spaces; cf. formula (2.5) and the corresponding discussions in section 2.

**COROLLARY 4.7** (relationship between the exact Lipschitzian bound and coderivative norm). *Let  $\bar{x} \in \mathcal{F}(0)$  for the infinite system (1.1) satisfying the strong Slater condition at  $p = 0$ . Assume that the decision space  $X$  is arbitrary Banach and that the coefficient set  $\{a_t^* \mid t \in T\}$  is bounded in  $X^*$ . Then*

$$(4.20) \quad \text{lip } \mathcal{F}(0, \bar{x}) = \|D^* \mathcal{F}(0, \bar{x})\|.$$

*Proof.* If  $\bar{x}$  is a strong Slater point for  $\mathcal{F}$  at  $p = 0$ , then we get equality (4.20) directly by comparing assertions (i) in Theorems 3.5 and 4.6 which ensure that

$$\text{lip } \mathcal{F}(0, \bar{x}) = \|D^* \mathcal{F}(0, \bar{x})\| = 0.$$

On the other hand, if  $\bar{x}$  is not a strong Slater point for  $\mathcal{F}$  at  $p = 0$ , then (4.20) follows from comparing assertions (ii) in Theorems 3.5 and 4.6 which justify the *same formula* for computing  $\|D^* \mathcal{F}(0, \bar{x})\|$  and  $\text{lip } \mathcal{F}(0, \bar{x})$  in (3.9) and (4.17), respectively.  $\square$

**Acknowledgments.** The main results of this paper were obtained in our research report [10], which was widely disseminated in the optimization community. We would like to thank many colleagues and friends for their helpful comments and remarks. Especially important contributions came from Radu Boţ, who suggested a new proof of Lemma 4.3 via Proposition 4.2, allowing us to remove a certain boundedness assumption imposed in [10], and from Constantin Zălinescu, whose various deep insights led us, in particular, to simplified and more elegant proofs of Theorems 3.5 and 4.6 in comparison with the original proofs given in [10]. We are also gratefully indebted to two anonymous referees and the Associate Editor Jiří Outrata for their valuable suggestions and remarks that allowed us to essentially improve the original presentation.

#### REFERENCES

- [1] E. J. ANDERSON AND P. NASH, *Linear Programming in Infinite-Dimensional Spaces*, Wiley, Chichester, UK, 1987.
- [2] G. BEER, *Topologies on Closed and Closed Convex Sets*, Kluwer, Dordrecht, The Netherlands, 1993.
- [3] J. M. BORWEIN AND Q. J. ZHU, *Techniques of Variational Analysis*, Springer, New York, 2005.
- [4] R. I. BOŢ, S. M. GRAD, AND G. WANKA, *Generalized Moreau-Rockafellar results for composed convex functions*, *Optimization*, 58 (2009), pp. 917–933.
- [5] B. BROSOWSKI, *Parametric Semi-Infinite Optimization*, Peter Lang, Frankfurt, 1982.
- [6] M. J. CÁNOVAS, A. L. DONTCHEV, M. A. LÓPEZ, AND J. PARRA, *Metric regularity of semi-infinite constraint systems*, *Math. Program.*, 104 (2005), pp. 329–346.
- [7] M. J. CÁNOVAS, F. J. GÓMEZ-SENENT, AND J. PARRA, *Regularity modulus of arbitrarily perturbed linear inequality systems*, *J. Math. Anal. Appl.*, 343 (2008), pp. 315–327.
- [8] M. J. CÁNOVAS, A. HANTOUTE, M. A. LÓPEZ, AND J. PARRA, *Lipschitz modulus in convex semi-infinite optimization via DC functions*, *ESAIM Control Optim. Calc. Var.*, 15 (2009), pp. 763–781.
- [9] M. J. CÁNOVAS, D. KLATTE, M. A. LÓPEZ, AND J. PARRA, *Metric regularity in convex semi-infinite optimization under canonical perturbations*, *SIAM J. Optim.*, 18 (2007), pp. 717–732.
- [10] M. J. CÁNOVAS, M. A. LÓPEZ, B. S. MORDUKHOVICH, AND J. PARRA, *Robust Stability and Optimality Conditions for Parametric Infinite and Semi-Infinite Programs*, Research report 12, Department of Mathematics, Wayne State University, Detroit, MI, 2008.
- [11] M. J. CÁNOVAS, M. A. LÓPEZ, B. S. MORDUKHOVICH, AND J. PARRA, *Variational analysis in semi-infinite and infinite programming, II: Necessary optimality conditions*, *SIAM J. Optim.*, submitted.

- [12] M. J. CÁNOVAS, M. A. LOPEZ, J. PARRA, AND M. I. TODOROV, *Stability and well-posedness in linear semi-infinite programming*, SIAM J. Optim., 10 (1999), pp. 82–98.
- [13] N. DINH, M. A. GOBERNA, AND M. A. LÓPEZ, *From linear to convex systems: Consistency, Farkas' lemma and applications*, J. Convex Anal., 13 (2006), pp. 279–290.
- [14] N. DINH, M. A. GOBERNA, AND M. A. LÓPEZ, *On the stability of the feasible set in optimization problems*, submitted (2008).
- [15] N. DINH, B. S. MORDUKHOVICH, AND T. T. A. NGHIA, *Subdifferentials of value functions and optimality conditions for DC and bilevel infinite and semi-infinite programs*, Math. Program., to appear.
- [16] N. DUNFORD AND J. T. SCHWARTZ, *Linear Operators. Part I: General Theory*, Wiley, New York, 1988.
- [17] M. FABIAN, P. HABALA, P. HÁJEK, V. MONTESINOS-SANTALUCÍA, J. PELANT, AND V. ZIZLER, *Functional Analysis and Infinite-Dimensional Geometry*, Springer, New York, 2001.
- [18] M. A. GOBERNA, V. JEYAKUMAR, AND M. A. LÓPEZ, *Necessary and sufficient constraint qualifications for solvability of systems of infinite convex inequalities*, Nonlinear Anal., 68 (2008), pp. 1184–1194.
- [19] M. A. GOBERNA AND M. A. LÓPEZ, *Linear Semi-Infinite Optimization*, Wiley, Chichester, UK, 1998.
- [20] A. D. IOFFE, *Metric regularity and subdifferential calculus*, Russian Math. Surveys, 55 (2000), pp. 501–558.
- [21] H. T. JONGEN, J.-J. RÜCKMANN, AND G.-W. WEBER, *One-parametric semi-infinite optimization: On the stability of the feasible set*, SIAM J. Optim., 4 (1994), pp. 637–648.
- [22] D. KLATTE AND R. HENRION, *Regularity and stability in nonlinear semi-infinite optimization*, in Semi-Infinite Programming, R. Reemtsen and J.-J. Rückmann, eds., Kluwer, Dordrecht, The Netherlands, 1998, pp. 69–102.
- [23] D. KLATTE AND B. KUMMER, *Nonsmooth Equations in Optimization. Regularity, Calculus, Methods and Applications*, Kluwer, Dordrecht, The Netherlands, 2002.
- [24] C. LI AND K. F. NG, *On constraint qualification for an infinite system of convex inequalities in a Banach space*, SIAM J. Optim., 15 (2005), pp. 488–512.
- [25] C. LI, K. F. NG, AND T. K. PONG, *Constraint qualifications for convex inequality systems with applications in constrained optimization*, SIAM J. Optim., 19 (2008), pp. 163–187.
- [26] B. S. MORDUKHOVICH, *Maximum principle in problems of time optimal control with nonsmooth constraints*, J. Appl. Math. Mech., 40 (1976), pp. 960–969.
- [27] B. S. MORDUKHOVICH, *Metric approximations and necessary optimality conditions for general classes of extremal problems*, Soviet Math. Dokl., 22 (1980), pp. 526–530.
- [28] B. S. MORDUKHOVICH, *Complete characterizations of openness, metric regularity, and Lipschitzian properties of multifunctions*, Trans. Amer. Math. Soc., 340 (1993), pp. 1–35.
- [29] B. S. MORDUKHOVICH, *Coderivatives of set-valued mappings: Calculus and applications*, Nonlinear Anal., 30 (1997), pp. 3059–3070.
- [30] B. S. MORDUKHOVICH, *Necessary conditions in nonsmooth minimization via lower and upper subgradients*, Set-Valued Anal., 12 (2004), pp. 163–193.
- [31] B. S. MORDUKHOVICH, *Variational Analysis and Generalized Differentiation, I: Basic Theory*, Springer, Berlin, 2006.
- [32] B. S. MORDUKHOVICH, *Variational Analysis and Generalized Differentiation, II: Applications*, Springer, Berlin, 2006.
- [33] J. V. OUTRATA, *Mathematical programs with equilibrium constraints: Theory and numerical methods*, in Nonsmooth Mechanics of Solids, J. Haslinger and G. E. Stavroulakis, eds., Springer, New York, 2006, pp. 221–274.
- [34] R. R. PHELPS, *Convex Functions, Monotone Operators, and Differentiability*, 2nd ed., Springer, Berlin, 1993.
- [35] R. T. ROCKAFELLAR AND R. J.-B. WETS, *Variational Analysis*, Springer, Berlin, 1998.
- [36] W. SCHIROTZKE, *Nonsmooth Analysis*, Springer, Berlin, 2007.
- [37] O. STEIN, *Bilevel Strategies in Semi-Infinite Programming*, Kluwer, Boston, 2003.
- [38] C. ZĂLINESCU, *Solvability results for sublinear functions and operators*, Z. Oper. Res. Ser. A-B, 31 (1987), pp. 79–101.
- [39] C. ZĂLINESCU, *Convex Analysis in General Vector Spaces*, World Scientific, Singapore, 2002.

## VARIATIONAL ANALYSIS IN SEMI-INFINITE AND INFINITE PROGRAMMING, I: STABILITY OF LINEAR INEQUALITY SYSTEMS OF FEASIBLE SOLUTIONS\*

M. J. CÁNOVAS<sup>†</sup>, M. A. LÓPEZ<sup>‡</sup>, B. S. MORDUKHOVICH<sup>§</sup>, AND J. PARRA<sup>†</sup>

**Abstract.** This paper concerns applications of advanced techniques of variational analysis and generalized differentiation to parametric problems of semi-infinite and infinite programming, where decision variables run over finite-dimensional and infinite-dimensional spaces, respectively. Part I is primarily devoted to the study of robust Lipschitzian stability of feasible solutions maps for such problems described by parameterized systems of infinitely many linear inequalities in Banach spaces of decision variables indexed by an arbitrary set  $T$ . The parameter space of admissible perturbations under consideration is formed by all bounded functions on  $T$  equipped with the standard supremum norm. Unless the index set  $T$  is finite, this space is intrinsically infinite-dimensional (nonreflexive and nonseparable) of the  $l_\infty$  type. By using advanced tools of variational analysis and exploiting specific features of linear infinite systems, we establish complete characterizations of robust Lipschitzian stability entirely via their initial data with computing the exact bound of Lipschitzian moduli. A crucial part of our analysis addresses the precise computation of the coderivative of the feasible set mapping and its norm. The results obtained are new in both semi-infinite and infinite frameworks.

**Key words.** semi-infinite and infinite programming, variational analysis, linear infinite inequality systems, robust stability, generalized differentiation, coderivatives

**AMS subject classifications.** 90C34, 90C05, 49J52, 49J53, 65F22

**DOI.** 10.1137/090765948

**1. Introduction.** This paper mainly deals with parameterized *infinite* systems of linear inequalities

$$(1.1) \quad \mathcal{F}(p) := \{x \in X \mid \langle a_t^*, x \rangle \leq b_t + p_t, t \in T\}, \quad p = (p_t)_{t \in T},$$

with an *arbitrary index set*  $T$ , where  $x \in X$  is a *decision* variable belonging to an arbitrary Banach space  $X$  (which may be finite-dimensional), and where  $p = (p_t)_{t \in T} \in P$  is a functional *parameter* taking values in the prescribed Banach space  $P$  of perturbations specified below. Infinite inequality systems of this type are important in various areas of mathematics and applications, while our primary interest in them is driven by applications to problems of *semi-infinite* and *infinite programming* corresponding to finite-dimensional and infinite-dimensional decision spaces  $X$ , respectively; see, e.g., the books [1, 5, 19, 37] and the references therein. Some applications of the results obtained in this paper to necessary optimality conditions for semi-infinite and infinite programs can be found in [11].

The data of (1.1) are given as follows:

---

\*Received by the editors July 22, 2009; accepted for publication (in revised form) September 23, 2009; published electronically December 16, 2009. This research was partially supported by grants MTM2005-08572-C03 (01-02) from MEC (Spain) and FEDER (EU), MTM2008-06695-C03 (01-02) from MICINN (Spain), and ACOMP/2009/047&133 from Generalitat Valenciana (Spain).  
<http://www.siam.org/journals/siopt/20-3/76594.html>

<sup>†</sup>Center of Operations Research, Miguel Hernández University of Elche, 03202 Elche (Alicante), Spain (canovas@umh.es, parra@umh.es).

<sup>‡</sup>Department of Statistics and Operations Research, University of Alicante, 03080 Alicante, Spain (marco.antonio@ua.es).

<sup>§</sup>Department of Mathematics, Wayne State University, Detroit, MI 48202 (boris@math.wayne.edu). The research of this author was partially supported by the National Science Foundation (USA) under grant DMS-0603846.



- $a_t^* \in X^*$  for all  $t \in T$ , where the space  $X^*$  is topologically dual to  $X$  with the canonical pairing  $\langle \cdot, \cdot \rangle$  between  $X$  and  $X^*$ . If no confusion arises, we use the *same notation*  $\| \cdot \|$  for the given norm in  $X$  and the corresponding dual norm in  $X^*$  defined by

$$\|x^*\| := \sup \{ \langle x^*, x \rangle \mid \|x\| \leq 1 \}, \quad x^* \in X^*.$$

We always assume that  $a_t^*$  are *fixed* and *arbitrary* in  $X^*$  for all  $t \in T$ .

- $b_t \in \mathbb{R}$  for all  $t \in T$ . We identify the collection  $\{b_t \mid t \in T\}$  with the real-valued function  $b: T \rightarrow \mathbb{R}$  and assume that it is *fixed* and *arbitrary*.

- $p_t = p(t) \in \mathbb{R}$  for all  $t \in T$ . These functional parameters  $p: T \rightarrow \mathbb{R}$  are our varying *perturbations*, which are taken from the Banach *parameter space*  $P := l_\infty(T)$  of all bounded functions on  $T$  with the supremum norm

$$(1.2) \quad \|p\|_\infty := \sup_{t \in T} |p_t| = \sup \{ |p(t)| \mid t \in T \}$$

(see, e.g., [17]), where the subscript “ $\infty$ ” is omitted if no confusion arises. When the index set  $T$  is compact (which is not assumed in this paper) and the perturbations  $p(\cdot)$  are restricted to be continuous on  $T$ , the maximum is realized in (1.2), and thus the parameter space  $l_\infty(T)$  reduces to the classical space  $\mathcal{C}(T)$  of continuous functions over a compact set.

The primary goal of this paper is to obtain comprehensive characterizations of *robust Lipschitzian stability* of infinite inequality systems (1.1) expressed entirely in terms of their initial data. By robust Lipschitzian stability we understand the fulfillment of the so-called *Lipschitz-like* (known also as Aubin) property of the mapping  $\mathcal{F}(p)$  in (1.1) *around* the reference point. This property is stable with respect to small perturbations of parameters and is crucial for both qualitative and quantitative/numerical aspects of optimization theory and applications; see, e.g., [31, 32, 35] and section 2 for more details and references.

To establish constructive characterizations of robust Lipschitzian stability in this paper and to derive efficient optimality conditions for semi-infinite and infinite programs in [11], we develop an advanced approach of *variational analysis* based on *generalized differentiation*. To the best of our knowledge, this approach is new in the literature on semi-infinite and infinite programming despite many publications related to various stability properties and applications of linear infinite inequality systems, most of which concern the case of finite-dimensional spaces  $X$  of decision variables (i.e., in the semi-infinite programming framework); see, e.g., [1, 19] for comprehensive overviews on this field and also [5], which is confined to the parameter space of continuous perturbations  $P = \mathcal{C}(T)$  when the index set  $T$  is a compact Hausdorff space. We refer the reader to [12] for the study of *qualitative stability* (formalized through certain semicontinuity properties of feasible solution and optimal solution mappings) in the framework of  $X = \mathbb{R}^n$ , an arbitrary index set  $T$ , and arbitrary perturbations. In the same semi-infinite context, for a *quantitative perspective* (through Lipschitzian properties), the reader is directed to [7], and to [6] for the case of continuous perturbations. Let us mention the recent paper [14] addressing the case of infinite linear programming from the viewpoint of qualitative stability. We also refer the reader to, e.g., [8, 9, 13, 15, 18, 24, 25] for the study of convex semi-infinite and infinite programs and to [21, 22, 37] for their smooth nonlinear counterparts.

The approach of this paper is mainly based on *coderivative analysis* of the parametric linear infinite inequality systems  $\mathcal{F}$  in (1.1), which eventually leads us to *com-*

plete characterizations of robust Lipschitzian stability for the parametric sets of feasible solutions in infinite/semi-infinite programming expressed entirely via their initial data with precise computing the *exact bound* of Lipschitzian moduli.

Coderivatives of set-valued mappings introduced in [27] have been well recognized as a powerful tool of variational analysis and its numerous applications, particularly to problems of optimization and control; see, e.g., [3, 23, 30, 31, 32, 33, 35, 36] and the references therein. However, we are not familiar with any implementation of coderivatives in problems of infinite or semi-infinite programming as well as with their application to analyzing stability of linear infinite inequality systems of type (1.1) in finite or infinite dimensions.

The power of coderivatives in variational analysis and its applications comes, first, from the possibility of obtaining in their terms verifiable *pointwise characterizations* of *robust Lipschitzian* properties of set-valued mappings (as well as of the equivalent properties of *metric regularity* and *linear openness* for the inverse mappings) and of deriving *necessary optimality conditions* in rather general settings. These developments are strongly supported by comprehensive pointwise *coderivative calculus* based on *variational/extremal principles* of advanced variational analysis; see [31, 32] and the references therein. However, a number of results in this vein are limited in infinite dimensions. In particular, the available coderivative characterizations of the Lipschitz-like property of closed-graph mappings  $F: Z \rightrightarrows Y$  obtained in [31, Theorem 4.10] require that both spaces  $Z$  and  $Y$  are *Asplund* (i.e., every separable subspace of them has a separable dual) while the precise coderivative formula for computing the *exact Lipschitzian bound* is established therein via the coderivative norm under the *finite-dimensionality* assumption on  $Z$ . But this is *never the case* for our infinite inequality system  $\mathcal{F}: P \rightrightarrows X$  from (1.1), where the parameter space ( $Z =$ )  $P = l_\infty(T)$  is always *infinite-dimensional* and *not Asplund* unless the index set  $T$  is finite!

This paper contains new and fairly comprehensive results in the aforementioned directions for the infinite/semi-infinite systems under consideration, which essentially take into account underlying specific features of the infinite inequality constraints (1.1) largely related to the possibility of employing an appropriate extended version of the fundamental *Farkas lemma* for infinite systems of linear inequalities in *general Banach spaces*.

The rest of paper is organized as follows. Section 2 presents some preliminary material from convex and variational analysis widely used in formulations and proofs of the subsequent main results. In section 3 we provide *precise calculations* of the basic *coderivative*  $D^*\mathcal{F}$  and its *norm* at the reference/nominal point for the feasible solution map  $\mathcal{F}: l_\infty(T) \rightrightarrows X$  in (1.1) via the initial data of  $\mathcal{F}$  in the general case of an arbitrary index set  $T$  and an arbitrary Banach space  $X$  of decision variables.

Section 4 is devoted to deriving *coderivative characterizations* of *robust stability* for the feasible solution system (1.1) of infinite inequalities with an arbitrary index set  $T$ , which are explicitly expressed in terms of the initial data  $\{a_t^*, b_t, t \in T\}$ . We establish verifiable *criteria* (i.e., necessary and sufficient conditions) for the fulfillment of the Lipschitz-like (and hence the classical local Lipschitzian) property of  $\mathcal{F}$  around the reference points and derive furthermore the *precise formulas* for computing the *exact bounds* of Lipschitzian moduli in the case of general Banach spaces  $X$ . It is worth mentioning that the criteria and exact bound formulas obtained in this section are represented in the conventional *coderivative form* of variational analysis as in [31, Theorem 4.10] for the case of abstract set-valued mappings, but with *no* Asplund space and finite-dimensionality requirements imposed therein. In fact, the latter requirements are *never* satisfied for the infinite linear inequality systems (1.1)

under consideration in either the infinite programming or semi-infinite programming framework.

Our notation is basically standard and conventional in the areas of variational analysis and infinite/semi-infinite programming; see, e.g., [19, 31, 35]. Unless otherwise stated, all the spaces under consideration are *Banach* with the corresponding norm  $\|\cdot\|$ . Recall that  $w^*$  indicates the weak\* topology of a dual space, and we use the symbol  $w^*$ -lim for the weak\* *topological* limit, which generally means the weak\* convergence of *nets* denoted usually by  $\{x_\nu^*\}_{\nu \in \mathcal{N}}$ . In the case of sequences we use the standard notation  $\mathbb{N} := \{1, 2, \dots\}$  for the collection of all natural numbers.

Given a subset  $\Omega \subset Z$  of a Banach space, the symbols  $\text{int } \Omega$ ,  $\text{cl } \Omega$ ,  $\text{co } \Omega$ , and  $\text{cone } \Omega$  stand, respectively, for the interior, closure, convex hull, and conic convex hull of  $\Omega$ ; the notation  $\text{cl}^* \Theta$  signifies the weak\* closure of a subset  $\Theta \subset Z^*$  in the dual space. Given a set-valued mapping  $F: Z \rightrightarrows Y$ , we denote its *domain*, *graph*, and *inverse* by, respectively,

$$\text{dom } F = \{z \in Z \mid F(z) \neq \emptyset\}, \quad \text{gph } F := \{(z, y) \in Z \times Y \mid y \in F(z)\},$$

and  $F^{-1}(y) := \{z \in Z \mid (z, y) \in \text{gph } F\}$ . Considering finally an arbitrary index set  $T$ , let  $\mathbb{R}^T$  be the *product space* of  $\lambda = (\lambda_t \mid t \in T)$  with  $\lambda_t \in \mathbb{R}$  for all  $t \in T$ , let  $\mathbb{R}^{(T)}$  be the collection of  $\lambda \in \mathbb{R}^T$  such that  $\lambda_t \neq 0$  for *finitely many*  $t \in T$ , and let  $\mathbb{R}_+^{(T)}$  be the *positive cone* in  $\mathbb{R}^{(T)}$  defined by

$$\mathbb{R}_+^{(T)} := \{\lambda \in \mathbb{R}^{(T)} \mid \lambda_t \geq 0 \text{ for all } t \in T\}.$$

**2. Basic definitions and preliminaries.** In this section we discuss the basic notions and tools needed for our subsequent analysis and results. As mentioned in section 1, a major focus of this paper is *robust Lipschitzian stability* of the feasible solution map given by (1.1). By such a robust stability we understand Lipschitzian behavior *around* (i.e., in a neighborhood) of the reference point. The most natural formalization of this behavior widely recognized in variational (as well as in general nonlinear) analysis is known as the Lipschitz-like or Aubin property, which can be viewed as a graphical localization (in the set-valued case) of the classical local Lipschitzian property of single-valued and set-valued mappings.

Given a set-valued mapping  $F: Z \rightrightarrows Y$  between Banach spaces, we say that  $F$  is *Lipschitz-like* around  $(\bar{z}, \bar{y}) \in \text{gph } F$  with modulus  $\ell \geq 0$  if there are neighborhoods  $U$  of  $\bar{z}$  and  $V$  of  $\bar{y}$  such that

$$(2.1) \quad F(z) \cap V \subset F(u) + \ell \|z - u\| \mathbb{B} \text{ for any } z, u \in U,$$

where  $\mathbb{B}$  stands for the closed unit ball in the space in question. The infimum of moduli  $\{\ell\}$  over all the combinations of  $\{\ell, U, V\}$  satisfying (2.1) is called the *exact Lipschitzian bound* of  $F$  around  $(\bar{z}, \bar{y})$  and is labeled as  $\text{lip } F(\bar{z}, \bar{y})$ . If  $V = Y$  in (2.1), this relationship signifies the classical (Hausdorff) *local Lipschitzian* property of  $F$  around  $\bar{z}$  with the exact Lipschitzian bound denoted by  $\text{lip } F(\bar{z})$  in this case.

It is worth mentioning that the Lipschitz-like property of an arbitrary mapping  $F: Z \rightrightarrows Y$  between Banach spaces is equivalent to two other fundamental properties in nonlinear analysis but for the *inverse* mapping  $F^{-1}: Y \rightrightarrows Z$ , namely, to the *metric regularity* of  $F^{-1}$  and to the *linear openness* of  $F^{-1}$  around  $(\bar{y}, \bar{z})$ , with the corresponding relationships between their exact bounds (see, e.g., [20, 31, 35]).

It is well known that the exact Lipschitzian bound of  $F$  around  $(\bar{z}, \bar{y})$  admits the following limiting representation via the *distance function* to a set:

$$(2.2) \quad \text{lip } F(\bar{z}, \bar{y}) = \limsup_{(z, y) \rightarrow (\bar{z}, \bar{y})} \frac{\text{dist}(y; F(z))}{\text{dist}(z; F^{-1}(y))},$$

where  $\inf \emptyset = \infty$  (and hence  $\text{dist}(x; \emptyset) = \infty$ ) as usual and where  $0/0 := 0$ . We have accordingly that  $\text{lip } F(\bar{z}, \bar{y}) = \infty$  if  $F$  is not Lipschitz-like around  $(\bar{z}, \bar{y})$ .

A remarkable fact consists of the possibility of *characterizing* pointwisely the (derivative-free) Lipschitz-like property of  $F$  around  $(\bar{z}, \bar{y})$ —and hence its local Lipschitzian, metric regularity, and linear openness counterparts—in terms of a *dual-space* construction of generalized differentiation called the *coderivative* of  $F$  at  $(\bar{z}, \bar{y}) \in \text{gph } F$ . The latter is a positively homogeneous multifunction  $D^*F(\bar{z}, \bar{y}): Y^* \rightrightarrows Z^*$  defined by

$$(2.3) \quad D^*F(\bar{z}, \bar{y})(y^*) := \{z^* \in Z^* \mid (z^*, -y^*) \in N((\bar{z}, \bar{y}); \text{gph } F)\}, \quad y^* \in Y^*,$$

where  $N(\cdot; \Omega)$  stands for the collection of generalized normals to a set at a given point known as the basic, or limiting, or Mordukhovich, normal cone; see, e.g., [26, 31, 35, 36] and the references therein. When both  $Z$  and  $Y$  are *finite-dimensional*, it is proved in [28] (cf. also [35, Theorem 9.40]) that a closed-graph mapping  $F: Z \rightrightarrows Y$  is Lipschitz-like around  $(\bar{z}, \bar{y}) \in \text{gph } F$  *if and only if*

$$(2.4) \quad D^*F(\bar{z}, \bar{y})(0) = \{0\},$$

and that the exact Lipschitzian bound of moduli  $\{\ell\}$  in (2.1) is computed by

$$(2.5) \quad \text{lip } F(\bar{z}, \bar{y}) = \|D^*F(\bar{z}, \bar{y})\| := \sup \{\|z^*\| \mid z^* \in D^*F(\bar{z}, \bar{y})(y^*), \|y^*\| \leq 1\}.$$

The situation is significantly more involved in infinite dimensions. It is proved in [29] (see also [31, Theorem 4.10]) that a closed-graph mapping  $F: Z \rightrightarrows Y$  is Lipschitz-like around  $(\bar{z}, \bar{y}) \in \text{gph } F$  *if and only if* the coderivative condition (2.4) holds in terms of the so-called *mixed coderivative* (which reduces to (2.3) in finite dimensions and the setting considered in this paper) together with a certain “partial sequential normal compactness” condition (which automatically holds in finite dimensions and in the setting of this paper) provided that both spaces  $Z$  and  $Y$  are *Asplund*. The latter property is defined in section 1; we also refer the reader to [17, 31, 34] for more details and various characterizations of this remarkable and well-investigated subclass of Banach spaces that includes, in particular, all reflexive spaces, while it does not include, e.g., the classical spaces  $\mathcal{C}$ ,  $l_1$ ,  $L_1$ ,  $l_\infty$ , and  $L_\infty$ .

The situation is even more complicated with infinite-dimensional extensions of the exact bound formula in (2.5). The aforementioned results of [29, 31] give merely upper and lower estimates for  $\text{lip } F(\bar{z}, \bar{y})$ , which ensure the precise equality in (2.5) in our setting here provided that  $Y$  is *Asplund* while  $Z$  is *finite-dimensional*.

The set-valued mapping  $F: Z \rightrightarrows Y$  considered in this paper is  $\mathcal{F}: l_\infty(T) \rightrightarrows X$  defined by the infinite system of linear inequalities (1.1); in what follows we always assume that the index set  $T$  is *infinite*, which is a characteristic feature of infinite and semi-infinite programs. In this setting the domain/parameter space  $Z = l_\infty(T)$  is an *infinite-dimensional* Banach that is *never Asplund*. Also, we do not suppose in this paper that our decision space  $X$  is anything but *arbitrary Banach*.

In this general setting for (1.1) we show that the coderivative condition (2.4) is *necessary and sufficient* for the Lipschitz-like property of  $F = \mathcal{F}$  around the reference/

nominal solution  $(\bar{p}, \bar{x}) \in \text{gph } \mathcal{F}$  and (2.5) is a *precise formula* for computing the exact Lipschitzian bound  $\text{lip } \mathcal{F}(\bar{p}, \bar{x})$ . This is *exactly what we have in finite dimensions*, while it is far removed from being a part of the infinite-dimensional variational theory in [31]. Moreover, we express the relationships in (2.4) and (2.5) *explicitly* in terms of the *initial data* of (1.1).

To proceed further, observe that the graph

$$(2.6) \quad \text{gph } \mathcal{F} = \{(p, x) \in l_\infty(T) \times X \mid \langle a_t^*, x \rangle \leq b_t + p_t \text{ for all } t \in T\}$$

of the mapping  $\mathcal{F}: l_\infty(T) \rightrightarrows X$  in (1.1) is *convex*. Hence the basic normal cone to  $\text{gph } \mathcal{F}$  at  $(\bar{p}, \bar{x}) \in \text{gph } \mathcal{F}$  reduces to

$$N((\bar{p}, \bar{x}); \text{gph } \mathcal{F}) = \{(p^*, x^*) \in l_\infty(T)^* \times X^* \mid \langle (p^*, x^*), (p, x) - (\bar{p}, \bar{x}) \rangle \leq 0 \text{ for } (p, x) \in \text{gph } \mathcal{F}\}$$

and the coderivative (2.3) of  $\mathcal{F}$  admits the representation

$$(2.7) \quad D^* \mathcal{F}(\bar{p}, \bar{x})(x^*) = \{p^* \in l_\infty(T)^* \mid \langle p^*, \bar{p} \rangle - \langle x^*, \bar{x} \rangle = \max_{(p,x) \in \text{gph } \mathcal{F}} [\langle p^*, p \rangle - \langle x^*, x \rangle]\}.$$

Let us now present two preliminary results that play an important role in our subsequent analysis. The first one is taken from [13, Lemma 2.4] and can be viewed as an *extended Farkas lemma* for linear infinite inequality systems in Banach spaces. An alternative proof can be derived from [38, Theorem 4].

LEMMA 2.1 (extended Farkas lemma). *Let  $p \in \text{dom } \mathcal{F}$  for the infinite system (1.1) with a Banach decision space  $X$ , and let  $(x^*, \alpha) \in X^* \times \mathbb{R}$ . The following are equivalent:*

(i) *We have  $\langle x^*, x \rangle \leq \alpha$  whenever  $x \in \mathcal{F}(p)$ , i.e.,*

$$[\langle a_t^*, x \rangle \leq b_t + p_t \text{ for all } t \in T] \implies [\langle x^*, x \rangle \leq \alpha].$$

(ii) *The pair  $(x^*, \alpha)$  satisfies the inclusion*

$$(x^*, \alpha) \in \text{cl}^* \text{cone}[\{(a_t^*, b_t + p_t) \mid t \in T\} \cup \{(0, 1)\}] \text{ with } 0 \in X^*.$$

Throughout the paper we largely use the parametric *characteristic sets*

$$(2.8) \quad C(p) := \text{co}\{(a_t^*, b_t + p_t) \mid t \in T\}, \quad p \in l_\infty(T),$$

and suppose with no loss of generality that our *nominal parameter* is the zero function  $\bar{p} = 0$  in the parameter space  $l_\infty(T)$ .

Let us recall a well-recognized *qualification condition* for linear infinite inequalities, which is often used in problems of semi-infinite and infinite programming.

DEFINITION 2.2 (strong Slater condition). *We say that the infinite system (1.1) satisfies the strong Slater condition (SSC) at  $p = (p_t)_{t \in T}$  if there is  $\hat{x} \in X$  such that*

$$(2.9) \quad \sup_{t \in T} [\langle a_t^*, \hat{x} \rangle - b_t - p_t] < 0.$$

*Furthermore, every point  $\hat{x} \in X$  satisfying condition (2.9) is a strong Slater point for system (1.1) at  $p = (p_t)_{t \in T}$ .*

The next result contains several equivalent descriptions and interpretations of the strong Slater condition used in what follows; the most important is the equivalence

(i) $\iff$ (ii). Note that a similar equivalence can be found in [14] for more general convex systems in locally convex spaces with different spaces of associated parameters. We include here a simplified proof of this equivalence for the reader's convenience.

LEMMA 2.3 (equivalent descriptions of the strong Slater condition). *Let  $X$  be a Banach space, and let  $p \in \text{dom } \mathcal{F}$  for the linear infinite inequality system (1.1). Then the following properties are equivalent:*

- (i)  $\mathcal{F}$  satisfies the strong Slater condition at  $p$ .
- (ii)  $(0, 0) \notin \text{cl}^* C(p)$  via the characteristic set from (2.8).
- (iii)  $p \in \text{int}(\text{dom } \mathcal{F})$ .
- (iv)  $\mathcal{F}$  is Lipschitz-like around  $(p, x)$  for all  $x \in \mathcal{F}(p)$ .

Moreover, if the set  $\{a_t^* \mid t \in T\}$  is bounded in  $X^*$ , the conditions above are equivalent to

- (v) there exists  $\hat{x} \in X$  such that  $(p, \hat{x}) \in \text{int}(\text{gph } \mathcal{F})$ .

*Proof.* We begin with the proof of (i) $\implies$ (ii). Arguing by contradiction, assume that  $(0, 0) \in \text{cl}^* C(p)$ . Then there is a net  $\{\lambda_\nu\}_{\nu \in \mathcal{N}} \in \mathbb{R}_+^{(T)}$  satisfying  $\sum_{t \in T} \lambda_{t\nu} = 1$  for all  $\nu \in \mathcal{N}$  and the limiting condition

$$(2.10) \quad (0, 0) = w^*\text{-}\lim_{\nu} \sum_{t \in T} \lambda_{t\nu} (a_t^*, b_t + p_t).$$

If  $\hat{x}$  is a strong Slater point for system (1.1) at  $p$ , we find  $\vartheta > 0$  such that

$$\langle a_t^*, \hat{x} \rangle - b_t - p_t \leq -\vartheta \quad \text{for all } t \in T.$$

Then (2.10) leads to the following contradiction:

$$0 = \langle 0, \hat{x} \rangle + 0 \cdot (-1) = \lim_{\nu} \sum_{t \in T} \lambda_{t\nu} (\langle a_t^*, \hat{x} \rangle + (b_t + p_t) \cdot (-1)) \leq -\vartheta.$$

Let us next justify the converse implication (ii) $\implies$ (i). By [13, Theorem 3.1] we have

$$p \in \text{dom } \mathcal{F} \iff (0, -1) \notin \text{cl}^* \text{cone}\{(a_t^*, b_t + p_t) \mid t \in T\}.$$

Then the *strong separation theorem* ensures the existence of  $(0, 0) \neq (v, \alpha) \in X \times \mathbb{R}$  with

$$(2.11) \quad \langle a_t^*, v \rangle + \alpha(b_t + p_t) \leq 0 \quad \text{for all } t \in T,$$

$$\langle 0, v \rangle + (-1)\alpha = -\alpha > 0.$$

At the same time by (ii) we have  $(0, 0) \neq (z, \beta) \in X \times \mathbb{R}$  and  $\gamma \in \mathbb{R}$  for which

$$(2.12) \quad \langle a_t^*, z \rangle + \beta(b_t + p_t) \leq \gamma < 0 \quad \text{whenever } t \in T.$$

Consider further the combination

$$(u, \eta) := (z, \beta) + \lambda(v, \alpha),$$

and select  $\lambda > 0$  to be sufficiently large to ensure that  $\eta < 0$ . Defining now  $\hat{x} := -\eta^{-1}u$ , we observe from (2.11) and (2.12) that

$$\langle a_t^*, \hat{x} \rangle - b_t - p_t = -\eta^{-1}(\langle a_t^*, u \rangle + \eta(b_t + p_t)) \leq -\eta^{-1}\gamma < 0.$$

This allows us to conclude that  $\hat{x}$  is a strong Slater point for system (1.1) at  $p$ .

To prove implication (i) $\implies$ (iii), assume that  $\hat{x}$  is a strong Slater point for system (1.1) at  $p$ , and find  $\vartheta > 0$  such that

$$\langle a_t^*, \hat{x} \rangle - b_t - p_t \leq -\vartheta \text{ for all } t \in T.$$

Then it is obvious that for any  $q \in l_\infty(T)$  with  $\|q\| < \vartheta$  we have  $\hat{x} \in \mathcal{F}(p + q)$ . Therefore,  $p + q \in \text{dom } \mathcal{F}$ , and thus (iii) holds.

Let us further proceed with justifying implication (iii) $\implies$ (i). If  $p \in \text{int}(\text{dom } \mathcal{F})$ , then  $p + q \in \text{dom } \mathcal{F}$  provided that  $q_t = -\vartheta$  as  $t \in T$  and that  $\vartheta > 0$  is sufficiently small. Thus every  $\hat{x} \in \mathcal{F}(p + q)$  is a strong Slater point for the infinite system (1.1) at  $p$ .

The equivalence between (iii) and (iv) is a consequence of the classical Robinson–Ursescu closed-graph/metric regularity theorem; see, e.g., [20] and [31, Chapter 4] and references therein for more discussion.

It remains to consider condition (v). We can easily observe that (v) always implies (iii) and hence the other conditions of the lemma. Suppose now that the set  $\{a_t^* \mid t \in T\}$  is bounded in  $X^*$ , and show that in this case (i) implies (v). Select  $M \geq 0$  such that  $\|a_t^*\| \leq M$  for every  $t \in T$ , and take  $\hat{x} \in X$  satisfying (2.9). Denote

$$\gamma := -\sup_{t \in T} [\langle a_t^*, \hat{x} \rangle - b_t - p_t] > 0,$$

and consider any pair  $(p', u) \in l_\infty(T) \times X$  satisfying the relationships

$$\|u\| \leq \eta := \gamma / (M + 1) > 0 \text{ and } \|p'\| \leq \eta.$$

It is easy to see that for such  $(p', u)$  and every  $t \in T$  we have

$$\langle a_t^*, \hat{x} + u \rangle - b_t - p_t - p'_t \leq -\gamma + M \|u\| + \|p'\| \leq \eta(M + 1) - \gamma = 0,$$

and so  $(p + p', \hat{x} + u) \in \text{gph } \mathcal{F}$ . Thus  $(p, \hat{x}) \in \text{int}(\text{gph } \mathcal{F})$ , which gives (i) $\implies$ (v) and completes the proof of the lemma.  $\square$

*Remark 2.4* (relationships between interiority and boundedness conditions). Note that we have the nonempty *interiority condition*  $\text{int}(\text{gph } \mathcal{F}) \neq \emptyset$  provided that the set  $\{a_t^* \mid t \in T\}$  is *bounded* in  $X^*$  and that  $\text{gph } \mathcal{F} \neq \emptyset$ . Also we have  $\text{int}(\text{dom } \mathcal{F}) \neq \emptyset$  if  $\text{gph } \mathcal{F} \neq \emptyset$  with no boundedness assumption.

To justify the first statement, take  $(p, x) \in \text{gph } \mathcal{F}$  and  $\|a_t^*\| \leq M < \infty$  for every  $t \in T$ , and define  $e \in l_\infty(T)$  by  $e_t := 1$  for all  $t \in T$ . Then arguing similarly to the proof of the last implication in Lemma 2.1, we get  $(p + e + p', x + u) \in \text{gph } \mathcal{F}$  whenever  $\|p'\| \leq \eta := 1 / (M + 1)$  and  $\|u\| \leq \eta$ . This gives  $(p + e, x) \in \text{int}(\text{gph } \mathcal{F})$ . If, furthermore,  $(p, x) \in \text{gph } \mathcal{F}$ , we can easily check that  $x \in \mathcal{F}(p + e + p')$  provided that  $\|p'\| \leq 1$ , and therefore  $p + e \in \text{int}(\text{dom } \mathcal{F})$ .

The major space for our consideration in this paper is the parameter space  $l_\infty(T)$  of bounded functions  $p: T \rightarrow \mathbb{R}$  on  $T$  with the supremum norm (1.2). It is obviously a Banach space that is *never finite-dimensional* when the index set  $T$  is infinite, which is our standing assumption. Let us show that it is *never Asplund*.

**PROPOSITION 2.5** (parameter space is never Asplund). *The parameter space  $l_\infty(T)$  is Asplund if and only if the index set  $T$  is finite.*

*Proof.* If  $T$  is countable (i.e.,  $T = \mathbb{N}$  and the parameter space is the classical space of sequences  $l_\infty$ ), the proof can be found in [34, Example 1.21]; in fact, this space is

not even weak Asplund. The same arguments can be adapted for any infinite index set  $T$ .  $\square$

Finally, in this section, we recall a convenient description of the topological dual space  $l_\infty(T)^*$  to the parameter space  $l_\infty(T)$ . According to [16], there is an isometric isomorphism between  $l_\infty(T)^*$  and the space of *bounded and additive measures*

$$ba(T) = \{\mu: 2^T \rightarrow \mathbb{R} \mid \mu \text{ is bounded and additive}\}$$

satisfying the relationship

$$\langle \mu, p \rangle = \int_T p_t \mu(dt) \quad \text{with } p = (p_t)_{t \in T}.$$

The dual norm on  $ba(T)$  corresponding to (1.2) is the *total variation* of  $\mu \in ba(T)$  on the index set  $T$  defined by

$$\|\mu\| := \sup_{A \subset T} \mu(A) - \inf_{B \subset T} \mu(B).$$

In what follows we always identify the measure space  $ba(T)$  with the dual parameter space  $l_\infty(T)^*$  and use, for the notational unification,  $p^* \in l_\infty(T)^*$  instead of  $\mu \in ba(T)$ .

**3. Computing coderivatives and coderivative norms for linear infinite inequality systems.** In this section we establish a constructive representation of the coderivative  $D^*\mathcal{F}(0, \bar{x})$  for the feasible solution map  $\mathcal{F}$  at the nominal point  $(0, \bar{x})$  and compute its norm  $\|D^*\mathcal{F}(0, \bar{x})\|$  in terms of the initial data of the linear infinite inequality system (1.1). Let us first describe the *normal cone* to the convex graph (2.6) employing the *extended Farkas lemma* presented above. In what follows  $\delta_t$  denotes the classical *Dirac measure* at  $t \in T$  satisfying

$$\langle \delta_t, p \rangle = p_t \quad \text{as } t \in T \quad \text{for } p = (p_t)_{t \in T} \in l_\infty(T).$$

**PROPOSITION 3.1** (computing normals to the graphical set of feasible solutions). *Let  $(\bar{p}, \bar{x}) \in \text{gph } \mathcal{F}$  for the graphical set (2.6) with a Banach decision space  $X$ , and let  $(p^*, x^*) \in l_\infty(T)^* \times X^*$ . Then we have  $(p^*, x^*) \in N((\bar{p}, \bar{x}); \text{gph } \mathcal{F})$  if and only if*

$$(3.1) \quad (p^*, x^*, \langle p^*, \bar{p} \rangle + \langle x^*, \bar{x} \rangle) \in \text{cl}^* \text{cone}[\{(-\delta_t, a_t^*, b_t) \mid t \in T\} \cup \{(0, 0, 1)\}],$$

where  $0 \in l_\infty(T)^*$  and  $0 \in X^*$  stand for the first and second entry of the last triple in (3.1), respectively. Furthermore, the inclusion  $(p^*, x^*) \in N((\bar{p}, \bar{x}); \text{gph } \mathcal{F})$  implies that  $p^* \leq 0$  in the space  $ba(T)$ , i.e.,  $p^*(A) \leq 0$  for all  $A \subset T$ .

*Proof.* Observe from (2.6) and from the definition of the Dirac measure that the graph of  $\mathcal{F}$  admits the representation

$$\text{gph } \mathcal{F} = \{(p, x) \in l_\infty(T) \times X \mid \langle a_t^*, x \rangle - \langle \delta_t, p \rangle \leq b_t \text{ for all } t \in T\}.$$

Therefore, we have  $(p^*, x^*) \in N((\bar{p}, \bar{x}); \text{gph } \mathcal{F})$  if and only if

$$(3.2) \quad \langle p^*, p \rangle + \langle x^*, x \rangle \leq \langle p^*, \bar{p} \rangle + \langle x^*, \bar{x} \rangle$$

for every  $(p, x) \in \text{gph } \mathcal{F}$ . Employing now the equivalence between (i) and (ii) in Lemma 2.1, we conclude that  $(p^*, x^*) \in N((\bar{p}, \bar{x}); \text{gph } \mathcal{F})$  if and only if inclusion (3.1) holds.



To justify the last statement of the proposition, for every set  $A \subset T$  consider its characteristic function  $\chi_A: T \rightarrow \{0, 1\}$  defined by

$$\chi_A(t) := \begin{cases} 1 & \text{if } t \in A, \\ 0 & \text{if } t \notin A. \end{cases}$$

It is obvious that the inclusion  $(p, x) \in \text{gph } \mathcal{F}$  implies that  $(p + \lambda\chi_A, x) \in \text{gph } \mathcal{F}$  for each  $\lambda > 0$ . Replacing now in (3.2) the pair  $(p, x)$  by  $(p + \lambda\chi_A, x)$ , keeping  $p$  and  $x$  fixed, dividing both sides of the inequality by  $\lambda$ , and letting  $\lambda \rightarrow \infty$ , we get the relationships

$$\langle p^*, \chi_A \rangle = \int_T \chi_A(t) p^*(dt) = p^*(A) \leq 0,$$

which complete the proof of the proposition.  $\square$

Based on the above proposition and the general coderivative definition, we now obtain a constructive representation of the coderivative  $D^*\mathcal{F}(0, \bar{x})$  in question.

**THEOREM 3.2** (coderivative of the feasible solution map). *Let  $\bar{x} \in \mathcal{F}(0)$  for the feasible solution map  $\mathcal{F}: l_\infty(T) \rightrightarrows X$  from (1.1) with a Banach decision space  $X$ . Then  $p^* \in D^*\mathcal{F}(0, \bar{x})(x^*)$  if and only if*

$$(3.3) \quad (p^*, -x^*, -\langle x^*, \bar{x} \rangle) \in \text{cl}^* \text{cone}\{(-\delta_t, a_t^*, b_t) \mid t \in T\}.$$

*Proof.* By the coderivative construction (2.3) applied to  $\mathcal{F}$  and by the normal cone formula from Proposition 3.1 as  $\bar{p} = 0$ , we get that  $p^* \in D^*\mathcal{F}(0, \bar{x})(x^*)$  if and only if

$$(3.4) \quad (p^*, -x^*, -\langle x^*, \bar{x} \rangle) \in \text{cl}^* \text{cone}\left[\{(-\delta_t, a_t^*, b_t) \mid t \in T\} \cup \{(0, 0, 1)\}\right].$$

To justify the coderivative representation claimed in the theorem, we need to show that inclusion (3.4) implies, in fact, the “smaller” one in (3.3). Assuming indeed that (3.4) holds, we find by the structure of the right-hand side on (3.4) some nets  $\{\lambda_\nu\}_{\nu \in \mathcal{N}} \subset \mathbb{R}_+^{(T)}$  and  $\{\gamma_\nu\}_{\nu \in \mathcal{N}} \subset \mathbb{R}_+$  satisfying the limiting relationship

$$(3.5) \quad (p^*, -x^*, -\langle x^*, \bar{x} \rangle) = w^* \text{-}\lim_\nu \left( \sum_{t \in T} \lambda_{t\nu} (-\delta_t, a_t^*, b_t) + \gamma_\nu (0, 0, 1) \right),$$

where  $\lambda_{t\nu}$  stands for the  $t$ -entry of  $\lambda_\nu = (\lambda_{t\nu})_{t \in T}$  as  $\nu \in \mathcal{N}$ . It follows directly from the component structure in (3.5) that

$$(3.6) \quad 0 = \langle p^*, 0 \rangle + \langle -x^*, \bar{x} \rangle + (-\langle x^*, \bar{x} \rangle)(-1) = \lim_\nu \left( \sum_{t \in T} \lambda_{t\nu} (\langle a_t^*, \bar{x} \rangle - b_t) - \gamma_\nu \right).$$

Finally, taking into account the definition of the positive cone  $\mathbb{R}_+^{(T)}$  and the fact that the pair  $(0, \bar{x})$  satisfies the infinite inequality system in (1.1), we conclude from (3.6) that  $\lim_\nu \gamma_\nu = 0$ . This justifies (3.3) and thus completes the proof of the theorem.  $\square$

The next consequence of Theorem 3.2 is useful in what follows.

**COROLLARY 3.3** (limiting descriptions of coderivatives). *If  $p^* \in D^*\mathcal{F}(0, \bar{x})(x^*)$  in the framework of Theorem 3.2, then there is a net  $\{\lambda_\nu\}_{\nu \in \mathcal{N}} \subset \mathbb{R}_+^{(T)}$  such that*

$$\begin{aligned} \sum_{t \in T} \lambda_{t\nu} &\rightarrow \|p^*\| = -\langle p^*, e \rangle, \\ \sum_{t \in T} \lambda_{t\nu} a_t^* &\xrightarrow{w^*} -x^*, \quad \text{and} \quad \sum_{t \in T} \lambda_{t\nu} b_t \rightarrow -\langle x^*, \bar{x} \rangle. \end{aligned}$$

*Proof.* It follows from Theorem 3.2 that there is a net  $\{\lambda_\nu\}_{\nu \in \mathcal{N}} \subset \mathbb{R}_+^{(T)}$  such that

$$\sum_{t \in T} \lambda_{t\nu} \delta_t \xrightarrow{w^*} -p^*, \quad \sum_{t \in T} \lambda_{t\nu} a_t^* \xrightarrow{w^*} -x^*, \quad \text{and} \quad \sum_{t \in T} \lambda_{t\nu} b_t \rightarrow -\langle x^*, \bar{x} \rangle,$$

which readily imply the relationships

$$\left\langle \sum_{t \in T} \lambda_{t\nu} \delta_t, e \right\rangle = \sum_{t \in T} \lambda_{t\nu} \rightarrow \langle p^*, -e \rangle =: \lambda \in [0, \infty).$$

Since the dual norm on  $X^*$  is  $w^*$ -lower semicontinuous, we have

$$\|p^*\| \leq \liminf_{\nu \in \mathcal{N}} \left\| \sum_{t \in T} \lambda_{t\nu} \delta_t \right\| \leq \liminf_{\nu \in \mathcal{N}} \sum_{t \in T} \lambda_{t\nu} = \lambda.$$

It follows at the same time from the norm definition that

$$\|p^*\| = \sup_{\|p\| \leq 1} \langle p^*, p \rangle \geq \langle p^*, -e \rangle = \lambda,$$

which finally yields  $\|p^*\| = -\langle p^*, e \rangle$  and completes the proof of the corollary.  $\square$

Our further intention is to provide the *exact calculation* of the *coderivative norm*

$$(3.7) \quad \|D^* \mathcal{F}(0, \bar{x})\| := \sup \{ \|p^*\| \mid p^* \in D^* \mathcal{F}(0, \bar{x})(x^*), \|x^*\| \leq 1 \}$$

in terms of the initial data of the linear infinite inequality system (1.1). A part of our analysis in this direction is the following lemma on properties of the characteristic set (2.8) at  $p = 0$ , which is also used in section 4 to compute the exact Lipschitzian bound  $\text{lip } \mathcal{F}(0, \bar{x})$ .

LEMMA 3.4 (properties of the characteristic set). *Let  $X$  be an arbitrary Banach space. The following assertions hold:*

(i) *Assume that  $\bar{x} \in \mathcal{F}(0)$  is not a strong Slater point for the infinite system (1.1) at  $p = 0$  and that the collection  $\{a_t^* \mid t \in T\}$  is bounded in  $X^*$ . Then the set*

$$(3.8) \quad S := \{x^* \in X^* \mid (x^*, \langle x^*, \bar{x} \rangle) \in \text{cl}^* C(0)\}$$

*built upon the characteristic set  $C(0)$  in (2.8) is nonempty and  $w^*$ -compact at  $X^*$ .*

(ii) *Assume that  $\bar{x} \in \mathcal{F}(0)$  is a strong Slater point of (1.1) at  $p = 0$ . Then the set  $S$  in (3.8) is empty.*

*Proof.* To justify (i), let  $\bar{x}$  be not a strong Slater point for the infinite system (1.1) at  $p = 0$ . Then there is a sequence  $\{t_k\}_{k \in \mathbb{N}} \subset T$  such that  $\lim_k (\langle a_{t_k}^*, \bar{x} \rangle - b_{t_k}) = 0$ . The boundedness of  $\{a_t^* \mid t \in T\}$  implies by the classical Alaoglu–Bourbaki theorem that this set is relatively  $w^*$ -compact in  $X^*$ ; i.e., there is a *subnet*  $\{a_{t_\nu}^*\}_{\nu \in \mathcal{N}}$  of the latter sequence that  $w^*$ -converges to some element  $u^* \in \text{cl}^* \{a_t^* \mid t \in T\}$ . This gives  $\lim_\nu b_{t_\nu} = \langle u^*, \bar{x} \rangle$ , and therefore

$$(u^*, \langle u^*, \bar{x} \rangle) = w^* \text{-} \lim_\nu (a_{t_\nu}^*, b_{t_\nu}) \in \text{cl}^* C(0),$$

which justifies the *nonemptiness* of the set  $S$  in (3.8). Next we prove that  $S$  is  $w^*$ -compact.

Indeed, by our assumption, the set  $A := \{a_t^* \mid t \in T\}$  is bounded in  $X^*$ , and so is  $\text{cl}^* \text{co } A$ ; the latter is actually  $w^*$ -compact due to its automatic  $w^*$ -closedness. Observe

further that the set  $S$  in (3.8) is a *preimage* of  $\text{cl}^*C(0)$  under the  $w^*$ -continuous mapping  $x^* \mapsto (x^*, \langle x^*, \bar{x} \rangle)$ , and thus it is  $w^*$ -closed in  $X^*$ . Since  $S$  is a subset of  $\text{cl}^*\text{co}A$ , it is also bounded and hence  $w^*$ -compact in  $X^*$ . This completes the proof of assertion (i).

To prove assertion (ii), let  $\bar{x}$  be a strong Slater point of system (1.1) at  $p = 0$ , and let

$$\gamma := -\sup_{t \in T} \{ \langle a_t^*, \bar{x} \rangle - b_t \}.$$

Then we have the inequality

$$\langle x^*, \bar{x} \rangle \leq \beta - \gamma \quad \text{whenever } (x^*, \beta) \in \text{cl}^*C(0),$$

which justifies (ii) and thus completes the proof of the lemma.  $\square$

Now we are ready to compute the coderivative norm  $\|D^*\mathcal{F}(0, \bar{x})\|$  at the reference point.

**THEOREM 3.5** (computing the coderivative norm). *Let  $\bar{x} \in \text{dom } \mathcal{F}$  for the infinite system (1.1) with an arbitrary Banach space  $X$  of decision variables. Assume that  $\mathcal{F}$  satisfies the strong Slater condition at  $p = 0$  and that the coefficient set  $\{a_t^* \mid t \in T\}$  is bounded in  $X^*$ . The following assertions hold:*

(i) *If  $\bar{x}$  is a strong Slater point for  $\mathcal{F}$  at  $p = 0$ , then  $\|D^*\mathcal{F}(0, \bar{x})\| = 0$ .*

(ii) *If  $\bar{x}$  is not a strong Slater point for  $\mathcal{F}$  at  $p = 0$ , then the coderivative norm (3.7) is positive and is computed by*

$$(3.9) \quad \|D^*\mathcal{F}(0, \bar{x})\| = \max \left\{ \|x^*\|^{-1} \mid (x^*, \langle x^*, \bar{x} \rangle) \in \text{cl}^*C(0) \right\}$$

via the  $w^*$ -closure of the characteristic set (2.8) at  $p = 0$ .

*Proof.* To justify assertion (i), assume that  $\bar{x}$  is a strong Slater point for the system  $\mathcal{F}$  at  $p = 0$ . It follows from the proof of implication (i) $\implies$ (v) in Lemma 2.3 that we have  $(0, \bar{x}) \in \text{int}(\text{gph } \mathcal{F})$ . It gives

$$N((0, \bar{x}); \text{gph } \mathcal{F}) = \{(0, 0)\},$$

and the conclusion in (i) follows from (2.3) and (3.7).

To prove assertion (ii), take  $x^* \in X^*$  such that  $(x^*, \langle x^*, \bar{x} \rangle) \in \text{cl}^*C(0)$ ; the latter set is nonempty according to Lemma 3.4. Then there exists a net  $\{\lambda_\nu\}_{\nu \in \mathcal{N}} \subset \mathbb{R}_+^{(T)}$  with  $\sum_{t \in T} \lambda_{t\nu} = 1$  for all  $\nu \in \mathcal{N}$  such that

$$\sum_{t \in T} \lambda_{t\nu} a_t^* \xrightarrow{w^*} x^* \quad \text{and} \quad \sum_{t \in T} \lambda_{t\nu} b_t \rightarrow \langle x^*, \bar{x} \rangle.$$

Form further net elements  $p_\nu^* \in l_\infty(T)^*$  by

$$p_\nu^* := -\sum_{t \in T} \lambda_{t\nu} \delta_t, \quad \nu \in \mathcal{N}.$$

Since  $\|p_\nu^*\| = \langle p_\nu^*, -e \rangle = 1$ , the Alaoglu–Bourbaki theorem allows us to select a subnet (without relabeling) such that  $p_\nu^* \xrightarrow{w^*} p^*$  with  $\|p^*\| \leq 1$ . Following the same reasoning as in the proof of Corollary 3.3, we conclude that

$$(3.10) \quad 1 = \lim_{\nu \in \mathcal{N}} \sum_{t \in T} \lambda_{t\nu} = \|p^*\| = \langle p^*, -e \rangle.$$

Moreover, we also obtain the inclusion

$$(p^*, x^*, \langle x^*, \bar{x} \rangle) \in \text{cl}^* \text{co} \{(-\delta_t, a_t^*, b_t) \mid t \in T\},$$

which implies by Theorem 3.2 that

$$(3.11) \quad p^* \in D^* \mathcal{F}(0, \bar{x})(-x^*).$$

Suppose now that  $x^* = 0$  in (3.11). Since  $p^* \neq 0$  by (3.10), we get from (3.11) that

$$(3.12) \quad D^* \mathcal{F}(0, \bar{x})(0) \neq \{0\},$$

which yields by [31, Theorem 1.44] that  $\mathcal{F}$  is *not Lipschitz-like* around  $(0, \bar{x})$ , and therefore it cannot satisfy the strong Slater condition by implication (i) $\implies$ (iv) in Lemma 2.3. This contradicts the assumption of the theorem.

Thus  $x^* \neq 0$  in (3.11), and we derive from the latter relationship that

$$\|x^*\|^{-1} p^* \in D^* \mathcal{F}(0, \bar{x}) \left( -\|x^*\|^{-1} x^* \right),$$

which gives in turn that

$$\|D^* \mathcal{F}(0, \bar{x})\| \geq \left\| \|x^*\|^{-1} p^* \right\| = \|x^*\|^{-1}$$

and hence justifies the inequality “ $\geq$ ” in (3.9).

It remains to prove the opposite inequality in (3.9). For the nonempty and  $w^*$ -compact set  $S$  in (3.8) we have  $0 \notin S$  by Lemma 2.3, which ensures the  $w^*$ -upper semicontinuity of the function  $x^* \mapsto \|x^*\|^{-1}$  on  $S$ . Thus the supremum in the right-hand side of (3.9) is attained and belongs to  $(0, \infty)$ . Then condition (v) in Lemma 2.3 implies that  $(0, \hat{x}) \in \text{int}(\text{gph } \mathcal{F})$  for some  $\hat{x} \in X$ , and so  $0 \in \text{int}(\text{dom } \mathcal{F})$ . Moreover, we have that  $p^* \in D^* \mathcal{F}(0, \bar{x})(-x^*)$  if and only if  $(p^*, x^*) \in N((0, \bar{x}); \text{gph } \mathcal{F})$ , which is equivalent to

$$(3.13) \quad \langle p^*, p \rangle + \langle x^*, x \rangle \leq \langle x^*, \bar{x} \rangle \quad \text{for all } (p, x) \in \text{gph } \mathcal{F}.$$

This allows us, by taking into account that  $0 \in \text{int}(\text{dom } \mathcal{F})$ , to arrive at the equivalences

$$(3.14) \quad p^* \in D^* \mathcal{F}(0, \bar{x})(0) \iff \langle p^*, p \rangle \leq 0 \text{ for all } p \in \text{dom } \mathcal{F} \iff p^* = 0.$$

Observe, furthermore, that, since  $\bar{x}$  is not a strong Slater point for  $\mathcal{F}$  at  $p = 0$ , we have  $(0, \bar{x}) \notin \text{int}(\text{gph } \mathcal{F})$  and thus conclude by the classical separation theorem that there is a pair  $(p^*, x^*) \neq (0, 0)$  for which relation (3.13) holds. Employing (3.14), we have  $x^* \neq 0$  and  $p^* \in D^* \mathcal{F}(0, \bar{x})(-x^*)$ .

Take now  $p^* \in D^* \mathcal{F}(0, \bar{x})(-x^*)$  with  $\|x^*\| \leq 1$  and suppose that  $x^* \neq 0$ ; the arguments of the previous paragraph ensure the existence of such an element. By Corollary 3.3 there is a net  $\{\lambda_\nu\}_{\nu \in \mathcal{N}} \subset \mathbb{R}_+^{(T)}$  for which

$$\begin{aligned} \gamma_\nu &:= \sum_{t \in T} \lambda_{t\nu} \rightarrow \|p^*\| = -\langle p^*, e \rangle, \\ x_\nu^* &:= \sum_{t \in T} \lambda_{t\nu} a_t^* \xrightarrow{w^*} x^*, \quad \text{and} \quad \sum_{t \in T} \lambda_{t\nu} b_t \rightarrow \langle x^*, \bar{x} \rangle. \end{aligned}$$

Taking  $M \geq \|a_t^*\|$  for every  $t \in T$ , we get the estimate

$$\|x_\nu^*\| \leq M\gamma_\nu \quad \text{whenever } \nu \in \mathcal{N}$$

and also the limiting relationships

$$0 < \|x^*\| \leq \liminf_{\nu \in \mathcal{N}} \|x_\nu^*\| \leq M \liminf_{\nu \in \mathcal{N}} \gamma_\nu = M \|p^*\|,$$

which ensure that  $p^* \neq 0$ . It follows, furthermore, that

$$\|p^*\|^{-1} (x^*, \langle x^*, \bar{x} \rangle) \in \text{cl}^* C(0).$$

Remembering, finally, that  $0 < \|x^*\| \leq 1$ , we arrive at the estimates

$$\|p^*\| \leq \left\| \|p^*\|^{-1} x^* \right\|^{-1} \leq \max \left\{ \|u^*\|^{-1} \mid (u^*, \langle u^*, \bar{x} \rangle) \in \text{cl}^* C(0) \right\},$$

which justify the inequality “ $\leq$ ” in (3.9) and complete the proof of the theorem.  $\square$

**4. Characterizations of robust Lipschitzian stability for feasible solution maps.** In this section we employ the above coderivative analysis combined with appropriate techniques developed in linear semi-infinite/infinite programming to establish a *coderivative characterization* of robust Lipschitzian stability, in the sense discussed in section 2, for the infinite inequality system  $\mathcal{F}$  in (1.1) at the reference point  $(0, \bar{x})$  with computing the *exact Lipschitzian bound*  $\text{lip } \mathcal{F}(0, \bar{x})$ .

The first result of this section establishes the coderivative *necessary and sufficient* condition in form (2.4) for the Lipschitz-like property of  $\mathcal{F}$  around  $(0, \bar{x}) \in \text{gph } \mathcal{F}$  in the general setting under consideration.

**THEOREM 4.1** (coderivative criterion for robust Lipschitzian stability of linear infinite inequalities). *Let  $\bar{x} \in \mathcal{F}(0)$  for the infinite inequality system (1.1) with a Banach space  $X$  of decision variables. Then  $\mathcal{F}$  is Lipschitz-like around  $(0, \bar{x})$  if and only if*

$$(4.1) \quad D^* \mathcal{F}(0, \bar{x})(0) = \{0\}.$$

*Proof.* The “only if” part follows from [31, Theorem 1.44] specified for the mapping  $\mathcal{F}: l_\infty(T) \rightrightarrows X$  under consideration. Let us now prove the “if” part of the theorem.

Arguing by contradiction, suppose that  $D^* \mathcal{F}(0, \bar{x})(0) = \{0\}$  while the mapping  $\mathcal{F}$  is *not* Lipschitz-like around  $(0, \bar{x})$ . Then, by the equivalence between properties (ii) and (iv) of Lemma 2.3, we get the inclusion

$$(0, 0) \in \text{cl}^* \text{co} \{ (a_t^*, b_t) \in X^* \times \mathbb{R} \mid t \in T \},$$

which means that there is a net  $\{\lambda_\nu\}_{\nu \in \mathcal{N}} \in \mathbb{R}_+^{(T)}$  such that  $\sum_{t \in T} \lambda_{t\nu} = 1$  for all  $\nu \in \mathcal{N}$  and

$$(4.2) \quad w^* \text{-} \lim_{\nu} \sum_{t \in T} \lambda_{t\nu} (a_t^*, b_t) = (0, 0).$$

Since the net  $\{\sum_{t \in T} \lambda_{t\nu} (-\delta_t)\}_{\nu \in \mathcal{N}}$  is obviously bounded in  $l_\infty(T)^*$ , the Alaoglu–Bourbaki theorem ensures the existence of its subnet (with no relabeling) that  $w^*$ -converges to some element  $p^* \in l_\infty(T)^*$ , i.e.,

$$(4.3) \quad p^* = w^* \text{-} \lim_{\nu} \sum_{t \in T} \lambda_{t\nu} (-\delta_t).$$

It follows from (4.3) by the Dirac function definition that

$$\langle p^*, -e \rangle = \lim_{\nu \in \mathcal{N}} \sum_{t \in T} \lambda_{t\nu} = 1, \text{ where } e = (e_t)_{t \in T} \text{ with } e_t = 1 \text{ for all } t \in T,$$

which implies that  $p^* \neq 0$ . It follows from (4.2) and (4.3) that

$$(p^*, 0, 0) = w^* \text{-}\lim_{\nu} \sum_{t \in T} \lambda_{t\nu} (-\delta_t, a_t^*, b_t) \text{ with } p^* \neq 0,$$

and therefore, by the explicit coderivative description of Theorem 3.2, we get the inclusion  $p^* \in D^* \mathcal{F}(0, \bar{x})(0) \setminus \{0\}$ , which contradicts the assumed condition (4.1). This justifies the sufficiency of the coderivative condition (4.1) for the Lipschitz-like property of  $\mathcal{F}$  around  $(0, \bar{x})$  and thus completes the proof of the theorem.  $\square$

Our further goal is to compute the *exact Lipschitzian bound*  $\text{lip } \mathcal{F}(0, \bar{x})$  of  $\mathcal{F}$  around  $(0, \bar{x})$ . We are going to do it according to the *distance representation* (2.2) for the exact Lipschitzian bound. To proceed, denote

$$(4.4) \quad H(x^*, \alpha) := \{x \in X \mid \langle x^*, x \rangle \leq \alpha\} \text{ for } (x^*, \alpha) \in X^* \times \mathbb{R},$$

and observe the following representation (known as the *Ascoli formula*; see, e.g., [2]):

$$(4.5) \quad \text{dist}(x; H(x^*, \alpha)) = \frac{[\langle x^*, x \rangle - \alpha]_+}{\|x^*\|},$$

where  $[\gamma]_+ := \max\{\gamma, 0\}$  for  $\gamma \in \mathbb{R}$ . Recall that we are under the convention  $0/0 := 0$ .

The next proposition and the subsequent lemma, which are certainly of their own interest, provide a significant extension of the Ascoli formula (4.4) to the case of *infinite* systems of linear inequalities instead of the single one as in (4.4). These results are essentially employed in what follows for computing the exact Lipschitzian bound  $\text{lip } \mathcal{F}(0, \bar{x})$ . We refer the reader to [6, Lemma 2.3] and [7, Lemma 1] for related results in the framework of semi-infinite programming and observe that in infinite dimensions we use the *w\*-closure* of the characteristic sets  $C(p)$  from (2.8); see also Example 4.4 below for more discussion.

From now on, given an extended-real-valued function  $f: X \rightarrow \overline{\mathbb{R}} := (-\infty, \infty]$  assumed to be *proper* (i.e., not identically equal to  $\infty$ ), consider its *Fenchel conjugate*  $f^*: X^* \rightarrow \overline{\mathbb{R}}$  defined by

$$f^*(x^*) := \sup \{ \langle x^*, x \rangle - f(x) \mid x \in X \} = \sup \{ \langle x^*, x \rangle - f(x) \mid x \in \text{dom } f \},$$

where  $\text{dom } f := \{x \in X \mid f(x) < \infty\}$  is the *effective domain* of  $f$ . In what follows we also use the standard notation  $\text{epi } f$  for the *epigraph* of  $f$  given by

$$\text{epi } f := \{(x, \gamma) \in X \times \mathbb{R} \mid x \in \text{dom } f, f(x) \leq \gamma\}.$$

The following result gives an *exact formula* for computing the distance to a set defined by a convex inequality via the corresponding conjugate function.

**PROPOSITION 4.2** (distance function for solutions of convex inequalities in Banach spaces). *Let  $g: X \rightarrow \overline{\mathbb{R}}$  be a proper convex function on a Banach space  $X$ , and let*

$$(4.6) \quad Q := \{y \in X \mid g(y) \leq 0\}.$$

Assume the fulfillment of the classical Slater condition: there is  $\hat{x} \in X$  such that  $g(\hat{x}) < 0$ . Then the distance function to the set  $Q$  in (4.6) is computed by

$$(4.7) \quad \text{dist}(x; Q) = \max_{(x^*, \alpha) \in \text{epi } g^*} \frac{[\langle x^*, x \rangle - \alpha]_+}{\|x^*\|}.$$

*Proof.* Observe that the nonemptiness of  $Q$  in (4.6) yields that  $\alpha \geq 0$  whenever  $(0, \alpha) \in \text{epi } g^*$  and that the possibility of  $x^* = 0$  is not an obstacle in (4.7) under our convention  $0/0 = 0$ . Obviously the distance function  $\text{dist}(x; Q)$  is nothing else but the optimal value function in the parametric *convex optimization* problem

$$(4.8) \quad \text{minimize } \|y - x\| \quad \text{subject to } g(y) \leq 0.$$

Since the classical Slater condition holds for problem (4.8) by our assumption, we have the *strong Lagrange duality* in (4.8) by, e.g., [39, Theorem 2.9.3], which gives

$$\begin{aligned} \text{dist}(x; Q) &= \max_{\lambda \geq 0} \inf_{y \in X} \{ \|y - x\| + \lambda g(y) \} \\ &= \max \left\{ \max_{\lambda > 0} \inf_{y \in X} \{ \|y - x\| + \lambda g(y) \}, \inf_{y \in X} \|y - x\| \right\} \\ &= \max \left\{ \max_{\lambda > 0} \inf_{y \in X} \{ \|y - x\| + \lambda g(y) \}, 0 \right\}. \end{aligned}$$

Applying the classical *Fenchel duality theorem* to the inner infimum problem above for a fixed  $\lambda > 0$  (observing to this end that Rockafellar’s regularity condition needed for Fenchel duality is satisfied for this problem due to the continuity of the norm function), we get

$$(4.9) \quad \inf_{y \in X} \{ \|y - x\| + \lambda g(y) \} = \max_{y^* \in X^*} \{ -\|\cdot - x\|^*(-y^*) - (\lambda g)^*(y^*) \}.$$

It is well known in convex analysis that

$$\|\cdot - x\|^*(-y^*) = \begin{cases} \langle -y^*, x \rangle & \text{if } \|y^*\| \leq 1, \\ \infty & \text{otherwise.} \end{cases}$$

Implementing this into formula (4.9) and employing elementary transformations, we obtain

$$\begin{aligned} \inf_{y \in X} \{ \|y - x\| + \lambda g(y) \} &= \max_{\|y^*\| \leq 1} \{ \langle y^*, x \rangle - (\lambda g)^*(y^*) \} \\ &= \max_{\|y^*\| \leq 1, (\lambda g)^*(y^*) \leq \eta} \{ \langle y^*, x \rangle - \eta \} \\ &= \max_{\|y^*\| \leq 1, \lambda g^*(y^*/\lambda) \leq \eta} \{ \langle y^*, x \rangle - \eta \} \\ &= \max_{\|y^*\| \leq 1, (1/\lambda)(y^*, \eta) \in \text{epi } g^*} \{ \langle y^*, x \rangle - \eta \}. \end{aligned}$$

The latter representation yields, by denoting  $x^* := (1/\lambda)y^*$  and  $\alpha := (1/\lambda)\eta$ , that

$$\inf_{y \in X} \{ \|y - x\| + \lambda g(y) \} = \max_{(x^*, \alpha) \in \text{epi } g^*, \|x^*\| \leq 1/\lambda} \lambda \{ \langle x^*, x \rangle - \alpha \}.$$

Combining this with the formulas above, we arrive at

$$(4.10) \quad \begin{aligned} \text{dist}(x; Q) &= \max \left\{ \max_{(x^*, \alpha) \in \text{epi } g^*, \|x^*\| \leq 1/\lambda} \lambda \{ \langle x^*, x \rangle - \alpha \}, 0 \right\} \\ &= \max_{(x^*, \alpha) \in \text{epi } g^*, \|x^*\| \leq 1/\lambda} \left\{ \lambda [\langle x^*, x \rangle - \alpha]_+ \right\}. \end{aligned}$$

It is easy to observe the following relationships held for any  $\lambda > 0$ :

$$\begin{aligned} \max_{(0,\alpha) \in \text{epi } g^*} \lambda \{ \langle 0, x \rangle - \alpha \} &= \max_{g^*(0) \leq \alpha} \lambda (\langle 0, x \rangle - \alpha) \\ &= \lambda (-g^*(0)) \leq \lambda \inf_{x \in X} g(x) \leq \lambda g(\hat{x}) < 0. \end{aligned}$$

Taking this into account, we have from (4.10) that

$$\begin{aligned} \text{dist}(x; Q) &= \max_{(x^*, \alpha) \in \text{epi } g^*, \|x^*\| \leq 1/\lambda} \left\{ \lambda [\langle x^*, x \rangle - \alpha]_+ \right\} \\ &= \max_{(x^*, \alpha) \in \text{epi } g^*} \max_{\|x^*\| \leq 1/\lambda} \left\{ \lambda [\langle x^*, x \rangle - \alpha]_+ \right\} \\ &= \max_{(x^*, \alpha) \in \text{epi } g^*} \frac{[\langle x^*, x \rangle - \alpha]_+}{\|x^*\|}, \end{aligned}$$

which gives (4.7) and thus completes the proof of the proposition.  $\square$

Now we are ready we establish the required extension of the Ascoli formula (4.5) to the case of the infinite inequality systems under consideration.

LEMMA 4.3 (distance to infinite linear inequalities in Banach spaces). *Assume that the infinite system  $\mathcal{F}(p)$  in (1.1) satisfies the strong Slater condition at  $p = (p_t)_{t \in T}$ . Then for any  $x \in X$  and any  $p \in l_\infty(T)$  we have the extended Ascoli distance formula*

$$(4.11) \quad \text{dist}(x; \mathcal{F}(p)) = \max_{(x^*, \alpha) \in \text{cl}^* C(p)} \frac{[\langle x^*, x \rangle - \alpha]_+}{\|x^*\|}.$$

If, furthermore,  $X$  is reflexive, then the distance formula (4.11) can be simplified as follows:

$$(4.12) \quad \text{dist}(x; \mathcal{F}(p)) = \sup_{(x^*, \alpha) \in C(p)} \frac{[\langle x^*, x \rangle - \alpha]_+}{\|x^*\|}.$$

*Proof.* It is obvious to observe that the infinite system (1.1) is represented in

$$(4.13) \quad \mathcal{F}(p) = \{x \in X \mid g(x) \leq 0\},$$

where the convex function  $g: X \rightarrow \overline{\mathbb{R}}$  is given in the supremum form

$$(4.14) \quad g(x) := \sup_{t \in T} (f_t(x) - p_t) \quad \text{with} \quad f_t(x) := \langle a_t^*, x \rangle - b_t.$$

The assumed strong Slater condition for  $\mathcal{F}(p)$  ensures the fulfillment of the classical Slater condition for  $g$  imposed in Proposition 4.2. To imply this proposition in the framework of (4.13), we need to compute the conjugate function to the supremum function in (4.14). The recent results in this direction [4, 18] yield that

$$\text{epi } g^* = \text{epi} \left\{ \sup_{t \in T} (f_t - p_t) \right\}^* = \text{cl}^* \text{co} \left( \bigcup_{t \in T} \text{epi} (f_t - p_t)^* \right) = \text{cl}^* C(p) + \mathbb{R}_+(0, 1)$$

with  $0 \in X^*$ , where the weak\* closedness of the set  $\text{cl}^* C(p) + \mathbb{R}_+(0, 1)$  is a consequence of the classical Dieudonné theorem; see, e.g., [39, Theorem 1.1.8]. Thus we get the distance formula (4.11) from Proposition 4.2 in general Banach spaces.



To prove the simplified distance formula (4.12) in the case of reflexive spaces, suppose, on the contrary, that it does not hold. Then there is a scalar  $\beta \in \mathbb{R}$  such that

$$(4.15) \quad \max_{(x^*, \alpha) \in \text{cl}^* C(p)} \frac{[\langle x^*, x \rangle - \alpha]_+}{\|x^*\|} > \beta > \sup_{(x^*, \alpha) \in C(p)} \frac{[\langle x^*, x \rangle - \alpha]_+}{\|x^*\|}.$$

This yields the existence of a pair  $(\bar{x}^*, \bar{\alpha}) \in \text{cl}^* C(p)$  with  $\bar{x}^* \in X^* \setminus \{0\}$  and  $\bar{\alpha} \in \mathbb{R}$  such that

$$\frac{[\langle \bar{x}^*, x \rangle - \bar{\alpha}]_+}{\|\bar{x}^*\|} > \beta.$$

Taking into account that the space  $X$  is reflexive and that the set  $C(p)$  is convex and employing the Mazur weak closure theorem, we can replace the weak\* closure of the  $C(p)$  above by its norm closure in  $X^*$ . This allows us to find a sequence  $(x_k^*, \alpha_k) \in C(p)$ ,  $k \in \mathbb{N}$ , converging in norm to  $(\bar{x}^*, \bar{\alpha})$  as  $k \rightarrow \infty$ . Thus we get

$$\lim_{k \rightarrow \infty} \frac{[\langle x_k^*, x \rangle - \alpha_k]_+}{\|x_k^*\|} = \frac{[\langle \bar{x}^*, x \rangle - \bar{\alpha}]_+}{\|\bar{x}^*\|} > \beta,$$

and therefore there is  $k_0 \in \mathbb{N}$  such that

$$\frac{[\langle x_{k_0}^*, x \rangle - \alpha_{k_0}]_+}{\|x_{k_0}^*\|} > \beta.$$

The latter surely contradicts (4.15), and this completes the proof of the lemma.  $\square$

The following example shows that the *reflexivity* of the decision space  $X$  is an *essential* requirement for the validity of the simplified distance formula (4.12), even in the framework of (nonreflexive) *Asplund* spaces.

*Example 4.4* (failure of simplified distance formula in nonreflexive Asplund spaces). Consider the classical space  $c_0$  of sequences of real numbers converging to zero endowed with the supremum norm. This space is well known to be *Asplund* while *not reflexive*; see, e.g., [17]. Let us show that the simplified distance formula (4.12) *fails* in  $X = c_0$  for a rather plain linear system of *countable* inequalities. Of course, we need to demonstrate that the inequality “ $\leq$ ” is generally violated in (4.12), since the opposite inequality holds in any Banach space. Form the infinite (countable) linear inequality system

$$(4.16) \quad \mathcal{F}(0) := \{x \in c_0 \mid \langle e_1^* + e_t^*, x \rangle \leq -1, t \in \mathbb{N}\},$$

where  $e_t^* \in l_1$  has 1 as its  $t$ th component while all the remaining components are 0. System (4.16) can be rewritten as

$$x \in \mathcal{F}(0) \iff x(1) + x(t) \leq -1 \text{ for all } t \in \mathbb{N}.$$

Observe that for the origin  $z = 0$  we have  $\text{dist}(0; \mathcal{F}(0)) = 1$ , and the distance is realized at, e.g.,  $u = (-1, 0, 0, \dots)$ . Indeed, passing to the limit in the inequality

$$x(1) + x(t) \leq -1 \text{ as } t \rightarrow \infty$$

and taking into account that  $x(t) \rightarrow 0$  as  $t \rightarrow \infty$ , by the structure of the space of  $c_0$ , we get  $x(1) \leq -1$ . Furthermore, it can be checked that

$$\begin{aligned} (e_1^*, -1) \in \text{cl}^*C(0), \quad \langle e_1^*, x - u \rangle \leq 0 \quad \text{for all } x \in \mathcal{F}(0), \\ \text{dist}(z; \mathcal{F}(0)) = \|z - u\| = \langle e_1^*, z - u \rangle = \frac{\langle e_1^*, z \rangle - (-1)}{\|e_1^*\|}. \end{aligned}$$

On the other hand, for the pair  $(x^*, \alpha) \in X^* \times \mathbb{R}$  given by

$$(x^*, \alpha) := \left( e_1^* + \sum_{t \in \mathbb{N}} \lambda_t e_t^*, -1 \right) \in C(0) \quad \text{with } \lambda \in \mathbb{R}_+^{(\mathbb{N})} \quad \text{and} \quad \sum_{t \in \mathbb{N}} \lambda_t = 1,$$

we can directly check that  $\|x^*\| = 2$  and hence

$$\frac{[\langle x^*, z \rangle - \alpha]_+}{\|x^*\|} = \frac{1}{2},$$

which shows that the equality in (4.12) is violated for the countable system (4.16) in the nonreflexive Asplund space  $X = c_0$  of decision variables.

Our next step is to derive a verifiable precise formula for computing the *exact Lipschitzian bound*  $\text{lip } \mathcal{F}(0, \bar{x})$  for the infinite system (1.1) in the general Banach space  $X$ . As a preliminary result we need the following technical lemma.

LEMMA 4.5 (closed-graph property of the characteristic set mapping). *The set-valued mapping  $l_\infty(T) \ni p \mapsto \text{cl}^*C(p) \subset X^* \times \mathbb{R}$  is closed-graph in the norm  $\times$  weak\* topology of  $l_\infty(T) \times (X^* \times \mathbb{R})$ ; i.e., for any nets*

$$\{p_\nu\}_{\nu \in \mathcal{N}} \subset l_\infty(T), \quad \{x_\nu^*\}_{\nu \in \mathcal{N}} \subset X^*, \quad \{\beta_\nu\}_{\nu \in \mathcal{N}} \subset \mathbb{R}$$

satisfying  $p_\nu \rightarrow p$ ,  $x_\nu^* \xrightarrow{w^*} x^*$ ,  $\beta_\nu \rightarrow \beta$ , and  $(x_\nu^*, \beta_\nu) \in \text{cl}^*C(p_\nu)$  for every  $\nu \in \mathcal{N}$  we have the inclusion  $(x^*, \beta) \in \text{cl}^*C(p)$ .

*Proof.* Arguing by contradiction, suppose that  $(x^*, \beta) \notin \text{cl}^*C(p)$ . Then the classical strict separation theorem allows us to find a pair  $(x, \alpha) \in X \times \mathbb{R}$  with  $(x, \alpha) \neq (0, 0)$  and real numbers  $\gamma$  and  $\gamma'$  satisfying

$$\langle x^*, x \rangle + \beta \alpha < \gamma' < \gamma \leq \langle a_t^*, x \rangle + (b_t + p_t) \alpha \quad \text{for all } t \in T.$$

Hence there exists a net index  $\nu_0 \in \mathcal{N}$  such that

$$\langle x_\nu^*, x \rangle + \beta_\nu \alpha < \gamma' \quad \text{and} \quad \|\alpha(p - p_\nu)\| \leq \gamma - \gamma' \quad \text{whenever } \nu \succeq \nu_0.$$

We have, therefore, that

$$\begin{aligned} \langle a_t^*, x \rangle + \alpha(b_t + p_{t\nu}) &= \langle a_t^*, x \rangle + \alpha(b_t + p_t) + \alpha(p_{t\nu} - p_t) \\ &\geq \gamma - \|\alpha(p_\nu - p)\| \geq \gamma' \quad \text{for all } t \in T. \end{aligned}$$

The latter implies that  $\gamma' \leq \langle z^*, x \rangle + \eta \alpha$  for all  $(z^*, \eta) \in \text{cl}^*C(p_\nu)$  whenever  $\nu \succeq \nu_0$ . Thus we arrive at the contradiction

$$\langle x_\nu^*, x \rangle + \beta_\nu \alpha < \gamma' \leq \langle x_\nu^*, x \rangle + \beta_\nu \alpha, \quad \nu \succeq \nu_0,$$

which completes the proof of the lemma.  $\square$

Now we are ready to establish the desired formula for computing the exact Lipschitzian bound of  $\mathcal{F}$  around  $(0, \bar{x})$ .

**THEOREM 4.6** (computing the exact Lipschitzian bound). *Let  $\bar{x} \in \mathcal{F}(0)$  for the linear infinite inequality system (1.1) with a Banach decision space  $X$ . Assume that  $\mathcal{F}$  satisfies the strong Slater condition at  $p = 0$  and that the coefficient set  $\{a_t^* \mid t \in T\}$  is bounded in  $X^*$ . The following assertions hold:*

(i) *If  $\bar{x}$  is a strong Slater point for  $\mathcal{F}$  at  $p = 0$ , then  $\text{lip } \mathcal{F}(0, \bar{x}) = 0$ .*

(ii) *If  $\bar{x}$  is not a strong Slater point for  $\mathcal{F}$  at  $p = 0$ , then the exact Lipschitzian bound of  $\mathcal{F}$  around  $(0, \bar{x})$  is computed by*

$$(4.17) \quad \text{lip } \mathcal{F}(0, \bar{x}) = \max \{ \|x^*\|^{-1} \mid (x^*, \langle x^*, \bar{x} \rangle) \in \text{cl}^* C(0) \} > 0$$

via the  $w^*$ -closure of the characteristic set (2.8) at  $p = 0$ .

*Proof.* Let us first justify (i). We have seen in the proof of Theorem 3.5(i) that our current assumptions imply that  $(0, \bar{x}) \in \text{int}(\text{gph } \mathcal{F})$ , which in turn yields by the definition of the exact Lipschitzian bound that  $\text{lip } \mathcal{F}(0, \bar{x}) = 0$  in this case.

Next we prove the more difficult assertion (ii) of the theorem, assuming that  $\bar{x}$  is not a strong Slater point for  $\mathcal{F}$  at  $p = 0$ . Observe that by Lemma 3.4 the set (3.8) under the maximum operation on the right-hand side in (4.17) is nonempty and  $w^*$ -compact in  $X^*$  and the maximum over this set is realized and hence it is finite. The inequality “ $\geq$ ” in (4.17) follows from the estimate

$$\text{lip } \mathcal{F}(0, \bar{x}) \geq \|D^* \mathcal{F}(0, \bar{x})\|$$

established for general mappings between Banach spaces in [31, Theorem 1.44] and from formula (3.9) for computing the coderivative norm of the inequality system  $\mathcal{F}$  in (1.1) derived above in Theorem 3.5. It remains to prove the opposite inequality “ $\leq$ ” in (4.17).

To proceed, let  $M := \sup_{t \in T} \|a_t^*\| < \infty$ , and observe that the inequality “ $\leq$ ” in (4.17) is obvious when  $L := \text{lip } \mathcal{F}(0, \bar{x}) = 0$ . Suppose now that  $L > 0$ , and consider any pair  $(p, x)$  sufficiently close to  $(0, \bar{x})$  in the limiting representation (2.2) of the exact Lipschitzian bound  $\text{lip } \mathcal{F}(0, \bar{x})$ . We can confine ourselves to the case of  $(p, x) \notin \text{gph } \mathcal{F}$  by  $L > 0$ . Furthermore, it follows from the structure of  $\mathcal{F}$  in (1.1) that

$$(4.18) \quad 0 < \text{dist}(p; \mathcal{F}^{-1}(x)) = \sup_{t \in T} [\langle a_t^*, x \rangle - b_t - p_t]_+.$$

Moreover, we have the relationships

$$\begin{aligned} \langle a_t^*, x \rangle - b_t - p_t &= \langle a_t^*, x - \bar{x} \rangle + \langle a_t^*, \bar{x} \rangle - b_t - p_t \\ &\leq M \|x - \bar{x}\| + \|p\| \quad \text{for all } t \in T, \end{aligned}$$

which allow us to conclude that

$$(4.19) \quad \begin{aligned} 0 &< \sup_{(x^*, \beta) \in \text{cl}^* C(p)} [\langle x^*, x \rangle - \beta]_+ = \sup_{(x^*, \beta) \in \text{cl}^* C(p)} \{ \langle x^*, x \rangle - \beta \} \\ &\leq M \|x - \bar{x}\| + \|p\|. \end{aligned}$$

Consider further the set

$$C_+(p, x) := \{ (x^*, \beta) \in \text{cl}^* C(p) \mid \langle x^*, x \rangle - \beta > 0 \},$$

which is obviously nonempty, and denote

$$M_{(p,x)} := \sup \{ \|x^*\|^{-1} \mid (x^*, \beta) \in C_+(p, x) \}.$$

In our setting we have  $0 \in \text{int}(\text{dom } \mathcal{F})$  (cf. Lemma 2.3 and Remark 2.4), and therefore  $p \in \text{dom } \mathcal{F}$  for all  $p \in l_\infty(T)$  sufficiently close to the origin. In this case the set  $C_+(p, x)$  cannot contain any element of the form  $(0, \beta)$ , since the contrary would yield by the definition of  $C_+(p, x)$  that  $\beta < 0$ , while Lemma 2.1 gives  $\beta \geq 0$ . Thus we get  $0 < \|x^*\| \leq M$  whenever  $(x^*, \beta) \in C_+(p, x)$  and, in particular,  $M_{(p,x)} \in (0, \infty]$ . It follows, furthermore, that

$$\frac{\sup_{(x^*, \beta) \in \text{cl}^* C(p)} \frac{[\langle x^*, x \rangle - \beta]_+}{\|x^*\|}}{\sup_{(x^*, \beta) \in \text{cl}^* C(p)} [\langle x^*, x \rangle - \beta]_+} = \frac{\sup_{(x^*, \beta) \in \text{cl}^* C(p)} \frac{\langle x^*, x \rangle - \beta}{\|x^*\|}}{\sup_{(x^*, \beta) \in \text{cl}^* C(p)} \{\langle x^*, x \rangle - \beta\}} \leq M_{(p,x)},$$

where the latter inequality implies the estimate

$$L \leq \limsup_{(p,x) \rightarrow (0, \bar{x}), x \notin \mathcal{F}(p) \neq \emptyset} M_{(p,x)} := K.$$

Considering next a sequence  $(p_k, x_k) \rightarrow (0, \bar{x})$  with  $x_k \notin \mathcal{F}(p_k) \neq \emptyset$  and

$$L \leq \lim_{k \rightarrow \infty} M_{(p_k, x_k)} = K,$$

we select a sequence  $\{\alpha_k\}_{k=1}^\infty \subset \mathbb{R}$  such that

$$\lim_{k \rightarrow \infty} \alpha_k = K \text{ and } 0 < \alpha_k < M_{(p_k, x_k)} \text{ as } k \in \mathbb{N}.$$

Take now  $(x_k^*, \beta_k) \in C_+(p_k, x_k)$  with  $\alpha_k < \|x_k^*\|^{-1}$  for all  $k \in \mathbb{N}$ . Since the sequence  $\{x_k^*\}_{k \in \mathbb{N}} \subset X^*$  is bounded, it contains a subnet  $\{x_\nu^*\}_{\nu \in \mathcal{N}}$  that  $w^*$ -converges to some  $x^* \in X^*$ . Denoting by  $\{p_\nu\}$ ,  $\{x_\nu\}$ ,  $\{\beta_\nu\}$ , and  $\{\alpha_\nu\}$  the corresponding subnets of  $\{p_k\}$ ,  $\{x_k\}$ ,  $\{\beta_k\}$ , and  $\{\alpha_k\}$ , we get from (4.19) that

$$0 < \langle x_\nu^*, x_\nu \rangle - \beta_\nu \leq M \|x_\nu - \bar{x}\| + \|p_\nu\|.$$

Thus  $\langle x_\nu^*, x_\nu \rangle - \beta_\nu \rightarrow 0$ , which implies by the construction above that  $\beta_\nu \rightarrow \langle x^*, \bar{x} \rangle$ . By Lemma 4.5 we get that

$$(x^*, \langle x^*, \bar{x} \rangle) \in \text{cl}^* C(0),$$

and then Lemma 2.3 ensures that  $x^* \neq 0$ .

To conclude proving the inequality “ $\leq$ ” in (4.17), observe that

$$\|x^*\| \leq \liminf_{\nu \in \mathcal{N}} \|x_\nu^*\| \leq \lim_{\nu} \frac{1}{\alpha_\nu} = \frac{1}{K}$$

due to  $\|x_\nu^*\| \leq \alpha_\nu^{-1}$  and  $\lim_{\nu} \alpha_\nu = K$ , which gives

$$L \leq K \leq \frac{1}{\|x^*\|} \leq \max \{ \|z^*\|^{-1} \mid (z^*, \langle z^*, \bar{x} \rangle) \in \text{cl}^* C(0) \}.$$

Remembering the notation above, we complete the proof of the theorem.  $\square$

Comparing, finally, the results on computing the coderivative norm in Theorem 3.5 and the exact Lipschitzian bound in Theorem 4.6 allows us to arrive at the *unconditional* relationship between the coderivative norm and exact Lipschitzian bound of the infinite inequality system  $\mathcal{F}$  with arbitrary Banach decision spaces. This

was known before only for set-valued mappings between finite-dimensional spaces; cf. formula (2.5) and the corresponding discussions in section 2.

**COROLLARY 4.7** (relationship between the exact Lipschitzian bound and coderivative norm). *Let  $\bar{x} \in \mathcal{F}(0)$  for the infinite system (1.1) satisfying the strong Slater condition at  $p = 0$ . Assume that the decision space  $X$  is arbitrary Banach and that the coefficient set  $\{a_t^* \mid t \in T\}$  is bounded in  $X^*$ . Then*

$$(4.20) \quad \text{lip } \mathcal{F}(0, \bar{x}) = \|D^* \mathcal{F}(0, \bar{x})\|.$$

*Proof.* If  $\bar{x}$  is a strong Slater point for  $\mathcal{F}$  at  $p = 0$ , then we get equality (4.20) directly by comparing assertions (i) in Theorems 3.5 and 4.6 which ensure that

$$\text{lip } \mathcal{F}(0, \bar{x}) = \|D^* \mathcal{F}(0, \bar{x})\| = 0.$$

On the other hand, if  $\bar{x}$  is not a strong Slater point for  $\mathcal{F}$  at  $p = 0$ , then (4.20) follows from comparing assertions (ii) in Theorems 3.5 and 4.6 which justify the *same formula* for computing  $\|D^* \mathcal{F}(0, \bar{x})\|$  and  $\text{lip } \mathcal{F}(0, \bar{x})$  in (3.9) and (4.17), respectively.  $\square$

**Acknowledgments.** The main results of this paper were obtained in our research report [10], which was widely disseminated in the optimization community. We would like to thank many colleagues and friends for their helpful comments and remarks. Especially important contributions came from Radu Boţ, who suggested a new proof of Lemma 4.3 via Proposition 4.2, allowing us to remove a certain boundedness assumption imposed in [10], and from Constantin Zălinescu, whose various deep insights led us, in particular, to simplified and more elegant proofs of Theorems 3.5 and 4.6 in comparison with the original proofs given in [10]. We are also gratefully indebted to two anonymous referees and the Associate Editor Jiří Outrata for their valuable suggestions and remarks that allowed us to essentially improve the original presentation.

#### REFERENCES

- [1] E. J. ANDERSON AND P. NASH, *Linear Programming in Infinite-Dimensional Spaces*, Wiley, Chichester, UK, 1987.
- [2] G. BEER, *Topologies on Closed and Closed Convex Sets*, Kluwer, Dordrecht, The Netherlands, 1993.
- [3] J. M. BORWEIN AND Q. J. ZHU, *Techniques of Variational Analysis*, Springer, New York, 2005.
- [4] R. I. BOŢ, S. M. GRAD, AND G. WANKA, *Generalized Moreau-Rockafellar results for composed convex functions*, *Optimization*, 58 (2009), pp. 917–933.
- [5] B. BROSOWSKI, *Parametric Semi-Infinite Optimization*, Peter Lang, Frankfurt, 1982.
- [6] M. J. CÁNOVAS, A. L. DONTCHEV, M. A. LÓPEZ, AND J. PARRA, *Metric regularity of semi-infinite constraint systems*, *Math. Program.*, 104 (2005), pp. 329–346.
- [7] M. J. CÁNOVAS, F. J. GÓMEZ-SENENT, AND J. PARRA, *Regularity modulus of arbitrarily perturbed linear inequality systems*, *J. Math. Anal. Appl.*, 343 (2008), pp. 315–327.
- [8] M. J. CÁNOVAS, A. HANTOUTE, M. A. LÓPEZ, AND J. PARRA, *Lipschitz modulus in convex semi-infinite optimization via DC functions*, *ESAIM Control Optim. Calc. Var.*, 15 (2009), pp. 763–781.
- [9] M. J. CÁNOVAS, D. KLATTE, M. A. LÓPEZ, AND J. PARRA, *Metric regularity in convex semi-infinite optimization under canonical perturbations*, *SIAM J. Optim.*, 18 (2007), pp. 717–732.
- [10] M. J. CÁNOVAS, M. A. LÓPEZ, B. S. MORDUKHOVICH, AND J. PARRA, *Robust Stability and Optimality Conditions for Parametric Infinite and Semi-Infinite Programs*, Research report 12, Department of Mathematics, Wayne State University, Detroit, MI, 2008.
- [11] M. J. CÁNOVAS, M. A. LÓPEZ, B. S. MORDUKHOVICH, AND J. PARRA, *Variational analysis in semi-infinite and infinite programming, II: Necessary optimality conditions*, *SIAM J. Optim.*, submitted.

- [12] M. J. CÁNOVAS, M. A. LOPEZ, J. PARRA, AND M. I. TODOROV, *Stability and well-posedness in linear semi-infinite programming*, SIAM J. Optim., 10 (1999), pp. 82–98.
- [13] N. DINH, M. A. GOBERNA, AND M. A. LÓPEZ, *From linear to convex systems: Consistency, Farkas' lemma and applications*, J. Convex Anal., 13 (2006), pp. 279–290.
- [14] N. DINH, M. A. GOBERNA, AND M. A. LÓPEZ, *On the stability of the feasible set in optimization problems*, submitted (2008).
- [15] N. DINH, B. S. MORDUKHOVICH, AND T. T. A. NGHIA, *Subdifferentials of value functions and optimality conditions for DC and bilevel infinite and semi-infinite programs*, Math. Program., to appear.
- [16] N. DUNFORD AND J. T. SCHWARTZ, *Linear Operators. Part I: General Theory*, Wiley, New York, 1988.
- [17] M. FABIAN, P. HABALA, P. HÁJEK, V. MONTESINOS-SANTALUCÍA, J. PELANT, AND V. ZIZLER, *Functional Analysis and Infinite-Dimensional Geometry*, Springer, New York, 2001.
- [18] M. A. GOBERNA, V. JEYAKUMAR, AND M. A. LÓPEZ, *Necessary and sufficient constraint qualifications for solvability of systems of infinite convex inequalities*, Nonlinear Anal., 68 (2008), pp. 1184–1194.
- [19] M. A. GOBERNA AND M. A. LÓPEZ, *Linear Semi-Infinite Optimization*, Wiley, Chichester, UK, 1998.
- [20] A. D. IOFFE, *Metric regularity and subdifferential calculus*, Russian Math. Surveys, 55 (2000), pp. 501–558.
- [21] H. T. JONGEN, J.-J. RÜCKMANN, AND G.-W. WEBER, *One-parametric semi-infinite optimization: On the stability of the feasible set*, SIAM J. Optim., 4 (1994), pp. 637–648.
- [22] D. KLATTE AND R. HENRION, *Regularity and stability in nonlinear semi-infinite optimization*, in Semi-Infinite Programming, R. Reemtsen and J.-J. Rückmann, eds., Kluwer, Dordrecht, The Netherlands, 1998, pp. 69–102.
- [23] D. KLATTE AND B. KUMMER, *Nonsmooth Equations in Optimization. Regularity, Calculus, Methods and Applications*, Kluwer, Dordrecht, The Netherlands, 2002.
- [24] C. LI AND K. F. NG, *On constraint qualification for an infinite system of convex inequalities in a Banach space*, SIAM J. Optim., 15 (2005), pp. 488–512.
- [25] C. LI, K. F. NG, AND T. K. PONG, *Constraint qualifications for convex inequality systems with applications in constrained optimization*, SIAM J. Optim., 19 (2008), pp. 163–187.
- [26] B. S. MORDUKHOVICH, *Maximum principle in problems of time optimal control with nonsmooth constraints*, J. Appl. Math. Mech., 40 (1976), pp. 960–969.
- [27] B. S. MORDUKHOVICH, *Metric approximations and necessary optimality conditions for general classes of extremal problems*, Soviet Math. Dokl., 22 (1980), pp. 526–530.
- [28] B. S. MORDUKHOVICH, *Complete characterizations of openness, metric regularity, and Lipschitzian properties of multifunctions*, Trans. Amer. Math. Soc., 340 (1993), pp. 1–35.
- [29] B. S. MORDUKHOVICH, *Coderivatives of set-valued mappings: Calculus and applications*, Nonlinear Anal., 30 (1997), pp. 3059–3070.
- [30] B. S. MORDUKHOVICH, *Necessary conditions in nonsmooth minimization via lower and upper subgradients*, Set-Valued Anal., 12 (2004), pp. 163–193.
- [31] B. S. MORDUKHOVICH, *Variational Analysis and Generalized Differentiation, I: Basic Theory*, Springer, Berlin, 2006.
- [32] B. S. MORDUKHOVICH, *Variational Analysis and Generalized Differentiation, II: Applications*, Springer, Berlin, 2006.
- [33] J. V. OUTRATA, *Mathematical programs with equilibrium constraints: Theory and numerical methods*, in Nonsmooth Mechanics of Solids, J. Haslinger and G. E. Stavroulakis, eds., Springer, New York, 2006, pp. 221–274.
- [34] R. R. PHELPS, *Convex Functions, Monotone Operators, and Differentiability*, 2nd ed., Springer, Berlin, 1993.
- [35] R. T. ROCKAFELLAR AND R. J.-B. WETS, *Variational Analysis*, Springer, Berlin, 1998.
- [36] W. SCHIROTZEK, *Nonsmooth Analysis*, Springer, Berlin, 2007.
- [37] O. STEIN, *Bilevel Strategies in Semi-Infinite Programming*, Kluwer, Boston, 2003.
- [38] C. ZĂLINESCU, *Solvability results for sublinear functions and operators*, Z. Oper. Res. Ser. A-B, 31 (1987), pp. 79–101.
- [39] C. ZĂLINESCU, *Convex Analysis in General Vector Spaces*, World Scientific, Singapore, 2002.