Revision of Genus *Enclisis* Townes and description of two new species (Hymenoptera: Ichneumonidae)

Bordera, S.¹, Kolarov, J.², Mazón, M.¹

Two new species of Enclisis Townes, 1970, from Spain and Balkan Region are described and illustrated: *E. castellana* sp.n. and *E. balcanica* sp.n. Females of the new species are distinguished between, and from the closest species *E. ornaticeps*, by microsculpture of first lateral area of propodeum and mesopleurum, head features and body colour. Male of *E. balcanica* is distinguished from the closest species *E. infernator* and *E. schwarzi* by its head, less transversal, and for a longer second metasomal tergite. Male of *E. castellana* is separated from *E. ruficeps* by its microsculpture on mesopleurum and mesonotum, always on shiny background, and for the position of subdiscoideus, which reach the postnervulus in the middle or above the centre. Furthermore, data on phenology of *E. castellana* and a new illustrated key for the *Enclisis* species, excluding *E. nigricoxis* and *E. ruficoxis*, is provided. The species *E. alpicola*, *E. ornaticeps* and *E. macilenta* are recorded for the first time for Bulgarian fauna. *E. infernator* is first record for Bulgaria and Spain.

This research was supported partially by the Projects 040/2002 from Ministerio de Medio Ambiente of Spanish Government and GV06/271 from Conselleria d'Empresa, Universitat i Ciència of Generalitat Valenciana.

Published in the book of abstracts of the X European Workshop on Insect Parasitoids, Erice (Sicily), Italy, 17-21 september, 2007, p. 73.

¹ Instituto de Investigación de Biodiversidad CIBIO, Alicante University, Ap. Correos 99, 03080 (Spain), s.bordera@ua.es,

² Faculty of Pedagogie, Plovdiv University, 24 Tsar Assen Str., 4000 Plovdiv (Bulgaria), j.kolarov@pu.acad.bg,

¹ Instituto de Investigación de Biodiversidad CIBIO, Alicante University, Ap. Correos 99, 03080 Alicante (Spain), mazon@ua.es.