2 Responses of two Mediterranean seagrasses to experimental 3 changes in salinity

4 Yolanda Fernández-Torquemada ·
5 José Luis Sánchez-Lizaso

6 Received: 27 October 2010 / Revised: 7 February 2011 / Accepted: 12 February 2011
7 © Springer Science+Business Media B.V. 2011

8 **Abstract** The aim of this study is to examine the
9 effects of variations in salinity levels on growth and
10 survival of two fast-growing Mediterranean seagrass-
11 es, *Cymodocea nodosa* and *Zostera noltii*. We also
12 tested the capacity of *C. nodosa* to acclimate to a
13 gradual increase in salinity and to discover how it
14 responds to a sharp rise in salinity in combination
15 with other factors, such as increases in temperature,
16 seasonality and different plant-population origins.
17 Several short-term (10 days) experiments were con-
18 ducted under controlled conditions. For each exper-
19 iment, ten marked shoots were placed in 5-l aquaria,
20 where they were exposed to different salinity treat-
21 ments (ranging from 2 to 72 psu). Growth and
22 survival of both species were significantly affected
23 by salinity. A significant effect between salinity and
24 temperature on the shoot growth rate of *C. nodosa*
25 was also detected, but not on shoot mortality. When
26 *C. nodosa* plants were acclimated by gradually
27 increasing the salinity level, it was observed that
28 acclimatisation improved tolerance to salinity
29 changes. A different response to salinity variations,
30 depending on the origin of the plants or the season of

the year, was also detected. These results indicated
that *Z. noltii* plants tolerate conditions of hyposalinity
better than *C. nodosa*, and that the tolerance range of
C. nodosa may change depending on the temperature,
the season or the population.

Keywords Salinity effects · Salinity tolerance ·
Desalination impact · *Cymodocea nodosa* ·
Zostera noltii

Introduction

Seagrasses have evolved from continental angio-
sperms that have returned and adapted to life in
completely submerged saline environments (den
Hartog, 1970), which they are able to tolerate as a
result of various biochemical, physiological and
morphological adaptations (Jagels, 1973; Tyerman,
1989; Arai et al., 1991; Pak et al., 1995; Fukuhara
et al., 1996; Fernández et al., 1999; Touchette, 2007).
These mechanisms have allowed seagrasses to occur
naturally in various aquatic environments with
different salinity values, including brackish and
oceanic waters (Walker, 1985; Adams & Bate,
1994; Tomasko & Hall, 1999), and from almost
stable to a more fluctuating environment.

To date, studies examining salinity influence or
tolerance in seagrasses have focused mainly on
estuarine species, which can be exposed to a wide
range of salinity levels, due to seasonal and natural

A1 Handling editor: Pierluigi Viaroli

A2 Y. Fernández-Torquemada (✉) · J. L. Sánchez-Lizaso
A3 Department of Marine Sciences and Applied Biology,
A4 University of Alicante, P.O. Box 99, 03080 Alicante,
A5 Spain
A6 e-mail: Yolanda.Fernandez@ua.es

variations occurring in their habitat (Wortmann et al., 1997; Benjamin et al., 1999; Kamermans et al., 1999). Results from these studies have shown that alterations to salinity levels can affect plant metabolism, growth, reproduction, survival and distribution (McMillan & Moseley, 1967; Zieman, 1975; Walker & McComb, 1990; Montague & Ley, 1993; Hillman et al., 1995; Ramage & Schiel, 1998; Vermaat et al., 2000; Chesnes & Montague, 2001). However, in marine environments, salinity is considered an approximately constant factor, so any alteration to salinity levels may lead to significant disturbances to open water seagrasses that are more stenohaline species. Salinity values in the Mediterranean Sea are not naturally subjected to large and rapid fluctuations. However, some Mediterranean coastal areas are currently exposed to anthropogenic alterations in salinity levels caused by the discharge of brine effluents from desalination plants (Fernández-Torquemada et al., 2005a, 2009; Gacia et al., 2007). These brine discharges have high salinity levels (40–70 psu), and sometimes a relatively high temperature and low pH (Lattemann & Höpner, 2003), so these and other environmental conditions occurring at the same time may have an interaction with increased salinity levels.

Previous studies have shown that the endemic species *Posidonia oceanica* is quite sensitive to increases in salinity levels (Fernández-Torquemada & Sánchez-Lizaso, 2005; Gacia et al., 2007; Sánchez-Lizaso et al., 2008), but little information exists with regard to the response and tolerance of other Mediterranean seagrasses, such as *Cymodocea nodosa* and *Zostera noltii*. *C. nodosa* and *Z. noltii* are relatively small and fast-growing seagrasses with similar behaviour and ecology, but different geographical distribution. Whereas *C. nodosa* is a common species in the Mediterranean and the eastern Atlantic, from south Portugal to Senegal and around the Canary Islands (den Hartog, 1970), *Z. noltii* is widespread along the European Atlantic coastline from Sweden to Mauritania, but is not common in the Mediterranean Sea. *C. nodosa* may also show a wider depth distribution (ranging from the intertidal to 33–35 m deep) than *Z. noltii*, which has a narrower vertical distribution (Drew, 1978; Vermaat et al., 1993; Reyes et al., 1995). On the other hand, both species have a high tolerance to a variety of environmental conditions, such as irradiance changes, seasonal

temperature fluctuations and different nutrient concentrations (Phillips & Meñez, 1988; Marbà et al., 1996), and can become established on a wide range of substrata, from sandy to muddy pristine as well as degraded coastal sites (Peduzzi & Vuković, 1990; Pavón-Salas et al., 2000; Charpentier et al., 2005). As the habitat preferences and environmental limits of *C. nodosa* and *Z. noltii* are very broad, these species can be expected to have higher salinity tolerance levels than *P. oceanica*. Throughout their distribution, both seagrasses can be found forming mixed meadows in areas where salinity values differ from marine conditions, such as estuaries and coastal lagoons (den Hartog, 1970; Mazzella et al., 1993; Vermaat et al., 2000; Greve & Binzer, 2004). Based on their field distribution, *C. nodosa* and *Z. noltii* have been classified as euryhaline species (den Hartog, 1970), although it is expected that individuals of both species that currently occur in coastal Mediterranean waters will have a different salinity tolerance to those from other populations occurring in habitats with greater fluctuations in salinity.

On the other hand, salinity tolerance in both species cannot be easily inferred from these field data, as the salinity range of normal growth and development for a species is usually narrower than its real tolerance limits (Kinne, 1964), and there can be multiple stressors in the field affecting the behaviour and distribution of a species. As a result, controlled experimental studies are needed to gather significant information about the environmental tolerance of a species. There are some experimental studies that refer to the effects of salinity variations on these seagrasses (Caye & Meinesz, 1986; Hootsmans et al., 1987; Loques et al., 1990; Caye et al., 1992; Vermaat et al., 2000; Pagès et al., 2010). Most of these studies have focused on the role that reduced salinity levels play in seed germination. Caye & Meinesz (1986) demonstrated that they could induce *C. nodosa* seed germination at any time of the year by reducing the salinity and using a temperature between 20 and 25°C. These same authors observed that seeds generally do not germinate at a salinity level of 38 psu, and the few that germinated did so very slowly (over months). Furthermore, seedlings that had germinated at lower salinity levels (15–30 psu) were the only ones that continued to develop. Other studies have demonstrated that low salinity levels also stimulate *Z. noltii* seed germination (Hootsmans

et al., 1987; Loques et al., 1990). Vermaat et al. (2000) carried out an experiment with adult *Z. noltii* plants from two separate populations that were exposed to 15 and 35 psu and the day-length regimes of two seasons, and observed that both populations suffered high mortality at 35 psu. In a recent paper, Pagès et al., (2010) estimated that *C. nodosa* was tolerant to moderate salinity increases (44 psu), while apparently only sub-lethal effects appeared at 54 psu, but they only studied four salinity treatments.

The first aim of the present study is to redress the lack of existing information on the tolerance of *Cymodocea nodosa* and *Zostera noltii* to salinity changes and to establish their possible range of salinity tolerance. Several experiments of a short duration (10 days) were conducted under controlled conditions to estimate shoot growth rate and survival in both species when exposed to different levels of salinity. A more thorough study of *C. nodosa* was also carried out to prove the effect of other factors associated with the discharge of desalination brine on this species. We thus tested the capacity of *C. nodosa* to acclimate to gradual increases in salinity and its response to sharp rises in salinity combined with other factors or conditions, such as temperature increases, seasonality and different plant populations.

Materials and methods

Plant material

Cymodocea nodosa and *Zostera noltii* shoots were carefully collected by scuba diving from a mixed shallow meadow (−2 m) at Almadra Beach (Alicante, SE Spain). Plants of a similar size and vitality were transported to the laboratory in a cooler containing ambient seawater from the sampling site (average salinity of 37.5 psu). Once transferred to the laboratory, the shoots were marked using a modified Zieman method (1974) and placed in plastic 5-l aquaria with sediment at different treatments for 10 days.

Experimental design

Six sets of experiments were conducted in aquaria under environmentally controlled conditions, with a daylight regime of 12 h of light and 12 h of darkness,

and an average daytime underwater photon flux of 30 $\mu\text{mol quanta m}^{-2} \text{s}^{-1}$ (measured with a LiCor 193 SA spherical quantum sensor), that is approximately the environmental light intensity observed in the plants sampling site (Almadra Beach, Alicante; 38°22'N, 0°26'W). For each experiment, three aerated aquaria (replicates) were assigned per treatment. With the exception of experiment VI, ten plants were placed in each aquarium with no acclimation period, in order to simulate sudden increases in salinity. In all the experiments carried out, temperature and salinity were measured daily using a conductimeter (model 1230; Thermo Orion), and salinity values were adjusted when necessary. Control-treatment salinity was the level measured in the area where the plants were collected (with a range of 36.8–38.1 psu, and an average of 37.5 psu). Increased salinity treatments were similar to hypersalinity values associated with a desalination brine discharge (Fernández-Torquemada et al., 2005a, 2009) and were prepared by adding natural salt produced by Santa Pola saltworks from seawater concentrate to ambient coastal seawater, while lower salinities were obtained by diluting seawater with freshwater. Once the aquaria were filled with treatment water, they were placed in a larger water-filled container, to keep temperatures as constant as possible during each experiment.

Leaf growth was measured, by using a hypodermic needle for *C. nodosa* or by using a water-insoluble marking pen for *Z. noltii*, as a daily elongation rate ($\text{cm}^2 \text{shoot}^{-1} \text{day}^{-1}$), and mortality as a percentage of dead shoots per aquaria at the end of each experiment. Shoots were considered dead when leaves died back and rhizomes and roots were totally degraded. In order to compare the results from selected experiments, some leaf growth rates were reported as a percentage relative to the growth obtained for the control plants.

The following section details the objectives, experimental design and methodology of the different experiments carried out.

Experiment I: salinity tolerance of *Zostera noltii*

To evaluate the effect of salinity on shoot growth rate and survival of *Z. noltii*, three consecutive and different tests of a short duration (10 days) were carried out during the summer period (July and August, temperature ≈ 25 – 27°C), working with a

247	wide range of salinities (21 different treatments	<i>Experiment V: differences between Cymodocea</i>	288
248	between 2 and 72 psu, each one replicated in three	<i>nodosa</i> populations	289
249	independent aquaria).		
250	<i>Experiment II: salinity tolerance of Cymodocea</i>		
251	<i>nodosa</i>		
252	This study was designed to evaluate the effect of	This experiment was performed during 10 days in	290
253	different salinity levels on shoot growth rate and	August (temperature $\approx 27^{\circ}\text{C}$) to determine whether	291
254	survival of <i>C. nodosa</i> . For this experiment, five short-	the response of this species to changes in salinity can	292
255	term (10 days) assays were carried out during spring	vary between individual plants from two populations	293
256	and summer months (April–July, tempera-	adapted to different conditions. The survival rates and	294
257	ture $\approx 22\text{--}25^{\circ}\text{C}$), with 27 different salinity levels	shoot growth rate among two <i>C. nodosa</i> populations	295
258	ranging from 2 to 72 psu (each salinity treatment	were compared: one from the western Mediterranean	296
259	replicated in three aquaria).	Sea (Alicante; $38^{\circ}22'\text{N}$, $0^{\circ}26'\text{W}$) and another from a	297
		near coastal lagoon located in southeast Spain (Mar	298
260	<i>Experiment III: seasonal response of C. nodosa</i>	Menor, Murcia; $37^{\circ}48'\text{N}$, $0^{\circ}46'\text{W}$) with a higher	299
261	<i>to increases in salinity</i>	salinity level (42–47 psu). The salinity levels applied	300
		were 37 psu (the ambient level at the Mediterranean	301
262	The aim of this experiment was to detect a possible	sampling site), 44 psu (the level at the lagoon when	302
263	seasonal variation of the response by <i>C. nodosa</i> to	and where the plants were collected), 47 and 50 psu.	303
264	hypersalinity (control, 42, 47 and 52 psu). Increased		
265	salinity experiments (10 days) were carried out at two	<i>Experiment VI: acclimation versus acute increases</i>	304
266	different times of the year. Winter time (20–30	<i>in C. nodosa</i>	305
267	January, temperature $\approx 15^{\circ}\text{C}$) was selected because		
268	of the reduced growth rate of <i>C. nodosa</i> during this	The aim of this experiment, also performed during	306
269	season, compared with summer (5–15 July, tempera-	August (temperature $\approx 26^{\circ}\text{C}$), was to determine	307
270	ture $\approx 25^{\circ}\text{C}$), when, due to the accumulation of	whether the acclimation of <i>Cymodocea nodosa</i> to	308
271	sucrose and starch and increases in irradiance and	gradual salinity variations allows for a greater	309
272	temperature, the species shows a higher rate of	tolerance to osmotic stress than an instantaneous	310
273	development.	transfer, as occurs with other seagrass species (Ralph,	311
		1998). In some treatments, plants were acclimated in	312
274	<i>Experiment IV: salinity and temperature interaction</i>	their respective aquaria to three different salinity	313
275	<i>effects on C. nodosa</i>	levels (42, 47, and 52 psu) in a stepwise manner	314
		(2.5 psu per day) up from the habitat salinity of 37	315
276	This experiment of 10 days was carried out to	psu, reaching the maximum salinity after 2 (42 psu),	316
277	examine the synergistic effects of increases in	4 (47 psu) and 6 days (52 psu). In other treatments,	317
278	temperature and salinity. Control aquaria were	plants were subjected to sudden increases in salinity	318
279	maintained at 20°C seawater temperature during that	(also 42, 47 and 52 psu), as performed in previous	319
280	period (spring, March), while, for the other aquaria, a	experiments. Finally, all plants were subjected to	320
281	commercial heater was used with a thermostat to	these salinity treatments during 10 days.	321
282	increase the temperature to 25°C , representing sum-		
283	mer temperatures in their natural environment.	Statistical analyses	322
284	Within each temperature regime (20 and 25°C), four		
285	salinity treatments were tested to represent an	One-way ANOVA was used in experiments I and II	323
286	increase in this variable: 37 (control), 43, 48 and 53	to test for differences in mortality among the different	324
287	psu.	salinity treatments used (including one as control).	325
		Nested ANOVA was used in the same experiments to	326
		test differences in shoot growth rate, with salinity	327
		considered as a fixed factor replicated in three aquaria	328
		(random factor and nested within salinity) and	329
		with ten marked shoots (replicates). In the rest of	330
		the experiments, an orthogonal factor was added	331

(temperature, original population, seasonality or acclimation) to determine possible interactions between salinity and these factors. Homogeneity of variance had been previously verified using the Cochran test (Underwood, 1997). The data were log-transformed if normality was not verified.

When analysis of variance identified a significant difference for any factor, the multiple comparison post-hoc test SNK (Student–Newman–Keuls) was applied to determine specific treatment differences. All calculations were performed using the GMAV.5 program (University of Sydney; Underwood & Chapman, 1997), with a minimum significance level established at $P < 0.05$.

Results

The growth and survival of *Zostera noltii* were both significantly affected by water salinity (Fig. 1). Shoot growth rate was considerably affected at higher salinities, but not at reduced salinities (Fig. 1). SNK test results revealed that elongation rates were similar

and maximal at 2–41 psu, decreasing significantly from 42 to 52 psu ($P < 0.05$). No growth was observed at 57 psu and higher. Hypersalinity also affected the mortality rate of this species (Fig. 1). *Z. noltii* plants sustained considerable mortality at salinity levels above 43 psu ($>16.7\%$), reaching 50% at ~50 psu and 100% mortality at 57 psu and higher.

Salinity also had a strong effect on leaf elongation rate and mortality in *Cymodocea nodosa* (Fig. 2). The relationship between shoot growth rate and salinity was bell-shaped (Fig. 2), with a distinct peak at 30–39 psu, and a significant reduction at salinity levels higher than 41 psu or lower than 16 psu. Mortality showed an opposite pattern (Fig. 2), with a minimum value at the control salinity level. Shoot mortality remained lower than 50% at salinity levels of less than 50 psu, but increased sharply above this level. All plants died when exposed to freshwater (0 psu) and to the highest salinity concentrations (≥ 57 psu).

The *C. nodosa* response to salinity increases during the two seasons studied (winter and summer) showed a significantly different behaviour (Fig. 3). The life

Fig. 1 *Zostera noltii* leaf growth (relative to the growth of the control plants) and mortality percentage at different salinity levels (bars represent standard errors)

Fig. 2 *Cymodocea nodosa* leaf growth (relative to the growth of the control plants) and mortality percentage at different salinity levels (bars represent standard errors)

cycle of this species is strongly marked by seasonality (Pérez & Romero, 1992), with the result that in summer, the plants presented a significantly ($P < 0.001$) higher leaf growth ($0.1963 \text{ cm}^2 \text{ day}^{-1} \text{ shoot}^{-1}$) than in winter ($0.0352 \text{ cm}^2 \text{ day}^{-1} \text{ shoot}^{-1}$). To compare the data obtained from the two seasons, the results were expressed as a percentage relative to the growth obtained at the control treatment (Fig. 3). Here, two-way ANOVA showed a significant interaction between salinity and the season of the year in which the experiment was carried out (Table 1). Subsequent SNK post-hoc tests indicated that, in summer, *C. nodosa* shoot growth rate was significantly higher at the control salinity level than at the other salinity treatment levels, whereas in winter, higher salinity (51 psu) is the only level that differs from the rest (Fig. 3). It was also observed that the effects of increased salinity on *C. nodosa* mortality was more evident in summer than in winter (Fig. 3), although no significant interaction was found between seasonality and salinity (Table 1). Shoot mortality was generally

lower ($\leq 20\%$) in winter, with higher mortality rates observed in summer (53.3% at 51 psu).

ANOVA revealed a significant effect of the interaction between temperature and increased salinity on the leaf growth of this species ($P < 0.01$). Plants subjected to salinity levels between 37 and 48 psu showed greater shoot growth rates at the highest temperature (25°C), but when salinity was increased to 53 psu, the increase in temperature did not affect leaf growth (Fig. 4). Although no interaction was found for mortality data (Table 1), mortalities associated with increases in salinity were lower at 25°C (Fig. 4). In fact, in this experiment, only salinity had a significant effect on the survival of *C. nodosa* shoots. Among the four salinity treatments, the highest mortality rates occurred at 53 psu (63.3–86.7%), followed by 48 psu (20–40%), with significantly lower values than 37 and 43 psu (3.3–6.7%).

Analysis of variance also showed significant differences in the response to salinity among the populations studied (Table 1). Individuals from the Mar Menor presented greater shoot growth and survival rates than those from the Mediterranean Sea (Fig. 5). Similar shoot growth and survival was observed in plants subjected from 37 to 47 psu, with significantly lower values reported at 50 psu. On the other hand, although no significant interaction was observed, plants from the Mediterranean Sea exposed to 50 psu clearly showed lower growth and survival rates than those from the Mar Menor at the same salinity treatment (Fig. 5).

On the other hand, *C. nodosa* shoots acclimated to gradual increases in salinity (2.5 psu per day) showed higher shoot growth rates at the four salinity treatments studied than those not subjected to acclimation (Fig. 6). No dead shoots were found at the control salinity level. However, at high salinity levels, the mortality rate was higher for plants exposed to abrupt salinity changes, although ANOVA did not detect any significant interaction.

Fig. 3 Leaf growth per shoot, relative to the growth of the control plants, and mortality percentage of *C. nodosa* plants at different salinity levels during summer and winter (bars represent standard errors)

Discussion

Effects of salinity on shoot growth rate and survival of *C. nodosa* and *Z. noltii*

Under these laboratory conditions, *Zostera noltii* was found to be more tolerant of conditions of hypersalinity than *Cymodocea nodosa*, but both species were

Table 1 Summary of the two-way ANOVAs testing the effects of different salinity treatments combined with other factors or conditions on *C. nodosa* shoot growth rate and shoot mortality

Experiment	Source of variation	Leaf elongation			Mortality		
		df	MS	F	df	MS	F
Exp. III	Salinity	7	0.0329	5.08***	7	723.81	3.66**
	Season of year	1	0.7594	117.47***	1	3,333.33	16.84**
	Salinity × season	7	0.0179	2.77*	7	280.95	1.42 ^{ns}
	Aquaria (salinity × season)	32	0.0065	2.31***	32	197.91	
	Residual	432	0.0028				
Exp. IV	Salinity	3	0.0404	94.32***	3	6,700.00	41.23***
	Temperature	1	0.0318	74.24***	1	600.00	3.69 ^{ns}
	Salinity × temperature	3	0.0031	7.16**	3	277.78	1.71 ^{ns}
	Aquaria (salinity × temperature)	16	0.0004	0.75 ^{ns}	16	162.50	
	Residual	216	0.0006				
Exp. V	Salinity	3	0.0003	5.73**	3	1,181.94	5.56**
	Population	1	0.0010	21.75***	1	2,204.17	10.37**
	Salinity × population	3	0.0001	1.32 ^{ns}	3	604.17	2.84 ^{ns}
	Aquaria (salinity × population)	16	0.0000	1.41 ^{ns}	16	212.50	
	Residual	216	0.0000				
Exp. VI	Salinity	3	0.0602	7.40**	3	1,693.06	8.13**
	Acclimation	1	0.0490	6.02*	1	2,204.17	10.58**
	Salinity × acclimation	3	0.0060	0.73 ^{ns}	3	293.06	1.41 ^{ns}
	Aquaria (salinity × acclimation)	16	0.0081	2.46**	16	208.33	
	Residual	216	0.0033				

ns non-significant, * $P < 0.05$, ** $P < 0.01$, *** $P < 0.005$

sensitive to increases in salinity. These experiments showed significant decreases in shoot growth rates in both seagrasses at salinity levels higher than 41 psu. A decrease in leaf growth was also observed at salinity levels lower than 15 psu for *Cymodocea nodosa*, whereas *Z. noltii* plants were found to be highly tolerant to salinity levels as low as 2 psu, which was the lowest salinity treatment tested for the 10-day period. This tolerance to low levels of salinity has also been found for other seagrasses, such as *Amphibolis antarctica* and *Posidonia sinuosa*, which can be highly tolerant to short-term reductions in salinity, even as low as 0–1 psu (Westphalen et al., 2005). The relationship between leaf growth and salinity was well described by a bell-shaped curve for *C. nodosa*, and by a sigmoid curve for *Z. noltii*, with maximum values recorded around ambient salinity levels. Similar responses, but with different relationships between salinity and plant growth, have been observed in several seagrasses (McMillan & Moseley, 1967; Walker, 1985; Walker & McComb,

1990; Kamermans et al., 1999) probably due to adverse effects on the meristematic tissues, interferences with carbon metabolism or negative alterations in the photosynthetic and respiratory rates (Ogata & Matsui, 1965; Biebl & McRoy, 1971; Kraemer et al., 1999). Mortality values for *C. nodosa* were significantly lower for salinity levels between 17 and 48 psu, with all plants dying at salinity levels of more than 56 psu. For *Z. noltii*, mortality was significantly lower at salinity levels below 47 psu, with 100% mortality at salinity levels above 56 psu. Previous studies have confirmed that salinity variations may affect seagrass survival under experimental conditions (Biebl & McRoy, 1971; Pinnerup, 1980; Vermaat et al., 2000; Fernández-Torquemada & Sánchez-Lizaso, 2005; Fernández-Torquemada et al., 2005b), as well as in their natural habitats (Robblee et al., 1991; Wortmann et al., 1997; Kamermans et al., 1999; van Katwijk et al., 1999). Water salinities outside the tolerance range of a species may alter its metabolism and therefore cause direct death of the plant at very

Fig. 4 *Cymodocea nodosa* shoot growth rate (cm² day⁻¹ shoot⁻¹) and shoot mortality (%) at different salinity levels and temperatures (bars represent standard errors)

Fig. 5 Shoot growth rate (cm² day⁻¹ shoot⁻¹) and mortality percentage of *C. nodosa* plants from two different populations exposed to different salinity levels (bars represent standard errors)

extreme levels (McMillan & Moseley, 1967; Adams & Bates, 1994), or death in the long term if causing starch-reserve depletion (Biebl & McRoy, 1971; Kerr & Strother, 1985).

These results are also in accordance with the field distribution of both seagrasses. *Z. noltii* is known to colonise estuary and delta areas with a higher influence of freshwater inputs than *C. nodosa* (Vermaat et al., 2000; Greve & Binzer, 2004). Our data are also coherent with the known colonisation of these species in the Mar Menor, a coastal lagoon located in the southeast of Spain. Until the end of the nineteenth century, this lagoon contained very high salinity levels (60–70 psu) that prevented it being colonised by macrophytes. However, a subsequent connection was established between the lagoon and the Mediterranean Sea, causing salinity to drop to the current levels (42–47 psu), allowing for the existing seagrass meadows to become established (Pérez Ruzafa et al., 1987).

It should be emphasised that all these experiments were carried out under controlled aquarium conditions and in short periods of time (10 days), so the response

observed here could differ to the one observed in the natural environment. However, Walker & McComb (1990) compared the effect of salinity variations on the growth of a tropical seagrass, both in situ and in aquaria, and found that its tolerance was similar in both cases. In the present study, aquaria experiments were also performed under low light levels (30 $\mu\text{mol quanta m}^{-2} \text{s}^{-1}$), but higher environmental light levels could imply a lower mortality or higher growth of plants under stressing salinity treatments, because those plants could be subjected to an additional co-stressor, the low light level. Moreover, we only worked with growth and survival of those species, and no information about photosynthesis or other descriptors is given. On the other hand, this study focussed solely on the effects of increases in salinity on the growth and survival of adult shoots, and the sensitivity to environmental variations may differ considerably between seedlings and adult plants, with young stages typically being more sensitive to stressors than adult ones. Future studies should therefore investigate the effects of increases in salinity, together with

Fig. 6 Shoot growth rate ($\text{cm}^2 \text{day}^{-1} \text{shoot}^{-1}$) and shoot mortality (%) of *C. nodosa* plants acclimated to salinity increases (2.5 psu day^{-1}) versus plants subjected to sudden salinity increases (bars represent standard errors)

variations in other conditions, on seedling development and seed germination of these species.

Nevertheless, if we compare the results obtained here for *C. nodosa* and *Z. noltii* with those obtained for *P. oceanica* (Fernández-Torquemada & Sánchez-Lizaso, 2005), it can be observed that these species were able to tolerate a wider range of salinity levels than *P. oceanica*. This is coherent with the fact that *C. nodosa* and *Z. noltii* are smaller and much more plastic species, with a higher capacity to adapt to environmental changes and recover from slight disturbances (Pérez et al., 1994; Kraemer & Mazzella, 1999).

Interactions of salinity with other factors on *Cymodocea nodosa*

Previous experiments have demonstrated the sensitivity of *C. nodosa* to salinity variations in the short term (10 days), but tolerance to salinity may also depend on other environmental or intra-specific

factors. Other studies have demonstrated that osmotic stress could modify the sensitivity of some seagrasses to other environmental conditions, such as temperature or pH increases, light levels or eutrophication (Biebl & McRoy, 1971; Ralph 1999; van Katwijk et al., 1999; Vermaat et al., 2000).

Mediterranean seawater temperatures vary considerably between seasons, but rapid temperature increases may be linked to brine discharges from certain desalination plants. For example, brine discharges from multi-stage flash desalination plants can cause increases of 15°C in the seawater temperature in the receiving environment (Lattemann & Höpner, 2003). On the other hand, brine discharges from reverse osmosis desalination plants do not imply temperature increases, though it has been observed that, in some of these facilities and during certain periods of the year, the temperature of the effluent can produce a slight increase in that of the receiving environment (Fernández-Torquemada et al., 2005a).

Temperature is considered an important abiotic factor that controls the production, growth and survival of a species (Drew, 1979), and it has been demonstrated that its influence on the metabolism of marine plants modifies their capacity to adapt to changes in other environmental factors, such as illumination or salinity (Zieman, 1975). Plant growth is most affected by light and temperature (Philippart, 1995; Marbà et al., 1996), and when these parameters are close to a species' optimum values, this species will tolerate a broader range of salinity levels (Kirst, 1989). It is known that, in several species of algae, such as *Macrocystis integrifolia*, *Alaria esculenta* and *Cladophora rupestris*, tolerance to saline stress is reduced at extreme levels within their temperature tolerance range (Druehl, 1981; Dring, 1992; Thomas et al., 1988). In seagrasses, Biebl & McRoy (1971) found an increase in plasmatic heat resistance with increasing levels of salinity for subtidal forms of *Zostera marina*, and Vermaat et al. (2000) observed increased mortality in *Z. noltii* when the temperature reached 20°C at higher experimental levels of salinity. Thorhaug & Marcus (1981) indicated that four seagrass species were less tolerant of low salinity levels at temperature above 29°C . The present work has also observed a significant interaction between the two factors. For salinities ranging from 37 to 48 psu, *C. nodosa* plants possessed a higher growth rate at the highest temperature (25°C), but when the

salinity was increased to 53 psu, the increase in temperature did not have a positive effect on leaf growth. This is probably because the increase in metabolic activity at high temperatures is insufficient for coping with the stress caused by high salinity levels.

Cymodocea nodosa is also affected by seasonality, showing minimum growth in winter due to lower irradiance and temperatures. In this study, it was observed that the salinity tolerance range of this species may also vary throughout the year. This could be related to maximum growth and, therefore, the higher energy and metabolic consumption that occur in summer due to the species' life cycle that makes more evident salinity effects. In winter, plants survive with smaller metabolic levels and are more able to resist salinity-related stress.

On the other hand, tolerance can differ considerably at one level of salinity depending on how such a level is reached. Other studies report that gradual increases in salinity are usually better resisted than a sudden or rapid rise (Ralph, 1998). In this study, *C. nodosa* plants exposed to slower increases of salinity (2.5 psu per day) were able to tolerate changes in salinity better than plants that had not been acclimated, showing higher rates of growth and survival.

The capacity of *C. nodosa* to improve its tolerance to slow increases in salinity can be explained by an acclimation of plants, and some structural or physiological reversible changes that occur when plants are exposed to stress. But when studying other stresses, it has been observed that some plant populations exposed to these same conditions during a longer period may be able to adapt by means of some genetic and irreversible changes (Peralta et al., 2005).

In fact, when conducting one of the experiments comparing the response of *Cymodocea nodosa* plants from the Mar Menor lagoon with plants from the Alicante (Mediterranean) near-shore coast, different behaviours to salinity variations were observed for both populations. For the Mediterranean population studied, salinity tolerance was narrower than for the Mar Menor population, as indicated by the high rates of mortality and the significant reduction in shoot growth rate at 50 psu. Plants from the Mar Menor, meanwhile, showed a lower sensitivity to increases in salinity, and similar rates of mortality and leaf growth

inside the salinity range studied. Salinity in the Mar Menor lagoon currently ranges between 42 and 47 psu, and western Mediterranean coasts are at levels of ~37 and 38 psu. So these differences in tolerance to salinity could be explained because, as occurs with other widespread species, individuals adapt to the different local conditions that occur naturally in their habitats (Doering & Chamberlain, 1998; Benjamin et al., 1999; Kamermans et al., 1999; van Katwijk et al., 1999; Vermaat et al., 2000). For example, Kamermans et al. (1999) and van Katwijk studies performed with *C. nodosa*, significant differences have also been found in the behaviour of the seeds of this species in populations from the Golfe Juan Bay in the French Mediterranean (Caye et al., 1992) and from the Island of Ischia in Naples (Pirc et al., 1986), although these differences were attributed to the genetic variability of this species inside the Mediterranean Sea. Consequently, it may be possible that the wide geographical distribution of a species could imply different tolerances among its populations.

Acknowledgements This research was financed by an ACUAMED contract and by an FPI grant (FPI 01 A 002) from the Generalitat Valenciana.

References

- Adams, J. B. & G. C. Bate, 1994. The ecological implications of tolerance to salinity by *Ruppia cirrhosa* (Petagna) Grande and *Zostera capensis* Setchell. *Botanica Marina* 37: 449–456.
- Arai, M., J. Y. Pak, K. Nomura & T. Nitta, 1991. Seawater-resistant, non-spherical protoplasts from seagrass leaves. *Physiologia Plantarum* 83: 551–559.
- Benjamin, K. J., D. I. Walker, A. J. McComb & J. Kuo, 1999. Structural response of marine and estuarine plants of *Halophila ovalis* (R.Br.) Hook. f. to long-term hypersalinity. *Aquatic Botany* 64: 1–17.
- Biebl, R. & C. P. McRoy, 1971. Plasmatic resistance of respiration and photosynthesis of *Zostera marina* at different salinities and temperatures. *Marine Biology* 8: 48–56.
- Caye, G. & A. Meinesz, 1986. Experimental study of seed germination in the seagrass *Cymodocea nodosa*. *Aquatic Botany* 26: 79–87.
- Caye, G., C. Bulard, A. Meinesz & F. Loquès, 1992. Dominant role of seawater osmotic pressure on germination in *Cymodocea nodosa*. *Aquatic Botany* 42: 187–193.
- Charpentier, A., P. Grillas, F. Lescuyer, E. Coulet & I. Auby, 2005. Spatio-temporal dynamics of a *Zostera noltii* dominated community over a period of fluctuating salinity in a shallow lagoon, Southern France. *Estuarine Coastal and Shelf Science* 64: 307–315.

- Chesnes, T. C. & C. L. Montague, 2001. The effects of salinity fluctuation on the productivity and osmoregulation of two seagrass species. Estuarine Research Federation Conference Abstracts, November 4–8, 2001, St. Petersburg, Florida.
- den Hartog, C., 1970. The Sea-Grasses of the World. North-Holland Publications Company, Amsterdam.
- Doering, P. H. & R. H. Chamberlain, 1998. Experimental studies on the salinity tolerance of turtle grass, *Thalassia testudinum*. In Bortone, S. A. (ed.), Workshop on Seagrasses. Subtropical and Tropical Seagrass Management Ecology: Responses to Environmental Stress. Fort Myers, Florida: 13.
- Drew, E. A., 1978. Factors affecting photosynthesis and its seasonal variation in the seagrasses *Cymodocea nodosa* (Ucria) Ascherson and *Posidonia oceanica* (L.) Delile in the Mediterranean. Journal of Experimental Marine Biology and Ecology 31: 173–194.
- Drew, E. A., 1979. Physiological aspects of primary production in seagrasses. Aquatic Botany 7: 139–150.
- Dring, M. J., 1992. The Biology of Marine Plants. Cambridge University Press, Cambridge.
- Druehl, L. D., 1981. Geographical distribution. In Lobban, L. S. & M. J. Wynne (eds), The Biology of Seaweeds. Blackwell, Oxford: 306–325.
- Fernández, J. A., M. J. García-Sánchez & H. H. Felle, 1999. Physiological evidence for a proton pump and sodium exclusion mechanisms at the plasma membrane of the marine angiosperm *Zostera marina* L. Journal of Experimental Botany 50: 1763–1768.
- Fernández-Torquemada, Y. & J. L. Sánchez-Lizaso, 2005. Effects of salinity on leaf growth and survival of the Mediterranean seagrass *Posidonia oceanica* (L.) Delile. Journal of Experimental Marine Biology and Ecology 320: 57–63.
- Fernández-Torquemada, Y., J. M. González-Correa & J. L. Sánchez-Lizaso, 2005a. Preliminary results of the monitoring of the brine discharge produced by the SWRO desalination plant of Alicante (SE Spain). Desalination 182: 395–402.
- Fernández-Torquemada, Y., M. J. Durako & J. L. Sánchez-Lizaso, 2005b. Effects of salinity and possible interactions with temperature and pH on growth and photosynthesis of *Halophila johnsonii* Eiseman. Marine Biology 148: 251–260.
- Fernández-Torquemada, Y., J. M. González-Correa, A. Loya, L. M. Ferrero, M. Díaz-Valdés & J. L. Sánchez-Lizaso, 2009. Dispersion of brine discharge from seawater reverse osmosis desalination plants. Desalination and Water Treatment 5: 137–145.
- Fukuhara, T., J. Y. Pak, Y. Ohwaki, H. Tsujimura & T. Nitta, 1996. Tissue-specific expression of the gene for a putative plasma membrane H^+ -ATPase in a seagrass. Plant Physiology 110: 35–42.
- Gacia, E., O. Invers, M. Manzanera, E. Ballesteros & J. Romero, 2007. Impact of the brine from a desalination plant on a shallow seagrass (*Posidonia oceanica*) meadow. Estuarine Coastal and Shelf Science 72: 579–590.
- Greve, T. M. & T. Binzer, 2004. Which factors regulate seagrass growth and distribution? In Borum, J., C. M. Duarte, D. Krause-Jensen & T. M. Greve (eds), European Seagrasses an Introduction to Monitoring and Management. The M&Ms Project, Hillerød: 19–23.
- Hillman, K., A. J. McComb & D. I. Walker, 1995. The distribution, biomass and primary production of the seagrass *Halophila ovalis* in the Swan/Canning Estuary, Western Australia. Aquatic Botany 51: 1–54.
- Hootsmans, M. J. M., J. E. Vermaat & W. van Vierssen, 1987. Seed bank development, germination and early seedling survival of two seagrass species from The Netherlands: *Zostera marina* L. and *Zostera noltii* Hornem. Aquatic Botany 28: 275–285.
- Jagels, R., 1973. Studies of a marine grass, *Thalassia testudinum* I. Ultrastructure of the osmoregulatory leaf cells. American Journal of Botany 60: 1003–1009.
- Kamermans, P., M. A. Hemminga & D. J. de Jong, 1999. Significance of salinity and silicon levels for growth of a formerly estuarine eelgrass (*Zostera marina*) population (Lake Grevelingen, The Netherlands). Marine Biology 133: 527–539.
- Kerr, E. A. & S. Strother, 1985. Effects of irradiance, temperature and salinity on photosynthesis of *Zostera muelleri*. Aquatic Botany 23: 177–183.
- Kinne, O., 1964. The effects of temperature and salinity on marine and brackish water animals. II. Salinity and temperature salinity combinations. In Barnes, H. (ed.), Oceanography and Marine Biology. An Annual Review. Haefner, New York: 281–339.
- Kirst, G. O., 1989. Salinity tolerance of eukaryotic marine algae. Annual Reviews of Plant Physiology. Plant Molecular Biology 40: 21–53.
- Kraemer, G. & L. Mazzella, 1999. Nitrogen acquisition, storage, and use by the co-occurring Mediterranean seagrasses *Cymodocea nodosa* and *Zostera noltii*. Marine Ecology Progress Series 183: 95–103.
- Kraemer, G., R. H. Chamberlain, P. H. Doering, A. D. Steinman & M. Hanisak, 1999. Physiological responses of transplant of the freshwater angiosperm *Vallisneria spiralis* along a salinity gradient in the Caloosahatchee Estuary (Southwestern Florida). Estuaries 22: 138–148.
- Lattemann, S. & T. Höpner, 2003. Seawater desalination. Impacts of brine and chemical discharges on the marine environment, Desalination Publications, L'Aquila.
- Loques, F., G. Caye & A. Meinesz, 1990. Germination in the marine phanerogam *Zostera noltii* Hornemann at Golfe Juan, French Mediterranean. Aquatic Botany 38: 249–260.
- Marbà, N., J. Cebrian, S. Enríquez & C. M. Duarte, 1996. Growth patterns of Western Mediterranean seagrasses: species-specific responses to seasonal forcing. Marine Ecology Progress Series 133: 203–215.
- Mazzella, L., M. B. Scipione, M. C. Gambi, M. C. Buia, M. Lorenti, V. Zupo & G. Cancemi, 1993. The Mediterranean seagrass *Posidonia oceanica* and *Cymodocea nodosa*. A comparative overview. First International Conference on the Mediterranean Coastal Environment, Antalya, Turkey: 103–116.
- McMillan, C. & F. N. Moseley, 1967. Salinity tolerances of five marine spermatophytes of Redfish Bay, Texas. Ecology 48: 503–506.
- Montague, C. L. & J. A. Ley, 1993. A possible effect of salinity fluctuation on abundance of benthic vegetation and

- associated fauna in Northeastern Florida Bay. *Estuaries* 16: 703–717.
- Ogata, E. & T. Matsui, 1965. Photosynthesis in several marine plants of Japan as affected by salinity, drying and pH, with attention to their growth habitats. *Botanica Marina* 8: 199–217.
- Pagès, J. F., M. Pérez & J. Romero, 2010. Sensitivity of the seagrass *Cymodocea nodosa* to hypersaline conditions: a microcosm approach. *Journal of Experimental Marine Biology and Ecology* 386: 34–38.
- Pak, J. Y., T. Fukuhara & T. Nitta, 1995. Discrete subcellular localization of membrane-bound ATPase activity in marine angiosperms and marine algae. *Planta* 196: 15–22.
- Pavón-Salas, N., R. Herrera, A. Hernández-Guerra & R. Haroun, 2000. Distributional pattern of seagrasses in the Canary Islands (Central-East Atlantic Ocean). *Journal of Coastal Research* 16: 328–335.
- Peduzzi, P. & A. Vuković, 1990. Primary production of *Cymodocea nodosa* in the Gulf of Trieste (northern Adriatic Sea): a comparison of methods. *Marine Ecology Progress Series* 64: 197–207.
- Peralta, G., F. G. Brun, I. Hernández, J. J. Vergara & J. L. Pérez-Lloréns, 2005. Morphometric variations as acclimation mechanisms in *Zostera noltii* beds. *Estuarine Coastal and Shelf Science* 64: 347–356.
- Pérez, M. & J. Romero, 1992. Photosynthetic response to light and temperature of the seagrass *Cymodocea nodosa* and the prediction of its seasonality. *Aquatic Botany* 43: 51–62.
- Pérez, M., C. M. Duarte, J. Romero, K. Sand-Jensen & T. Alcoverro, 1994. Growth plasticity in *Cymodocea nodosa* stands: the importance of nutrient supply. *Aquatic Botany* 47: 249–264.
- Pérez Ruzafa, A., C. Marcos, I. M. Pérez Ruzafa & J. D. Ros, 1987. Evolución de las características ambientales y de los doblamientos del Mar Menor (Murcia, SE España). *Anales de Biología* 12: 53–65.
- Philippart, C. J. M., 1995. Seasonal variation in growth and biomass of an intertidal *Zostera noltii* stand in the Dutch Wadden Sea. *Netherlands Journal of Sea Research* 33: 205–218.
- Phillips, R. C. & E. G. Meñez, 1988. *Seagrasses*. Smithsonian Institution Press, Washington.
- Pinnerup, S. P., 1980. Leaf production of *Zostera marina* L. at different salinities. *Ophelia Supplement* 1: 219–224.
- Pirc, H., M. C. Buia & L. Mazzella, 1986. Germination and seedling development of *Cymodocea nodosa* (Ucria) Ascherson under laboratory conditions and “in situ”. *Aquatic Botany* 26: 181–188.
- Ralph, P. J., 1998. Photosynthetic responses of *Halophila ovalis* (R. Br.) Hook. f. to osmotic stress. *Journal of Experimental Marine Biology and Ecology* 227: 203–220.
- Ralph, P. J., 1999. Photosynthetic response of *Halophila ovalis* (R.Br.) Hook. f. to combined environmental stress. *Aquatic Botany* 65: 83–96.
- Ramage, D. L. & D. R. Schiel, 1998. Reproduction in the seagrass *Zostera novaezelandica* on intertidal platforms in southern New Zealand. *Marine Biology* 130: 479–489.
- Reyes, J., M. Sansón & J. Afonso-Carrillo, 1995. Distribution and reproductive phenology of the seagrass *Cymodocea nodosa* (Ucria) Ascherson in the Canary Islands. *Aquatic Botany* 50: 171–180.
- Robblee, M. B., T. R. Barber, P. R. Carlson, M. J. Durako, J. W. Fourqurean, L. K. Muehlstein, D. Porter, L. A. Yarbro, R. T. Zieman & J. C. Zieman, 1991. Mass mortality of the tropical seagrass *Thalassia testudinum* in Florida Bay (USA). *Marine Ecology Progress Series* 71: 297–299.
- Sánchez-Lizaso, J. L., J. Romero, J. Ruiz, E. Gacia, J. L. Buceta, O. Invers, Y. Fernández-Torquemada, J. Mas, A. Ruiz-Mateo & M. Manzanera, 2008. Salinity tolerance of the Mediterranean seagrass *Posidonia oceanica*: recommendations to minimize the impact of brine discharges from desalination plants. *Desalination* 221: 602–607.
- Thomas, D. N., J. C. Collins & G. Russell, 1988. Interactive effects of temperature and salinity upon net photosynthesis of *Cladophora glomerata* (L.) Kütz. and *C. rupestris* (L.) Kütz. *Botanica Marina* 31: 73–77.
- Thorhaug, A. & J. Marcus, 1981. Mortality of *Thalassia testudinum* (Banks ex König) when exposed to the extremes of temperature, salinity and light. *American Journal of Botany* 68: 1102.
- Tomasko, D. A. & M. O. Hall, 1999. Productivity and biomass of the seagrass *Thalassia testudinum* along a gradient of freshwater influence in Charlotte Harbor, Florida. *Estuaries* 22: 592–602.
- Touchette, B. W., 2007. Seagrass-salinity interactions: physiological mechanisms used by submersed marine angiosperms for a life at sea. *Journal of Experimental Marine Biology and Ecology* 350: 194–215.
- Tyerman, S. D., 1989. Solute and water relations of seagrasses. In Larkum, A. W. D., A. J. McComb & S. A. Shepherd (eds), *Biology of Seagrasses*. Elsevier, Amsterdam: 729–759.
- Underwood, A. J., 1997. *Experiments in Ecology. Their Logical Design and Interpretation using Analysis of Variance*. Cambridge University Press, Cambridge.
- Underwood, A. J. & M. G. Chapman, 1997. Statistical program GMAV.5 for Windows. Institute of Marine Ecology, University of Sidney, Australia.
- van Katwijk, M. M., G. H. W. Schmitz, A. P. Gasseling & P. H. van Avesaath, 1999. Effects of salinity and nutrient load and their interaction on *Zostera marina*. *Marine Ecology Progress Series* 190: 155–165.
- Vermaat, J. E., J. A. J. Beijer, R. Gijlstra, M. J. M. Hootsmans, C. J. M. Philippart, N. W. van den Brink & W. van Vierrsen, 1993. Leaf dynamics and standing stocks of intertidal *Zostera noltii* Hornem and *Cymodocea nodosa* (Ucria) Ascherson on the Banc d’Arguin (Mauritania). *Hydrobiologia* 258: 59–72.
- Vermaat, J. E., F. C. A. Verhagen & D. Lindenburg, 2000. Contrasting responses in two populations of *Zostera noltii* Hornem. to experimental photoperiod manipulation at two salinities. *Aquatic Botany* 67: 179–189.
- Walker, D. I., 1985. Correlations between salinity and growth of the seagrass *Amphibolis antarctica* (Labill.) Sonder & Aschers., in Shark Bay, Western Australia, using a new method for measuring production rate. *Aquatic Botany* 23: 13–26.
- Walker, D. I. & A. J. McComb, 1990. Salinity response of the seagrass *Amphibolis antarctica* (Labill.) Sonder et

- Aschers.: an experimental validation of field results. Aquatic Botany 36: 359–366.
- Westphalen, G., E. O'Loughlin, G. Collings, J. Tanner, Y. Eglinton & S. Bryars, 2005. Responses to reduced salinities of the meadow forming seagrasses *Amphibolis* and *Posidonia* from the Adelaide metropolitan coast. ACWS technical report no 9 prepared for the Adelaide Coastal Waters Study Steering Committee. South Australian Research and Development Institute (Aquatic Sciences) Publication No. RD01/020814, Adelaide. <http://www.epa.sa.gov.au/pdfs/acws9.pdf>.
- Wortmann, J., J. W. Hearne & J. B. Adams, 1997. A mathematical model of an estuarine seagrass. Ecological Modeling 98: 137–149.
- Zieman, J. C., 1974. Methods for the study of the growth and production of turtle grass, *Thalassia testudinum* König. Aquaculture 4: 139–143.
- Zieman, J. C., 1975. Seasonal variation of turtle grass, *Thalassia testudinum* König, with reference to temperature and salinity effects. Aquatic Botany 1: 107–123.

UNCORRECTED PROOF