Alleviation of Zn toxicity by low water availability

Journal:	<i>Physiologia Plantarum</i>
Manuscript ID:	PPL-2013-00011.R2
Manuscript Type:	Regular manuscript - Ecophysiology, stress and adaptation
Date Submitted by the Author:	n/a
Complete List of Authors:	Disante, Karen; Departamento de Ecología. Universidad de Alicante, Ap 99, 03080 Alicante., ; Instituto Multidisciplinar para el Estudio del Medio "Ramon Margalef" (IMEM). Universidad de Alicante, Ap 99, 03080 Alicante., Cortina, Jordi; Departamento de Ecología. Universidad de Alicante, Ap 99, 03080 Alicante., ; Instituto Multidisciplinar para el Estudio del Medio "Ramon Margalef" (IMEM). Universidad de Alicante, Ap 99, 03080 Alicante., Vilagrosa, Alberto; Fundación Centro de Estudios Ambientales del Mediterráneo (CEAM), Joint Research Unit University of Alicante-CEAM, PO Box 99, 03080 Alicante., Fuentes, David; Fundación Centro de Estudios Ambientales del Mediterráneo (CEAM), Joint Research Unit University of Alicante-CEAM, PO Box 99, 03080 Alicante., Hernández, Encarni; Departamento de Ecología. Universidad de Alicante, PO Box 99, 03080 Alicante., Ljung, Karin; Umeå Plant Science Center, Department of Forest Genetics and Plant Physiology, Swedish University of Agricultural Sciences, SE-901 83 Umeå.,
Key Words:	heavy metal, drought, Zn application, auxin, Quercus suber, multiple stresses

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 Alleviation of Zn toxicity by low water availability
2
3 Karen B. Disante^{a,b*}, Jordi Cortina^{a,b}, Alberto Vilagrosa^c, David Fuentes^c, Encarni I. Hernández^b, Karin
4 Ljung^c
5
6 ^a Instituto Multidisciplinar para el Estudio del Medio “Ramon Margalef” (IMEM). Universidad de
7 Alicante, PO Box 99, 03080 Alicante, Spain
8 ^b Departamento de Ecología. Universidad de Alicante, PO Box 99, 03080 Alicante, Spain
9 ^c Fundación Centro de Estudios Ambientales del Mediterráneo (CEAM), Joint Research Unit
10 University of Alicante-CEAM, PO Box 99, 03080 Alicante, Spain.
11 ^d Umeå Plant Science Center, Department of Forest Genetics and Plant Physiology, Swedish
12 University of Agricultural Sciences, SE-901 83 Umeå, Sweden
13
14 * Corresponding author: kb.disante@ua.es; Tel.: + 34 965909564; fax: + 34 965 909 825.

17 Abstract

18 Heavy metal contamination and drought are expected to increase in large areas worldwide. However,
19 their combined effect on plant performance has been scantily analyzed. This study examines the effect
20 of Zn supply at different water availabilities on morpho-physiological traits of *Quercus suber* L in
21 order to analyze the combined effects of both stresses. Seedlings were treated with four levels of zinc
22 from 3 to 150 μM and exposed to low (LW) or high (HW) frequency of watering in hydroponic
23 culture, using a growth chamber. Under both watering regimes, Zn concentration in leaves and roots
24 increased with Zn increment in nutrient solution. Nevertheless, at the highest Zn doses, Zn tissue
25 concentrations were almost twice in HW than in LW seedlings. Functional traits as leaf photosynthetic
26 rate and root hydraulic conductivity, and morphological traits as root length and root biomass
27 decreased significantly in response to Zn supply. Auxin levels increased with Zn concentrations,
28 suggesting the involvement of this phytohormone in the seedling response to this element. LW
29 seedlings exposed to 150 μM Zn showed higher root length and root biomass than HW seedlings
30 exposed to the same Zn dose. Our results suggest that low water availability could mitigate Zn toxicity
31 by limiting internal accumulation. Morphological traits involved in the response to both stresses
32 probably contributed to this response.

34 Key words: heavy metal, drought, Zn application, auxin, *Quercus suber*, multiple stresses

36 Introduction

38 In nature, plants are commonly exposed to a combination of stress factors. Multiple stresses may
39 affect plants in ways that are hardly predictable from studies of single stress factors, as synergistic and
40 antagonistic interactions are widespread (Nilsen and Orcutt 1996, Larcher 2003). Water is the most
41 limiting factor for plant productivity worldwide and several studies have evaluated the combined
42 effect of drought and other sources of stress such as extreme temperature and radiation (Valladares
43 and Percy 1997, Aranda et al. 2005, Gimeno et al. 2008). In contrast, the combined effect of drought
44 and heavy metal contamination has received less attention, despite the potential interaction between
45 both stress factors (Poschenrieder and Barceló 2004, Brady et al. 2005; Sardans and Peñuelas 2007).
46 Several studies have shown that the effect of heavy metals on plant performance may be modulated by
47 other sources of stress as sulphur and proton concentration in rainfall, heat and frost (Kukkola et al.
48 1997, Öncel et al. 2000).

50 Among heavy metals, Zn is an essential micronutrient for higher plants and it is required for several
51 physiological processes (Marschner 1995, Cakmak 2000, Broadley et al. 2007). Zinc abundance is
52 increasing in natural systems as a consequence of human activities (Kabata-Pendias and Pendias 1992,
53 Fuentes et al. 2007a; Dominguez et al. 2008). When Zn concentration exceeds a critical level, it

1
2
3 54 becomes toxic (Balsberg Pålsson 1989, Woolhouse 1983). Symptoms of Zn toxicity include
4 55 alteration on biomass partitioning (Ruano et al. 1988), morphological changes in root system (Fuentes
5 56 et al. 2007b, Disante et al. 2010), reductions in photosynthetic rate and chlorophyll measurements (Di
6 57 Baccio et al. 2009, Disante et al. 2011), unbalance mineral nutrition (Kabata-Pendias and Pendias
7 58 1992, Monet et al. 2001) and modifications in xylem anatomy (Kasim 2007). These changes may
8 59 affect plant capacity to uptake and transport water to above ground parts (Barceló and Poschenrieder
9 60 1990, Kasim 2007).
10
11
12
13
14
15 62 High levels of Zn and low water availability may induce similar responses in vascular plants,
16 63 including stomatal closure and decreased CO₂ uptake, reduced chlorophyll fluorescence and foliar
17 64 area, and activation of defence mechanisms against oxidative damage (Vaillant et al. 2005, Valladares
18 65 et al. 2005). Conversely, plants may exhibit contrasted responses to Zn toxicity and drought. While
19 66 reduced root length and root biomass have been described in seedlings exposed to Zn (Wolhouse
20 67 1983), adaptive mechanisms in response to water limitation include deeper root systems and increased
21 68 proportion of biomass allocated belowground (Nicotra et al. 2002).
22
23
24
25
26 69 The interaction between Zn and drought may result in synergistic or antagonistic responses. Thus,
27 70 small increases in Zn availability (20 mg l⁻¹ of Zn) promoted growth in water-stressed soybeans
28 71 (Gadallah 2000). Similarly, heavy metal-enriched sewage sludge induced a positive effect on morpho-
29 72 physiological traits of seedlings subjected to drought (Pascual et al. 2004, Fuentes et al. 2007a; Santala
30 73 and Ryser 2009). Finally, in a previous study we observed that exposition of *Quercus suber* seedlings
31 74 to Zn delayed the effects of a short drought (Disante et al. 2011). High Zn concentrations could
32 75 hamper seedling capacity to absorb and transport water and compromise seedling ability to withstand
33 76 drought due to reduction of root length and biomass (Disante et al. 2010). Limited water availability
34 77 may also predispose leaves to cellular damage and photoinhibition, altering the carbon balance
35 78 (Werner et al. 2001, Vilagrosa et al. 2010). Finally, drought may impair uptake and acropetal
36 79 translocation of nutrients, including Zn (Hu and Schmidhalter 2005).
37
38
39
40
41
42
43
44 81 Despite the potential interaction between Zn and drought, studies dealing with the simultaneous effect
45 82 of both stress factors are scarce, and they frequently show conflictive results (Balsberg Pålsson
46 83 1989). The objective of this study is to evaluate the combined effect of Zn supply and water
47 84 availability on plant performance, and contribute to unveil the interactions between both sources of
48 85 stress. The underlying hypothesis is that morpho-physiological responses to drought may mitigate the
49 86 effects of high Zn availability. For this study we used cork oak (*Quercus suber* L.) seedlings, a
50 87 drought-resistant tree widely distributed in the western Mediterranean basin. Cork oak woodlands
51 88 produce valuable goods and services, including cork, and they are protected by the European Union
52 89 (Habitat directive 92/43/EEC; (Pausas et al. 2009)). In addition, we performed this experiment with
53 90 young seedlings because this stage is one of the most sensitive for plant capacity to resist stress
54
55
56
57
58
59
60

conditions (Vallejo et al. 2000, Vilagrosa et al. 2003a). To account for this objective, we analyzed plant physiological status, root capacity to supply water to leaves, and phytohormonal response in several Zn supply rates under two levels of water availability.

Materials and methods

Plant material and growing conditions

We cultivated *Q. suber* seedlings from acorns. Seeds from local origin were supplied by the Forest Service Seed Bank (Banco de Semillas El Serranillo, Ministerio de Medio Ambiente y Medio Rural y Marino). *Quercus suber* acorns were sown in silica sand and kept at $24 \pm 2^\circ\text{C}$. Two weeks after germination, when the radicle was approximately 2 cm long, acorns were transplanted into 5 cm diameter and 30 cm length tubes filled with fine, medium and coarse silica sand (the size of sand ranged from 0.5 to 2.2 mm) in a proportion corresponding to 1:2:3 by volume, respectively. The substrate was tested in the lab before the onset of the experiment to ensure proper drainage. The bottom of the tubes was closed with a 1 x 1 mm mesh, which prevented substrate loss. Tubes were placed in a growth chamber under controlled conditions of temperature ($22 \pm 1^\circ\text{C}/18 \pm 1^\circ\text{C}$, day/night), 12 h photoperiod ($600 \mu\text{mol m}^{-2} \text{s}^{-1}$ of photosynthetic photon flux density during the day) and 50% relative humidity. During this period, seedlings were irrigated with a nutrient solution containing 500 μM KCl, 435 μM NaNO_3 , 300 μM $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 45 μM NaH_2PO_4 , 300 μM CaCl_2 , 0.37 μM $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 0.012 μM $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, 1.6 μM H_3BO_3 , 0.006 μM KI, 0.6 μM $\text{MnSO}_4 \cdot 7\text{H}_2\text{O}$ and 0.0004 μM $(\text{NH}_4)_6\text{Mo}_7\text{O}_{24}$. This solution was a modification of the one used by Arduini et al. (1994) to simulate forest soil conditions.

Experimental design

Eight weeks after germination, we added ZnSO_4 to the nutrient solution to achieve a Zn concentration of 3 (Control), 10, 50 and 150 μM , corresponding to those commonly found in uncontaminated and contaminated soils (Knight et al. 1997, Kabata-Pendias and Pendias 1992, González et al. 2011). We applied two irrigation regimes using this solution: high watering (HW) and low watering (LW). HW seedlings were irrigated three times a week (Monday, Wednesday and Friday), whereas LW seedlings were irrigated every 5 days. During the experiment, seedlings were irrigated with 60 ml of nutrient solution. Substrate moisture content (SMC) was estimated by measurements of water content by gravimetric method. SMC for HW and LW treatments ranged from $2.0 \pm 0.1\%$ to $7.4 \pm 0.1\%$ and from $0.3 \pm 0.1\%$ to $6.40 \pm 0.2\%$ respectively. The moisture content (SMC) for the substrate during 18 consecutive days is shown in Fig. 1 as example of SMC dynamics.

Each Zn level (hereafter treatments) was replicated 10 times, i.e., 10 seedlings per Zn treatments and irrigation level. Ion speciation calculations for solutions containing different amounts of Zn were conducted using MINTEQA2 software (Allison Geoscience Consultants, Inc. Flowery Branch, Georgia, USA and HydroGeoLogic, Inc. Herndon, Virginia, USA). The concentration in mol/L of all anions and cations present in the nutrient solution (described above) were entered in MINTEQA2 software to estimate the equilibrium composition of the experimental solution. pH was fixed at 5.5 (the pH of the nutrient solution was adjusted to this value) and quartz was added as the mineral present in the equilibrium. Calculations showed that 98% of Fe was present as Fe (II) and 81% of Zn was present as free ion and 18% as ZnNO_3^{3+} . Both elements were highly available for plants, as only small precipitations of PO_4^{-3} (0.2%) and Mn^{+2} (16.2%) were found by chemical equilibrium estimations using MINTEQA2 (see Appendix S1 in Supporting information). Leachates obtained immediately after watering showed that Zn concentration in the solution was always close to the intended concentration, independently of water availability and plant uptake. For 3 (Control), 10, 50 and 150 μM Zn were: 4.2 ± 0.4 , 10.3 ± 3.7 , 47.6 ± 8.7 ; 150.2 ± 12.9 μM Zn and 2.8 ± 0.4 , 7.8 ± 1.3 , 46.8 ± 6.8 , 162.9 ± 16.3 μM Zn for HW and LW seedlings, respectively. Twelve weeks after the onset of the treatments, when seedlings were 5-months-old, we assessed their physiological status, and measured hydraulic conductivity and IAA concentration. Finally, we quantified their morphological traits.

Gas exchange and chlorophyll fluorescence

Physiological status was assessed 5 months after the onset of the experiment on 5 seedlings per treatment level (i.e. 40 seedlings in total). Gas exchange was analysed with a Li-Cor 6400 portable infrared gas analyser system (Li-Cor Inc. Lincoln, Nebraska, USA). We determined photosynthetic rate (A), stomatal conductance (g_s) and transpiration rate (E) in one fully expanded leaf per seedling. From these data, we calculated the instantaneous water use efficiency (IWUE). We maintained gas flow rate at 350 mL min^{-1} , CO_2 concentration at 400 ppm and a photon flux density of 600 $\mu\text{mol m}^{-2} \text{s}^{-1}$ during the course of the measurements. Previous determinations of light response curves showed that a value of 600 $\mu\text{mol m}^{-2} \text{s}^{-1}$ was sufficient to saturate the PSII photosystem in seedlings of this species. Measurements were conducted early in the morning (09:00–11:00 h solar time). A portable fluorometer (pulse-amplitude modulated photosynthesis yield analyzer, PAM-2010 Walz, Effeltrich, Germany) was used to measure chlorophyll fluorescence from intact leaves. In dark adapted leaves, we measured the minimum fluorescence yield (F_0) under weak red light and the maximum fluorescence (F_m) after a saturating pulse of white light. Maximal PSII photochemical efficiency ($F_v:F_m$) was calculated as $F_v:F_m = (F_m - F_0) / F_m$ according to Genty et al. (1989)

Root hydraulic conductance

165

166 Four to five seedlings per Zn dose and irrigation regime were used to measure water transport capacity
 167 of complete and intact root systems by means of a High Pressure Flowmeter (HPFM, Dynamax Inc.,
 168 Houston, USA). Measures were carried out at the same time that gas exchange variables were
 169 measured. The night before the measurements, seedlings were watered to pot capacity, which is the
 170 amount of water remaining in a pot after irrigation and visible drainage had ceased. Root hydraulic
 171 conductance (K_R) measurements were carried out early in the morning. Root systems were kept in
 172 their substrate and perfused with distilled and de-gassed ultra-pure water filtered through a 0.2 μm
 173 water filtration membrane using the high pressure flow meter method (HPFM, Dynamax, USA)
 174 connected to the stump. K_R was measured in a transient way within a range of 0–0.5 MPa pressure at
 175 a constant increment rate of 3–5 kPa s^{-1} and the flow rate (F) was recorded every 3 s as described in
 176 Tyree et al. (1995). K_R ($\text{Kg m s}^{-1} \text{MPa}^{-1}$) was calculated as the slope of ΔF (Kg s^{-1}) versus ΔP (MPa) as
 177 $K_R = dF/dP$. K_R was corrected for variations in water temperature. Morphological variables were used
 178 to normalize root hydraulic conductance (K_R) on root surface area basis (K_{R-RA} ; $\text{Kg m}^{-2} \text{s}^{-1} \text{MPa}^{-1}$).

180 Seedling morphology

181

182 Five months after the onset of the experiment, seedlings were harvested, stems cut above the root
 183 collar and the rooting system carefully washed. Roots and leaves were digitized by scanning on an A3
 184 flatbed scanner (Epson Expression 1680 Pro, Long Beach, California, USA) fitted with a transparency
 185 adaptor at 300 dpi, using an 8-bit greyscale. Images were analyzed with specific software (WinRhizo,
 186 Regent Instruments, Québec, Canada) to evaluate projected leaf area, total root length and total root
 187 surface area. All biomass fractions were dried at 65°C for 48 h, and then weighed. Specific root length
 188 (SRL) was calculated as the ratio between total root length and root biomass.

190 Zn concentration in plant tissues

191

192 Dry roots and leaves were digested in a heating block at 250°C with a mixture of sulphuric acid and
 193 hydrogen peroxide (Jones and Case 1990). Digests were analyzed for Zn by ICP-OES (Perkin Elmer
 194 Optima 4300 Inductively Coupled Plasma Optical Emission Spectrometry). *Olea europaea* L. leaf
 195 standard reference material (BCR: CRM 062, Commission of the European Communities Bureau of
 196 Reference, Brussels) was digested and analysed for quality control.

197

198 Quantification of indole-3-acetic-acid (IAA)

199

200 Frozen leaves (10 mg) were homogenised in 0.5 ml of 0.05 M sodium phosphate buffer (pH 7.0)
 201 containing 0.02% diethyldithiocarbamic acid as an antioxidant and 500 pg^{13}C_6 -IAA as an internal

standard, using the Retsch vibration mill (Retsch MM 200 mixer mill, Retsch GmbH, Haan, Germany) and a 3-mm tungstencarbide bead at a frequency of 30 Hz for 3 min. The pH was adjusted to 2.7 with 1 M HCl, and the samples were purified by solid phase extraction (SPE) and derivatized as described Andersen et al. (2008). Finally, samples were dissolved in 30 µl heptane and analyzed by GC-SRM-MS (gas chromatography - selected reaction monitoring - mass spectrometry) as described in Edlund et al. 1995.

Data analysis

Treatment effects on physiological and morphological variables were evaluated using two-way ANOVA. When the interaction term was significant, one-way ANOVA and Student-t were used to evaluate the effect of Zn at each watering level, and the effect of increased water availability at each Zn level, respectively. Root biomass and instantaneous water use efficiency were raised to a power of 0.5 and root Zn concentration was transformed using natural logarithm prior to analysis, to homogenize the variance. Factor levels were compared pairwise (Tukey's HSD test) when ANOVA showed a significant treatment effect. We used regression analysis to evaluate the significance of the relationship between K_R and A and g_s . All analyses were performed using the SPSS v.15.0 statistical package (SPSS Inc., Chicago, Illinois, USA).

Results

Zn bioaccumulation

Foliar Zn concentration increased in HW and LW seedlings with increasing Zn concentration in the growing medium (Fig. 2A). The magnitude of the increase was dependent on water availability (Table 1, Table 2). Thus, at the highest Zn dose, its concentration was two-fold higher in leaves of HW seedlings than in leaves of LW seedlings. The Zn bioaccumulation in roots was affected by the increase in Zn load and by the interaction between both factors (Fig 2B, Table 1).

Morphological response

Foliar biomass was not affected by Zn application, water availability or the interaction between both (Table 1 and 3). On the contrary, root biomass showed a significant effect of the increase in Zn load, and the interaction between Zn application and water availability. While root biomass in HW seedlings decreased with increasing Zn dose, LW seedlings showed the opposite trend (Table 2). At the highest Zn dose, root biomass accumulation was 33% lower in HW seedlings than in LW

seedlings. This response was similar for root length, but changes in root length in LW seedlings with increasing Zn concentration were not statistically significant (Table 2). We found a marginally significant trend towards lower specific root length as Zn dose increased at any watering level (Table 1 and 3).

Physiological response

Maximum PSII photochemical efficiency ($F_v:F_m$) ranged between 0.65 and 0.72. $F_v:F_m$ was significantly decreased by Zn but not by water availability (Table 1). The interaction between Zn and watering regime was marginally significant as the effect of drought on this variable decreased as Zn dose increased above 3 μM Zn (Table 1, Appendix S2 in Supporting Information). In HW seedlings, $F_v:F_m$ decreased with the increase of Zn while F_o showed the opposite trend (Table 2). We found no significant effect of Zn application in $F_v:F_m$ and F_o in LW seedlings.

Zinc application, water availability and their interaction significantly affected gas exchange (Table 1, Fig. 3A and 3B). HW seedlings receiving 3 μM Zn, had 2.8 and 3.9 times higher net photosynthetic rate (A) and stomatal conductance (g_s) than LW seedlings, respectively. Stomatal conductance and A decreased with increasing Zn dose in HW (Table 2). In contrast, A in LW seedlings was weakly affected by Zn application, and g_s showed a unimodal response to increase in Zn load (Table 2). Instantaneous water-use-efficiency (IWUE) increased as Zn dose increased and, in contrast to other variables, this effect was independent of the irrigation regime (see Appendix S3 in Supporting Information).

Hydraulic conductance

Root hydraulic conductance in a root area basis (K_{R-RA}) was sensitive to Zn doses, watering regime and their interaction (Fig. 4, Table 1). K_{R-RA} gradually decreased as Zn availability increased in HW seedlings, whereas K_{R-RA} was not affected by Zn doses in LW seedlings (Fig. 4, Table 2). K_{R-RA} was significantly higher in HW seedlings than in LW seedlings at low Zn availability, but this difference disappeared at Zn concentrations of 50 μM and higher (Appendix S4 in Supporting Information). Root hydraulic conductance per unit of leaf area and root hydraulic conductance per unit of root length showed similar trends (data not shown).

IAA concentration

Foliar IAA concentration was significantly affected by the increase in Zn load, and marginally affected by the interaction between Zn application and water availability (Fig. 5, Table 1). This was

the result of a substantial increase in IAA in HW seedlings receiving 150 μM of Zn compared to HW seedlings receiving lower Zn doses and LW seedlings receiving any Zn dose (Table 2).

Discussion

In Mediterranean type ecosystems, water stress is the main constraint to plant performance (Di Castri 1973, Rhizopoulou and Mitrakos 1990). In these areas, prolonged summer drought, low water storage capacity of forest soils and high summer temperatures could amplify the effects of other sources of stress such as heavy metals, high temperatures or high radiation levels. In addition, the deleterious impact of both sources of stress may be enlarged in the future as a consequence of climate change and increased inputs of heavy metals (Sardans and Peñuelas 2007, Micó et al. 2006, IPCC 2001). Results of the present study show that water availability and heavy metal contamination may indeed interact in complex ways, and they suggest that the combined effects of both sources of stress may not be additive simply.

Zn effects on seedling performance

Zinc concentration in leaves of seedlings exposed to 3 μM Zn was similar to concentrations found in 1-year-old *Q. suber* (Robert et al., 1996). Foliar Zn concentration reached critical levels (e.g., 200 $\mu\text{g g}^{-1}$ d.w.; Balsberg Pahlsson, 1989) only after exposition to 150 μM Zn. Maximum Zn concentration belowground in HW seedlings was 1.7 higher than in LW seedlings. Similar values of Zn bioaccumulation have been found in other woody species grown in hydroponic culture (e.g., 3444 $\mu\text{g g}^{-1}$ to 5700 $\mu\text{g g}^{-1}$; Reichman et al., 2001; Fuentes et al., 2007b).

Aboveground biomass accumulation showed no evidence of Zn toxicity. Similar response had been observed previously (Disante et al. 2011). This could be a consequence of the time span of the experiment and low Zn concentration in shoots, as belowground biomass accumulation and ecophysiological status (as leaf photosynthetic rate) were affected by Zn application. In addition, stem elongation in oaks follows successive flushes (Johnson et al. 2001). Thus, we might observe changes in shoot biomass in response to the increase in Zn load after a new flush of growth. Reductions in aboveground biomass accumulation and chlorosis in seedlings exposed to high levels of Zn have been reported in other woody species in experiments of different time span (Reichman et al. 2001, Fuentes et al. 2007b, Disante et al. 2010).

On the contrary, belowground parts were more sensitive to increase in Zn load than aboveground parts as reflected by the reduction in root biomass and the marginally significant decrease in root length of well watered seedlings receiving increasing doses of heavy metal. The reduction in root biomass accumulation is a common and quick response to high Zn availability (Ruano et al. 1988). Indeed Zn

concentration in roots was up to ten times higher than in shoots. This may be a result of the strong control of Zn translocation of *Q. suber*, possibly excluder species (Disante et al. 2010).

Functionality of *Q. suber* leaves was negatively affected by the increase in Zn load. Previous studies suggested that Zn increases plant sensitivity to photoinhibition (Joshi and Mohanty 2004, Mateos-Naranjo et al. 2008). The decline observed in $F_v:F_m$ may be due to damage in the reaction centre and down-regulation processes (Osmond et al. 1999). In our experiment, this reduction was caused by a weak increase in F_o that can be induced by damages to the PSII reaction centre (Moustakas et al. 1997), including reductions in the energy transfer from the antennae to the reaction centre associated with chronic photoinhibition (Ralph and Burchett 1998, Popovic et al. 2003). Similarly, we observed a reduction in A and g_s in seedlings exposed to Zn application, as described in previous works (Myśliwa-Kurziel et al. 2004, Disante et al. 2011). Impaired carbon fixation may partly reflect photoinhibition and photon damage, but also the reduction in g_s .

Another direct effect over g_s may be associated with the decrease in water transport capacity in roots affected by increment of this heavy metal. Supporting this hypothesis, we found a reduction in hydraulic conductance of roots and a positive relationship between K_R and gas exchange rates ($R^2=0.64$, $F=10.66$, $P=0.0171$, and $R^2=0.73$, $F=15.99$, $P=0.0071$ for A and g_s respectively). These results are in agreement with observations on stomata sensitivity to decreasing K_R , and the prevention of xylem cavitation (Brodribb and Holbrook 2003, Vilagrosa et al. 2003b, Otieno et al. 2007). The negative effect of the increase in Zn load altering root morphology supports our observations in root functionality (i.e. decreases of K_R). Heavy metals may modify root hydraulic conductance by different mechanisms, but studies on this topic are relatively scarce (Poschenrieder and Barceló 2004). Thus, previous studies found that Hg can block aquaporins in cell membranes (Lee et al. 2005), and Zn and Cd may foster the deposition of phenolic compounds and the reduction in vessel size (Fuhrer 1982, Poschenrieder et al. 1989). Independently of the mechanisms involved, the reduction of K_R and the concomitant decrease in A and g_s , together with changes in root biomass accumulation and root morphology, suggest that Zn may hamper the ability of *Q. suber* seedlings to access and transport soil resources to aboveground parts. The development of extended rooting system is important to ensure plant survival and early growth in Mediterranean conditions, and especially for seedling establishment in areas subjected to intense droughts (Padilla and Pugnaire, 2007, León et al. 2011).

Hormonal response to heavy metals is another response poorly studied, especially in woody species. We found a substantial increase in IAA in leaves of well watered seedlings as Zn concentration increased. Auxins are mainly synthesized in shoots, and then translocated and accumulated in roots (Marschner 1995, Ljung et al. 2005). This hormone is involved in several process related to the

1
2
3 350 coordination of root and shoot growth and development, participating in processes as cell elongation,
4 351 formation and differentiation of vascular tissue, and in the shaping of root architecture (Mattsson et al.
5 352 2003; Aloni et al. 2006, Robert and Friml 2009). Our results show that the increase in Zn load may
6 353 elicit alterations in endogenous auxin levels and suggest that this could promote the observed
7 354 alterations in root morphology and functionality. Indeed, other studies have found changes in auxin
8 355 accumulation after Cu exposition, which was related to morphological changes as the formation of
9 356 lateral roots (Lequeux et al. 2010, Pető et al. 2011).

10 357
11
12 358 Combined effects of Zn and water availability

13 359
14
15 360 We found no significant effect of the increase in Zn load on aboveground biomass accumulation in
16 361 HW seedlings, whereas aboveground biomass showed a marginally significant increase with Zn
17 362 application in LW seedlings. Several studies have found increases in plant growth in response to small
18 363 amounts of available Zn (Gadallah 2000, Fuentes et al. 2007b). In contrast, belowground biomass
19 364 increased and decreased in LW and HW seedlings, respectively, as Zn availability increased. The
20 365 marginal increase in root biomass in LW seedlings was probably due to the combination of small
21 366 changes in root diameter (data not shown) and root length (Table 3): 5% and 12% increase,
22 367 respectively, in seedlings exposed to 150 μ M Zn compared to seedlings exposed to 3 μ M Zn. We may
23 368 note that root biomass and root length were 1.5 times higher in LW seedlings than in HW seedlings
24 369 when both sets received 150 μ M Zn. A study combining Zn and salinity found similar results, as Zn
25 370 application reduced root length and root biomass accumulation in unstressed seedlings, but it increased
26 371 root growth in seedlings subjected to saline stress (Gadallah and Ramadan 1997).
27 372 *Quercus suber* displays intermediate tolerance to drought (Ramírez-Valiente et al. 2009). In our
28 373 experiment, we did not test intense and extended drought that could probably affect root anatomy. Our
29 374 objective was to study how the reduction in the frequency of irrigation influences seedling response to
30 375 different Zn doses. This may explain why contrasted watering frequency did not induce changes in
31 376 root length or root biomass in seedlings receiving 3 μ M Zn. Data on Table 1 show that the effects of
32 377 Zn and watering frequency on several seedling traits depend on the levels of the other factors. The
33 378 interaction between drought and Zn points out at a synergistic or antagonistic effect of the two factors.
34 379 Table 2 and Table 3 show that Zn had opposite effects on seedlings depending on the watering level.
35 380 Despite that the effect of Zn on root architecture, (i.e., root biomass) was marginally significant in LW
36 381 seedlings ($P= 0.052$), we consider this effect to be important and worth discussing. Besides, low
37 382 watering induced a marginal decrease in root average diameter compared to HW (Appendix S5 in
38 383 Supporting information). Results of a two-way ANOVA showed the lack of a significant interaction
39 384 between Zn and drought, and indicate that roots of seedlings receiving low irrigation were thinner than
40 385 HW seedlings. In other words, LW induced changes in root morphology.

Physiological response to increasing Zn availability also differed in LW and HW seedlings. Low irrigation frequency decreased gas exchange rates, and probably limited Zn uptake and accumulation. This may explain the lower foliar and root Zn concentrations found in LW seedlings compared with HW seedlings, and may be partially responsible for the smaller effect of Zn doses observed in the former seedlings. Zinc is incorporated by specific transporters (Lasat et al. 2000, Takahashi et al. 2012) and to our knowledge there is no report about the effect of water availability on these transporters. But drought generally reduces nutrient transport because it impairs transpiratory flow and active transport (Hu and Schmidhalter 2005).

All physiological traits measured in *Q. suber* decreased as Zn concentration increased in HW seedlings, whereas most traits showed a unimodal response, albeit not statistically significant, to increasing Zn in LW seedlings. Stomatal conductance and carbon assimilation rates of LW seedlings receiving the lowest Zn corresponded to a values obtained in *Q. suber* seedling exposed to mild drought (Disante et al. 2011), but they were not further reduced by the increase in Zn load. As a consequence, physiological status of LW and HW seedlings strongly differed in seedlings receiving the lowest doses of Zn, but differences gradually disappeared as Zn availability increased, or Zn concentration in roots increased above ca. $1000 \mu\text{g g}^{-1}$ (Appendix S2). When water was limiting, the addition of $50 \mu\text{M}$ Zn increased A and g_s in a 67% and 73% as compared with $3 \mu\text{M}$ Zn, respectively. Moreover, in the last case this increment was significant. These results are in agreement with observed trends in morphological variables.

In the same way, K_{R-RA} showed a similar behaviour as gas exchange rates. K_{R-RA} decreased with increasing Zn concentration in LW seedlings and increases a 30% after $50 \mu\text{M}$ Zn. This may be a consequence of the morphological changes occurring in LW plants subjected to increased Zn availability. As a result, K_{R-RA} values converged in HW and LW seedlings at root Zn concentration above ca. $1000 \mu\text{g g}^{-1}$ (see Appendix S4 in Supporting Information). Similar results were observed when we analysed K_R on a leaf area and a root length basis. Limitations to water flow through xylem vessels can increase the risk of cavitation by increasing water tension from roots to leaves (Sperry et al. 2002). However, some limitation to water flow may increase IWUE of plants by regulating water losses. This response would be beneficial to maintain a positive water balance, especially for plants living in water limited ecosystems (Vilagrosa et al. 2003a, Hernández et al. 2010).

The effect of heavy metals on IWUE has been scarcely studied, but an increase in IWUE can alleviate the effects of drought. We found that the increase in Zn load enhanced carbon assimilation per unit of water loss independently of the irrigation regime. This is remarkable, as A and g_s behaved in quite different ways in both groups of seedlings.

To what extent was the differential morpho-physiological response of LW seedlings to Zn a simple consequence of reduced Zn uptake? The relationships between root Zn concentration, on the one hand,

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

424 and g_s and K_R , on the other, did not overlap in LW and HW seedlings (see Appendix S4 in Supporting
425 Information). These results suggest that the response of seedlings to low water availability influence
426 the response of *Q. suber* seedlings to Zn application.
427
428

For Peer Review

Conclusion

Our results demonstrate that the combination of the increase in Zn load and moderate water stress did not induce a synergistic response in *Q. suber* seedlings, supporting our initial hypothesis. The combination of both stress factors, may contribute to alleviate their negative impact. On the one hand, low water availability prevented Zn accumulation, and probably buffered its negative effects. On the other hand, when water was limiting, intermediate Zn concentrations may have contributed to the improvement of traits as water use efficiency and root growth, buffering the effects of water limitation. Low water availability expanded the range of Zn concentrations where this heavy metal may have a positive effect on plant performance, shifting the toxicity level up. Our results suggest that legal and practical regulations of Zn levels should take into account the water regime of each zone.

Acknowledgements

We thank Jose Huesca for their assistance in the lab, Banco de Semillas El Serranillo for the seeds used in this experiment and the University of Alicante for an FPU (Formacion del Profesorado Universitario) grant awarded to K. Disante. This research was funded by the Spanish Ministry of Science and Innovation (Project GRACCIE, Programa Consolider-Ingenio 2010 (CSD 2007-00067) and SURVIVE (CGL-2011-30531-CO2-02)) and Generalitat Valenciana (FEEDBACKS-PROMETEO/2009/006). E.I. Hernández thanks the University of Alicante for her FPU research fellowship. CEAM is supported by Generalitat Valenciana.

References

- Aloni R, Aloni E, Langhans M, Ullrich CI (2006) Role of cytokinin and auxin in shaping root architecture: regulating vascular differentiation, lateral root initiation, root apical dominance and root gravitropism. *Ann Bot* 97: 883–893
- Andersen SU, Buechel S, Zhao Z, Ljung K, Novák O, Busch W, Schuster C, Lohmann JU (2008) Requirement of B2-type cyclin-dependent kinases for meristem integrity in *Arabidopsis thaliana*. *Plant Cell* 20: 88 – 100
- Aranda I, Castro L, Alia R, Pardos JA, Gil L (2005) Low temperature during winter elicits differential responses among populations of the Mediterranean evergreen cork oak (*Quercus suber*). *Tree Physiol* 25: 1085 – 1090

1
2
3 465 Arduini I, Godbold DL, Onnis A (1994) Cadmium and copper change root growth and morphology of
4 466 *Pinus pinea* and *Pinus pinaster* seedlings. *Physiol Plant* 92: 675 – 680
5 467
6
7 468 Balsberg Pålsson AM (1989) Toxicity of heavy metals (Zn, Cu, Cd, Pb) to vascular plants. A
8 469 literature review. *Water Air Soil Poll* 47: 287 – 319
9 470
10
11 471 Barceló J, Poschenrieder Ch (1990) Plant water relations as affected by heavy metal stress: a review. *J*
12 472 *Plant Nutr* 13: 1 – 37
13 473
14
15 474 Brady KU, Kruckeberg AR, Bradshaw Jr HD (2005) Evolutionary ecology of plant adaptation to
16 475 serpentine soils. *Ann Rev Ecol Evol S* 36: 243 – 66
17 476
18 477 Broadley MR, White PJ, Hammond JP, Zelko I, Lux, A (2007) Zinc in plants. *New Phytol* 173: 677 –
19 478 702
20 479
21 480 Brodribb TJ, Holbrook NM (2003) Stomatal closure during leaf dehydration, correlation with other
22 481 leaf physiological traits. *Plant Physiol* 132: 2166 – 2173
23 482
24 483 Cakmak I (2000) Possible roles of Zinc in protecting plant cells from damage by reactive oxygen
25 484 species. *New Phytol* 146: 185 – 205
26 485
27 486 Di Baccio D, Tognetti R, Minnocci A, Sebastiani L (2009) Responses of the *Populus × euramericana*
28 487 clone I-214 to excess zinc: Carbon assimilation, structural modifications, metal distribution and
29 488 cellular localization. *Environ Exp Bot* 67: 153 – 163
30 489
31 490 Di Castri F (1973) Climatographical comparisons between Chile and the western coast of North
32 491 America. In: Castri F, Mooney HA (eds) *Mediterranean type-ecosystems*, Springer-Verlag, Berlin,
33 492 Germany, pp 21 – 36
34 493
35 494 Disante KB, Fuentes D, Cortina J (2010) Sensitivity to zinc of Mediterranean woody species important
36 495 for restoration. *Sci Tot Environ* 408: 2216 – 2225
37 496
38 497 Disante KB, Fuentes D, Cortina J (2011) Response to drought of Zn-stressed *Quercus suber* L.
39 498 seedlings. *Environ Exp Bot* 70: 96 – 103
40 499
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- Domínguez MT, Marañón T, Murillo JM, Schulin R, Robinson BH (2008) Trace element accumulation in woody plants of the Guadiamar Valley, SW Spain: a large-scale phytomanagement case study. *Environ Pollut* 152: 50 – 59
- Edlund A, Eklof S, Sundberg B, Moritz T, Sandberg G (1995) A microscale technique for gas chromatography-mass spectrometry measurements of picogram amounts of indole-3-acetic acid in plant tissues. *Plant Physiol* 108: 1043 – 1047
- Fuhrer J (1982) Early effects of excess cadmium uptake in *Phaseolus vulgaris*. *Plant Cell Environ* 5: 263 – 270
- Fuentes D, Disante KB, Valdecantos A, Cortina J, Vallejo VR (2007a) Response of *Pinus halepensis* Mill. seedlings to biosolids enriched with Cu, Ni and Zn in three Mediterranean forest soils. *Environ Pollut* 145: 316 – 323
- Fuentes D, Disante KB, Valdecantos A, Cortina J, Vallejo VR (2007b) Sensitivity of Mediterranean woody seedlings to copper, nickel and zinc. *Chemosphere* 66: 412 – 420
- Gadallah MAA (2000) Effects of indole-3-acetic acid and zinc on the growth, osmotic potential and soluble carbon and nitrogen components of soybean plants growing under water deficit. *J Arid Environ* 44: 451 – 467
- Gadallah MAA, Ramadan T (1997) Effects of zinc and salinity on growth and anatomical structure of *Carthamus tinctorius* L. *Biol Plantarum* 39: 411 – 418
- Genty B, Briantais JM, Baker NR (1989) The relationship between the quantum yield of photosynthetic electron transport and quenching of chlorophyll fluorescence. *Biochim Biophys Acta* 990: 87 – 92
- Gimeno TE, Pas B, Lemos-Filho JP, Valladares F (2008) Plasticity and stress tolerance override local adaptation in the responses of Mediterranean holm oak seedlings to drought and cold. *Tree Physiol* 29: 87 – 98
- Hernández EI, Vilagrosa A, Pausas JG, Bellot J (2010) Morphological traits and water use strategies in seedlings of Mediterranean coexisting species. *Plant Ecol* 207: 233 – 244

1
2
3 535 Hu Y, Schmidhalter U (2005) Drought and salinity: A comparison of their effects on mineral nutrition
4 536 of plants. J Plant Nutr Soil Sc 168: 541 – 549
5
6 537
7 538 IPCC (2001) Climate change 2001: the scientific basis. Contribution of working group I to the Third
8
9 539 Assessment Report of the Intergovernmental Panel on Climate Change. In: Houghton JT, Dung Y
10 540 Griggs DJ, Noguer M, Van der Linden PJ, Dui X, Maskell K, Johson CA (eds.) Third Assessment
11 541 Report of Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge.
12
13 542
14
15 543 Jones Jr JB, Case VW (1990) Sampling, handling, and analyzing plant tissue samples. In: Westerman
16 544 RL (ed) Soil Testing and Plant Analysis, SSSA Book Series 3, Madison, Wisconsin, pp 389 – 427
17
18 545
19
20 546 Johnson PS, Shifley SR, Rogers R (2001) The ecology and silviculture of oaks. CABI Publishing.
21 547 Oxon, UK
22
23 548
24 549 Joshi MK, Mohanty P (2004) Chlorophyll a fluorescence as a probe of heavy metal ion toxicity in
25 550 plants. In: Papageorgiou GC, Govindjee (eds) Chlorophylla fluorescence. A signature of
26 551 photosynthesis, Springer Netherlands, pp 637 – 661
27
28 552
29
30 553 Kabata-Pendias A, Pendias H (1992) Trace elements in soils and plants. CRC Press, Boca Raton, FL
31 554
32
33 555 Kasim WA (2007) Physiological consequences of structural and ultra-structural changes induced by
34 556 Zn stress in *Phaseolus vulgaris*. I. Growth and photosynthetic apparatus. Int J Bot 3: 15 – 22
35
36 557
37
38 558 Kukkola E, Huttunen S, Bäck J, Rautio P (1997) Scots pine needle injuries at subarctic industrial sites.
39 559 Trees 11: 378 – 387
40
41 560
42
43 561 Larcher W (2003) Physiological plant ecology: Ecophysiology and stress physiology of functional
44 562 groups. 4th edition, Springer, Berlin, pp 513
45
46 563
47 564 Lasat MM, Pence NS, Garvin DF, Ebbs SD, Kochian LV (2000) Molecular physiology of zinc
48 565 transport in the Zn hyperaccumulator *Thlaspi caerulescens*. J Exp Bot 51: 71 – 79
49
50 566
51
52 567 Lee SH, Chung GC, Steudle E (2005) Low temperature and mechanical stresses differently gate
53 568 aquaporins of root cortical cells of chilling-sensitive cucumber and resistant figleaf gourd. Plant Cell
54 569 Environ 28: 1191 – 1202
55
56 570
57
58
59
60

- 571 León MF, Squeo FA, Gutiérrez JR, Holmgren M (2011) Rapid root extension during water pulses
572 enhances establishment of shrub seedlings in the Atacama Desert. *J Veg Sci* 22: 120 – 129
573
- 574 Lequeux H, Hermans C, Lutts S, Verbruggen N (2010) Response to copper excess in *Arabidopsis*
575 *thaliana*: Impact on the root system architecture, hormone distribution, lignin accumulation and
576 mineral profile. *Plant Physiol Biochem* 48: 673 – 682
577
- 578 Ljung K, Hull AK, Celenza J, Yamada M, Estelle M, Normanly J, Sandberg, G (2005) Sites and
579 regulation of auxin biosynthesis in *Arabidopsis* roots. *Plant Cell* 17: 1090 – 1104
580
- 581 Marschner H (1995) Mineral Nutrition of Higher Plants. Academic Press Inc, London
582
- 583 Mateos-Naranjo E, Redondo-Gómez S, Cambrollé J, Luque T, Figueroa ME (2008) Growth and
584 photosynthetic responses to zinc stress of an invasive cordgrass, *Spartina densiflora*. *Plant Biol* 10:
585 754 – 762
- 586 Mattsson J, Ckurshumova W, Berleth T (2003) Auxin signaling in *Arabidopsis* leaf vascular
587 development. *Plant Physiol* 131: 1327 – 1339
588
- 589 Micó C, Recatala L, Peris M, Sánchez J (2006) Assessing heavy metal sources in agricultural soils of
590 an European Mediterranean area by multivariate analysis. *Chemosphere* 65: 863 – 872
591
- 592 Monnet F, Vaillant N, Vernay P, Coudret A, Sallanon H, Hitmi A (2001) Relationship between PSII
593 activity, CO₂ fixation, and Zn, Mn and Mg contents of *Lolium perenne* under zinc stress. *J Plant*
594 *Physiol* 158: 1137 – 1144
595
- 596 Moustakas M, Eleftheriou EP, Ouzounidou G (1997) Short-term effects of aluminium at alkaline pH
597 on the structure and function of the photosynthetic apparatus. *Photosynthetica* 34: 169 – 177
598
- 599 Myśliwa-Kurdziel B, Prasad MNV, Strzalka K (2004) Photosynthesis in heavy metal stressed plants.
600 In: Prasad MNV (ed) Heavy Metal Stress in Plants. From Molecule to Ecosystems, Springer, Berlin,
601 pp 146 – 181
602
- 603 Nicotra AB, Babicka N, Westoby M (2002) Seedling root anatomy and morphology: an examination
604 of ecological differentiation with rainfall using phylogenetically independent contrasts. *Oecologia*
605 130: 136 – 145

1
2
3 606
4 607 Nilsen ET, Orcutt DM (1996) The physiology of plants under stress: abiotic factors. John Wiley and
5 608 Sons, New York
6 609
7
8 610 Öncel I, Keles Y, Üstün AS (2000) Interactive effects of temperature and heavy metal stress on the
9 611 growth and some biochemical compounds in wheat seedlings. Environ Pollut 107: 315 – 320
10 612
11
12 613 Osmond B, Schwartz O, Gunning B (1999) Photoinhibitory printing on leaves, visualised by
13 614 chlorophyll fluorescence imaging and confocal microscopy, is due to diminished fluorescence from
14 615 grana. Aust J Plant Physiol 26: 717 – 724
15 616
16
17 617 Otieno DO, Schmidt MWT, Kurz-Besson C, Vale RLD, Pereira JS, Tenhunen JD (2007) Regulation
18 618 of transpirational water loss in *Quercus suber* trees in a Mediterranean-type ecosystem. Tree Physiol
19 619 27: 1179 – 1187
20 620
21
22 621 Pascual I, Antolín MC, García C, Polo A, Sánchez-Díaz M (2004) Plant availability of heavy metals in
23 622 a soil amended with a high dose of sewage sludge under drought conditions. Biol Fert Soils 40: 291 –
24 623 299
25 624
26
27 625 Pausas J, Pereira J, Aronson J (2009) The tree. In: Aronson J, Pereira JS, Pausas JG (eds) Cork Oak
28 626 Woodlands on the Edge. Ecology, Adaptive Management and Restoration, Island Press, Washington
29 627 DC, pp 11 – 25
30
31
32 628 Padilla FM, Pugnaire FI (2007) Rooting depth and soil moisture control Mediterranean woody
33 629 seedling survival during drought. Funct Ecol 21: 489 – 495
34
35
36
37 630 Pető A, Lehotai N, Lozano-Juste J, León J, Tari I, Erdei L, Kolbert Z (2011) Involvement of nitric
38 631 oxide and auxin in signal transduction of copper-induced morphological responses in *Arabidopsis*
39 632 seedlings. Ann Bot 108: 449 – 457
40
41
42
43
44
45
46
47 633 Poschenrieder C, Gunse B, Barceló J (1989) Influence of cadmium on water relations, stomatal
48 634 resistance, and abscisic acid content in expanding bean leaves. Plant Physiol 90: 1365 – 1371
49
50
51 635 Poschenrieder C, Barceló J (2004) Water relations in heavy metal stressed plants. In: Prasad MNV
52 636 (ed) Heavy Metal Stress in Plants. From Molecule to Ecosystems, Springer, Berlin, pp 249 – 271
53
54 637
55
56
57
58
59
60

- 638 Popovic R, Dewez D, Juneau P (2003) Application of chlorophyll fluorescence in ecotoxicology:
639 heavy metals, herbicides, and air pollutants. In: Dell JR, Toivinen PMA (eds) Practical applications of
640 chlorophyll fluorescence in plant biology, Kluwer Academic Publishers, Boston, pp 151 – 184
641
- 642 Ralph PJ, Burchett MD (1998) Photosynthetic response of *Halophila ovalis* to heavy metal stress.
643 Environ Pollut 103: 91 – 101
644
- 645 Ramírez-Valiente JA, Valladares F, Gil L, Aranda I (2009) Population differences in juvenile survival
646 under increasing drought are mediated by seed size in cork oak (*Quercus suber* L.). Forest Ecology
647 and Management 257: 1676 – 1683.
648
- 649 Reichman SM, Asher CJ, Mulligan DR, Menzies NW (2001) Seedling responses of three Australian
650 tree species to toxic concentrations of zinc in solution culture. Plant Soil 235: 151 – 158
651
- 652 Rhizopoulou S, Mitrakos K (1990) Water relations of evergreen sclerophylls. I. Seasonal changes in
653 the water relations of eleven species from the same environment. Ann Bot 65: 171 – 178
654
- 655 Robert B, Berton G, Sayag D, Masson P (1996) Assessment of mineral nutrition of cork oak through
656 foliar analysis. Commun. Soil Sci Plant Anal. 27 (9–10): 2091–2109.
- 657 Robert HS, Friml J (2009) Auxin and other signals on the move in plants. Nat Chem Biol 5: 325 – 332
658
- 659 Ruano A, Poschenrieder Ch, Barceló J (1988) Growth and biomass partitioning in zinc-toxic bush
660 beans. J Plant Nutr 11: 577 – 588
661
- 662 Santala KR, Ryser P (2009) Influence of heavy-metal contamination on plant response to water
663 availability in white birch, *Betula papyrifera*. Environ Exp Bot 66: 334 – 340
664
- 665 Sardans J, Peñuelas J (2007) Drought changes the dynamics of trace element accumulation in a
666 Mediterranean *Quercus ilex* forest. Environ Pollut 147: 567 – 583
667
- 668 Sperry JS, Hacke UG, Oren R, Comstock JP (2002) Water deficits and hydraulic limits to leaf water
669 supply. Plant Cell Environ 25: 251 – 263
670
- 671 Takahashi R, Ishimaru Y, Shimo H, Ogo Y, Senoura T, Nishizawa NK, Nakanishi H (2012) The
672 OsHMA2 transporter is involved in root-to-shoot translocation of Zn and Cd in rice. Plant Cell
673 Environ 35: 1948 – 1957

1
2
3 674
4 675 Tyree MT, Patiño S, Bennink J, Alexander J (1995) Dynamic measurements of roots hydraulic
5 676 conductance using a high-pressure flowmeter in the laboratory and field. J Exp Bot 46: 83 – 94
6 677
7
8 678 Vaillant N, Monnet F, Hitmi A, Sallanon H, Coudret A (2005) Comparative study of responses in four
9 679 *Datura* species to a zinc stress. Chemosphere 59: 1005 – 1013
10 680
11
12 681 Valladares F, Dobarro I, Sánchez-Gómez D, Pearcy RW (2005) Photoinhibition and drought in
13 682 Mediterranean woody saplings: scaling effects and interactions in sun and shade phenotypes. J Exp
14 683 Bot 56: 483 – 494
15 684
16
17 685 Valladares F, Pearcy RW (1997) Interactions between water stress, sun-shade acclimation, heat
18 686 tolerance, and photoinhibition in the sclerophyll *Heteromeles arbutifolia*. Plant Cell Environ 20: 25 –
19 687 36
20 688
21
22 689 Vallejo VR, Bautista S, Cortina J (2000) Restoration for soil protection after disturbances. In: Trabaud
23 690 L (ed) Life and Environment in the Mediterranean, WIT Press, Southampton, pp 199 – 208
24 691
25
26 692 Vilagrosa A, Cortina J, Gil-Pelegrin E, Bellot J (2003a) Suitability of drought preconditioning
27 693 techniques in Mediterranean climate. Restor Ecol 11: 208 – 216
28 694
29
30 695 Vilagrosa A, Vallejo VR, Bellot J, Gil-Pelegrin E (2003b) Cavitation, stomatal conductance, and leaf
31 696 dieback in seedlings of two co-occurring Mediterranean shrubs during an intense drought. J Exp Bot
32 697 54: 2015 – 2024
33 698
34
35 699 Vilagrosa A, Morales F, Abadía A, Bellot J, Cochard H, Gil-Pelegrin E (2010) Are symplast tolerance
36 700 to intense drought conditions and xylem vulnerability to cavitation coordinated? An integrated
37 701 analysis of photosynthetic, hydraulic and leaf level processes in two Mediterranean drought-resistant
38 702 species. Environ Exp Bot 69: 233 – 242
39 703
40
41 704 Werner C, Ryel RJ, Correia O, Beyschlag W (2001) Effects of photoinhibition on whole-plant carbon
42 705 gain assessed with a photosynthesis model. Plant Cell Environ 24: 27 – 40
43 706
44
45 707 Woolhouse HW (1983) Toxicity and tolerance in the responses of plants to metals. In: Lange OL (ed)
46 708 Encyclopedia of plant physiology, Springer-Verlag, Berlin, pp 245 – 300
47 709
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1. Results of two-way ANOVA to evaluate the effects of Zn and water availability (w) on morpho-physiological traits of *Quercus suber* seedlings. Foliar Zn: foliar Zn concentration, Root Zn: root Zn concentration, SRL: specific root length, $F_v:F_m$: predawn maximum photochemical efficiency, F_o : minimum fluorescence, A : photosynthesis; g_s : stomatal conductance, $IWUE$: instantaneous water use efficiency, K_{R-RA} : root hydraulic conductance per root area, IAA: auxin concentration in leaves.

Variables	F_{Zn}	P_{Zn}	F_w	P_w	$F_{Zn \times w}$	$P_{Zn \times w}$
Foliar Zn	$F_{3,26} = 105.34$	<0.001	$F_{1,26} = 12.78$	0.001	$F_{3,26} = 3.34$	0.034
Root Zn	$F_{3,26} = 46.67$	<0.001	$F_{1,26} = 0.01$	0.920	$F_{3,26} = 2.86$	0.056
Foliar biomass	$F_{3,48} = 1.95$	0.135	$F_{1,48} = 0.01$	0.905	$F_{1,48} = 2.00$	0.127
Root biomass	$F_{3,50} = 2.95$	0.041	$F_{1,50} = 0.94$	0.337	$F_{3,50} = 5.11$	0.004
Root length	$F_{3,48} = 0.25$	0.858	$F_{1,48} = 0.02$	0.870	$F_{3,48} = 3.29$	0.028
SRL	$F_{3,45} = 2.44$	0.076	$F_{1,45} = 0.93$	0.339	$F_{3,45} = 0.13$	0.943
$F_v:F_m$	$F_{3,30} = 5.09$	0.006	$F_{1,30} = 2.14$	0.154	$F_{3,30} = 2.36$	0.091
F_o	$F_{3,30} = 2.66$	0.066	$F_{1,30} = 1.18$	0.287	$F_{3,30} = 2.85$	0.054
A	$F_{3,30} = 3.74$	0.021	$F_{1,30} = 32.52$	<0.001	$F_{3,30} = 3.97$	0.017
g_s	$F_{3,30} = 4.04$	0.016	$F_{1,30} = 31.35$	<0.001	$F_{3,30} = 4.00$	0.017
$IWUE$	$F_{3,30} = 2.86$	0.054	$F_{1,30} = 2.71$	0.111	$F_{3,30} = 0.40$	0.756
K_{R-RA}	$F_{3,32} = 2.96$	0.047	$F_{1,32} = 22.76$	<0.001	$F_{3,32} = 6.96$	0.001
IAA	$F_{3,23} = 3.58$	0.029	$F_{1,23} = 0.08$	0.773	$F_{3,23} = 2.84$	0.060

Table 2. Results of one-way ANOVA to evaluate the effects of Zn on morpho-physiological traits of *Quercus suber* seedlings subjected to high and low watering frequency (HW and LW, respectively). Foliar Zn: foliar Zn concentration, Root Zn: root Zn concentration, $F_v:F_m$: predawn maximum photochemical efficiency, F_o : minimum fluorescence; A : photosynthesis; g_s : stomatal conductance, K_{R-RA} : root hydraulic conductance per root area, IAA: auxin concentration in leaves.

Variables	F_{HW}	P_{HW}	F_{LW}	P_{LW}
Foliar Zn	$F_{3,13} = 92.33$	<0.001	$F_{3,13} = 31.51$	<0.001
Root Zn	$F_{3,14} = 99.27$	<0.001	$F_{3,14} = 35.55$	<0.001
Root biomass	$F_{3,23} = 4.23$	0.016	$F_{3,27} = 2.93$	0.052
Root length	$F_{3,22} = 2.57$	0.080	$F_{3,26} = 1.35$	0.281
$F_v:F_m$	$F_{3,15} = 6.66$	0.004	$F_{3,15} = 0.86$	0.485
F_o	$F_{3,15} = 4.88$	0.015	$F_{3,15} = 0.31$	0.819
A	$F_{3,16} = 5.02$	0.012	$F_{3,15} = 2.46$	0.106
g_s	$F_{3,16} = 4.61$	0.017	$F_{3,15} = 3.34$	0.050
K_{R-RA}	$F_{3,15} = 7.30$	0.003	$F_{3,18} = 0.63$	0.602
IAA	$F_{3,11} = 4.39$	0.029	$F_{3,12} = 0.90$	0.468

Table 3. Morphological attributes of 5-months-old *Quercus suber* seedlings grown under a range of Zn supply rates and contrasted irrigation regimes. HW and LW correspond to high and low watering frequency, respectively. Means and standard errors of N= 6-7 seedlings are showed.

Different capital letters and lowercase letters indicate significant differences at $P < 0.05$ for HW seedlings and LW seedling respectively (Tukey's HSD test). Asterisks indicate significant differences at $P < 0.05$ between seedlings HW and LW exposed to the same Zn dose (t-Student).

	Zn application rate (μM)							
	3		10		50		150	
	HW	LW	HW	LW	HW	LW	HW	LW
Foliar biomass (g)	0.47 $\pm$ 0.10	0.45 $\pm$ 0.05	0.50 $\pm$ 0.05	0.56 $\pm$ 0.05	0.71 $\pm$ 0.09	0.54 $\pm$ 0.03	0.48 $\pm$ 0.08	0.64 $\pm$ 0.01
Root biomass (g)	2.43 $\pm$ 0.33 ^{AB}	2.02 $\pm$ 0.15	2.93 $\pm$ 0.33 ^B	2.43 $\pm$ 0.15	2.86 $\pm$ 0.28 ^B	2.35 $\pm$ 0.16	1.75 $\pm$ 0.18 ^A	2.59 $\pm$ 0.13*
Root length (cm)	1479 $\pm$ 119	1306 $\pm$ 123	1283 $\pm$ 180	1199 $\pm$ 105	1398 $\pm$ 217	1193 $\pm$ 94	976 $\pm$ 105	1482 $\pm$ 132*
Specific root length (cm g ⁻¹)	593 $\pm$ 72	660 $\pm$ 42	481 $\pm$ 70	493 $\pm$ 41	494 $\pm$ 42	549 $\pm$ 76	560 $\pm$ 42	578 $\pm$ 48

Figure 1. Changes in substrate moisture content (SMC) for HW and LW seedlings during 18 consecutive days. HW and LW correspond to high and low watering frequency, respectively. Means and standard errors of N= 6-7 tubes per treatment are shown.

Figure 2. Foliar (A) and root (B) Zn concentration of *Quercus suber* seedlings growing at four Zn doses and two contrasting water availability levels (HW and LW for high and low watering frequency, respectively). Bars are means and standard errors of N=5 seedlings. Note the different scale used for leaves and roots. Note that data distribution along the X-axis is not proportional.

Different capital letters and lowercase letters indicate significant differences at $P < 0.05$ for HW seedlings and LW seedling respectively (Tukey's HSD test). Asterisks indicate significant differences at $P < 0.05$ between seedlings HW and LW exposed to the same Zn dose (t-Student).

Figure 3. Photosynthesis (A) and stomatal conductance (B) of *Quercus suber* seedlings exposed to different levels of Zn availability and contrasting watering frequency. Bars are means and standard errors of N=5 seedlings. Note that data distribution along the X-axis is not proportional.

Different capital letters and lowercase letters indicate significant differences at $P < 0.05$ for HW seedlings and LW seedling respectively (Tukey's HSD test). Asterisks indicate significant differences at $P < 0.05$ between seedlings HW and LW exposed to the same Zn dose (t-Student).

Figure 4. Specific hydraulic conductance per root surface area in *Quercus suber* seedlings exposed to different levels of Zn availability and contrasting watering frequency. Bars are means and standard errors of N=4-5 seedlings. Note that data distribution along the X-axis is not proportional.

Different capital letters and lowercase letters indicate significant differences at $P < 0.05$ for HW seedlings and LW seedling respectively (Tukey's HSD test). Asterisks indicate significant differences at $P < 0.05$ between seedlings HW and LW exposed to the same Zn dose (t-Student).

Figure 5. Indole-3-acetic acid (IAA) concentration in leaves of *Quercus suber* seedlings exposed to different levels of Zn availability and contrasting watering frequency. Bars are

means and standard errors of N=3-4 seedlings. Note that data distribution along the X-axis is not proportional.

Different capital letters and lowercase letters indicate significant differences at $P < 0.05$ for HW seedlings and LW seedling respectively (Tukey's HSD test). Asterisks indicate significant differences at $P < 0.05$ between seedlings HW and LW exposed to the same Zn dose (t-Student).

For Peer Review

Figure 1

Figure 2.

Figure 3

Figure 4

Figure 5