

**ENANTIOSELECTIVE SYNTHESIS OF TETRAHYDROQUINOLINE
ALKALOIDS (-)-ANGUSTUREINE AND (-)-CUSPAREINE FROM
CHIRAL *TERT*-BUTANESULFINYL IMINES.**

Juan A. Sirvent, Francisco Foubelo,* and Miguel Yus*

Department of Organic Chemistry, Faculty of Science, and Institute of Organic
Synthesis (ISO), University of Alicante, Apdo. 99, 03080 Alicante, Spain

Abstract – The addition of a Grignard reagent to both enantiomeric *N-tert*-butanesulfinyl imines derived from 3-(2-bromophenyl)propanal **8** proceeded with high diastereoselectivity. The resulting sulfinamides **9** and **12** were easily transformed into tetrahydroquinoline alkaloids (-)-angustureine (**4**) and (-)-cuspatureine (**5**) after three steps: *N*-desulfinylation, intramolecular *N*-arylation and *N*-methylation.

INTRODUCTION

The enantioselective synthesis of 2-substituted tetrahydroquinolines has attracted much attention among organic chemists because they are widespread subunits in natural alkaloids and pharmaceuticals as well. For instance, simple synthetic 1,2,3,4-tetrahydroquinolines with interesting biological properties includes oxamniquine (**1**), which has been used to combat parasitic disease schistosomiasis,¹ the novel antibiotic virantmycin (**2**),² and the tetrahydroquinoline L-689,560 (**3**),³ one of the most potent NMDAR (*N*-methyl *d*-aspartate receptor) antagonists yet described (Figure 1). On the other hand, among natural alkaloids, (-)-angustureine (**4**), (-)-cuspatureine (**5**), (-)-galipeine (**6**) and (-)-galipinine (**7**) were isolated from *Galipea officinalis* Hancock (Figure 1),⁴ a Venezuelan shrubby tree used in folk medicine as tonic in dyspepsia, dysentery, chronic diarrhea and also as antipyretic.⁵ The majority of the enantioselective synthesis of these tetrahydroquinolines reported until today are based on asymmetric catalytic hydrogenation of the corresponding quinolines.⁶ More recently, Helmchen provided a synthesis of (+)-angustureine and (-)-cuspatureine involving a highly regio- and enantioselective intermolecular iridium-catalyzed allylic amination as a key step.⁷ Continuing our interest in the use of *N-tert*-butanesulfinyl imines⁸ as





Figure 1.

electrophiles, we envisioned a straightforward synthesis of (-)-angustureine (4) and (-)-cuspareine (5), based on the diastereoselective addition of Grignard reagents to these chiral imines,⁹ which have found high applicability in synthesis because both enantiomers are accessible in large-scale processes¹⁰ and because the chiral auxiliary is easily removed under acidic conditions. In addition, practical processes for recycling the *tert*-butanesulfinyl group upon deprotection of *N*-*tert*-butanesulfinyl amines have also been reported.¹¹ In our synthetic strategy, a palladium-catalyzed intramolecular *N*-arylation will be also involved (Scheme 1).




Scheme 1.

RESULTS AND DISCUSSION

As depicted in Scheme 1, our synthesis commenced with the diastereoselective addition of the corresponding Grignard reagent to (*E*)-*N*-[3-(2-bromophenyl)propylidene]-*tert*-butanesulfinamide (8). Chiral imines 8 were easily accessible from the corresponding *tert*-butanesulfinamide and 3-(2-bromophenyl)pronal (obtained from commercially available 1,2-bromiodobenzene and allylic alcohol through a type-Heck coupling reaction).¹² Ellman and co-workers reported for the first time the addition of Grignard reagents to *N*-*tert*-butanesulfinyl aldimines.¹³ Interestingly, they found a marked


solvent effect on the stereoselectivity of the addition, no coordinating solvents, such as dichloromethane, leading to the best results.¹⁴ In addition, and considering the configuration of the resulting diastereomers, a chelated transition state **I** was proposed, which was also consistent with the observed solvent effect (competitive coordination of ethereal solvents such as THF would interfere with the formation of the six-membered-ring transition state). According to this model, the attack of the Grignard reagent occurred on the *Re* face of the imine with the *S* configuration at the sulfur atom (Scheme 2). For that reason, we studied first the addition of 1M solution of pentylmagnesium bromide in ether to the imine (*S*)-**8** in different solvents (Scheme 2). The addition was carried out at -78 °C, and after that the reaction mixture was allowed to reach room temperature. The highest diastereoselectivity was obtained performing the reaction in toluene (94:6 dr), although similar levels of diastereocontrol were achieved using dichloromethane and diethyl ether (92:8 and 93:7 dr, respectively). As it could be anticipated, the lowest diastereoselectivity was reached in THF as solvent. The diastereomeric ratios were easily determined by ¹H-NMR analysis of the crude reaction mixtures through the comparison of the integrals of the *t*-Bu group and the N–H for each of the diastereoisomer (the largest chemical shift difference was always observed for the diastereomeric signals of the N–H). In all cases complete conversion was observed using two equivalents of the Grignard reagent, giving rise to the expected product **9** with *R* configuration at the newly created stereogenic center (Scheme 2).



Scheme 2.


Free amine **10** was generated in quantitative yield from sulfinamide **9** upon reaction with 6M HCl in THF at room temperature. Intramolecular *N*-arylation to produce the tetrahydroquinoline derivative **11** was

achieved in 81% yield under palladium catalysis, using Cs₂CO₃ as a base in toluene in a high pressure tube at 115 °C for 20 hours.^{12,15} Finally, *N*-methylation of **11** was achieved with paraformaldehyde and sodium cyanoborohydride leading to (-)-angustureine (**4**) in 82% yield (Scheme 3).



The same strategy was applied to the synthesis of (-)-cuspareine (**5**), starting in this case from the enantiomeric aldimine (*R*)-**8**, since the configuration at the C2 position of the tetrahydroquinoline unit is the opposite to that of the (-)-angustureine (**4**). The reaction of 1.5 equivalents of 2-(3,4-dimethoxyphenyl)ethylmagnesium bromide (prepared from the corresponding alkyl bromide and magnesium in THF at 67 °C) with aldimine (*R*)-**8** in toluene proceeded in this case in moderate yield (47%) and stereoselectivity (86:14 dr). Acidic hydrolysis of sulfonamide **12** led to free amine **13** in high yield. Further palladium-catalyzed intramolecular *N*-arylation gave tetrahydroquinoline derivative **14** in lower yield when comparing with the angustureine precursor shown above. Methylation of **14** produced (-)-cuspareine (**5**) in 80% yield (Scheme 4).

In summary, 2-substituted tetrahydroquinoline alkaloids (-)-angustureine (**4**) and (-)-cuspareine (**5**) were easily accessible from the imines derived from 3-(2-bromophenyl)propanal and (*S*)- or (*R*)-*tert*-butanesulfinamide, respectively. The methodology presented here comprised as key steps a diastereoselective addition of a Grignard reagent to a chiral sulfinyl imine, and a palladium-catalyzed intramolecular *N*-arylation.



Scheme 4.

EXPERIMENTAL

All chemicals were commercially available (Acros, Aldrich). TLC was performed on Merck silica gel 60 F₂₅₄, using aluminum plates and visualized with phosphomolybdic acid (PMA) stain. Chromatographic purification was performed by flash chromatography using Merck silica gel 60 (0.040-0.063 mm) and hexane/EtOAc as eluent. GC-MS was developed in an Agilent 6890N spectrometer with FID detector, helium gas transportation (2 mL/min), 12 psi injection pressure, 270 °C temperature in detection and injection blocks, 1.0 μL volume of sample and 5 mm/min registration speed. Program temperature: 60 °C, initial temperature for 3 min; heating 15 °C/min; 270 °C, final temperature. Column type HP-1, 12 m long, 0.22 mm internal diameter, 0.25 μm thickness methylsilicone rubber and OV-101 stationary phase. IR spectra were measured (film) with a Nicolet Impact 510 P-FT Spectrometer. NMR spectra were recorded with a Bruker AC-300 using CDCl₃ as the solvent and TMS as internal standard. Optical rotations were measured on a Perkin Elmer 341 polarimeter.

(3*R,S*)-*N*-(*tert*-Butanesulfinyl)-1-(2-bromophenyl)octan-3-amine (9).- To solution of aldimine (*S*)-**8** (322 mg, 1.0 mmol) in dry toluene (4 mL) was added dropwise a 1M solution of pentylmagnesium bromide in diethyl ether (2.0 mmol, 2.0 mL) at -78 °C. The reaction was allowed to reach room temperature and after 4 h, it was cooled down to 0 °C, hydrolyzed with water (5 mL) and extracted with EtOAc (4 × 15 mL). The organic layers were successively washed with water (15 mL), brine (10 mL) and then dried over magnesium sulfate and concentrate under vacuum (15 Torr). The residue was purified by column chromatography (silica gel, hexane/EtOAc: 4/1) to yield 291 mg of pure product **9** (75% yield, 94:6 dr). Physical and spectroscopic data follow: Yellow oil; $[\alpha]_D^{20} +28$ (*c* 1.15, CH₂Cl₂); *R*_f 0.48 (hexane/EtOAc: 1/1); IR ν (film) 1622, 1471, 1084, 1025, 749 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 0.88 (3H, t, *J* = 6.8 Hz, CH₃), 1.25 [9H, s, (CH₃)₃], 1.27-1.47 (6H, m, CH₂), 1.53-1.93 (4H, m, CH₂), 2.70 (1H, ddd, *J* = 13.5, 11.1, 5.6 Hz, CH₂), 2.90 (1H, ddd, *J* = 13.5, 11.1, 5.1 Hz, CH₂), 3.08 (1H, br d, *J* = 6.9 Hz, NH), 3.24-3.37 (1H, m, CH), 7.02-7.09 (1H, m, ArH), 7.19-7.24 (2H, m, ArH), 7.52 (1H, d, *J* = 7.9 Hz, ArH); ¹³C NMR (75 MHz, CDCl₃) δ 14.2 (CH₃), 22.7 (CH₂), 22.9 (CH₃), 25.5 (CH₂), 31.8 (CH₂), 32.6 (CH₂), 36.0 (CH₂), 36.6 (CH₂), 56.0 (C), 56.8 (CH), 124.4 (C), 127.7 (CH), 127.8 (CH), 130.4 (CH), 133.0 (CH), 141.5 (C); LRMS (EI) *m/z* 331 [*M*⁺ - 58 (⁸¹Br), 5%], 329 (5), 260 (7), 258 (7), 250 (24), 185 (33), 183 (33), 171 (46), 169 (48), 147 (100), 130 (19), 104 (29), 103 (17), 91 (31), 90 (17), 89 (10), 83 (14), 77 (28), 70 (11), 55 (10).

(3*R*)-1-(2-Bromophenyl)octan-3-amine (10).- To solution of sulfinamide **9** (245 mg, 0.63 mmol) in THF (1.5 mL) was added dropwise a 6M solution of HCl (0.6 mmol, 0.10 mL) at 0 °C. The reaction mixture was stirred at 20 °C for 1 h. Then it was cooled down to 0 °C and basified with 2M NaOH (1.0 mmol, 0.5 mL). The reaction mixture was extracted with EtOAc (3 × 15 mL), dried over magnesium sulfate, filtered through a short pad of Celite and concentrated under vacuum (15 Torr) to give 177 mg of pure product **10** (99%). Physical and spectroscopic data follow: Yellow oil; $[\alpha]_D^{20} -6$ (*c* 1.07, CH₂Cl₂); *R*_f 0.30 (CH₂Cl₂/methanol: 9/1); IR ν (film) 2925, 1469, 1022, 747 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 0.89 (3H, t, *J* = 6.7 Hz, CH₃), 1.20-1.40 (7H, m, CH₂), 1.40-1.63 (2H, m, CH₂), 1.67-1.90 (3H, m, CH₂), 2.69-2.92 (3H, m, CH₂, NH₂), 6.99-7.09 (1H, m, ArH), 7.18-7.25 (2H, m, ArH), 7.52 (1H, d, *J* = 7.7 Hz, ArH); ¹³C NMR (75 MHz, CDCl₃) δ 14.2 (CH₃), 22.8 (CH₂), 25.9 (CH₂), 32.1 (CH₂), 33.0 (CH₂), 38.0 (CH₂), 38.3 (CH₂), 51.2 (CH), 124.5 (C), 127.6 (2 x CH), 130.4 (CH), 132.9 (CH), 141.8 (C); LRMS (EI) *m/z* 284 [*M*⁺ (⁸¹Br), 1%], 282 (1), 214 (60), 212 (62), 204 (43), 171 (34), 169 (35), 100 (100).

(*R*)-2-Pentyltetrahydroquinoline (11).- To a solution of amine **10** (152 mg, 0.53 mmol) in dry toluene (5 mL) in a high pressure flask was successively added triphenylphosphine (21 mg, 0.081 mmol), Pd(OAc)₂ (6

mg, 0.027 mmol) and Cs₂CO₃ (345 mg, 1.06 mmol). The reaction mixture was stirred at 115 °C for 20 h. The mixture was allowed to reach room temperature and hydrolyzed, first with 3M HCl (3.0 mmol, 1 mL) and then with 2M NaOH (8.0 mmol, 4 mL), and extracted with EtOAc (4 × 15 mL). The organic layers were washed with brine (15 mL) and then dried over magnesium sulfate and concentrate under vacuum (15 Torr). The residue was purified by column chromatography (silica gel, hexane/EtOAc: 20/1) to yield 87 mg of pure product **11** (81% yield). Physical and spectroscopic data follow: Colorless oil; $[\alpha]_D^{20} +75$ (*c* 1.10, CH₂Cl₂); *R_f* 0.44 (hexane/EtOAc: 16/1); IR ν (film) 3400, 2925, 2853, 1606, 1483, 743 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 0.90 (3H, t, *J* = 6.8 Hz, CH₃), 1.25-1.52 (8H, m, CH₂), 1.53-1.65 (1H, m, CH₂), 1.95 (1H, dddd, *J* = 12.7, 5.5, 4.1, 3.0 Hz, CH₂), 2.61-2.87 (2H, m, CH₂), 3.22 (1H, dtd, *J* = 9.4, 6.2, 3.0 Hz, CH), 6.46 (1H, dd, *J* = 8.4, 1.2 Hz, ArH), 6.58 (1H, td, *J* = 7.4, 1.2 Hz, ArH), 6.90-6.98 (2H, m, ArH); ¹³C NMR (75 MHz, CDCl₃) δ 14.2 (CH₃), 22.8 (CH₂), 25.5 (CH₂), 26.6 (CH₂), 28.3 (CH₂), 32.1 (CH₂), 36.8 (CH₂), 51.7 (CH), 114.2 (CH), 117.0 (CH), 121.5 (C), 126.8 (CH), 129.4 (CH), 144.9 (C); LRMS (EI) *m/z* 203 [M⁺, 22%], 133 (13), 132 (100), 130 (11), 117 (11).

(-)-Angustureine [(R)-1-Methyl-2-pentyltetrahydroquinoline (4)].- To a solution of tetrahydroquinoline **11** (36 mg, 0.18 mmol) in acetonitrile (2 mL) in a 10 mL round-bottom flask was successively added paraformaldehyde (27 mg, 0.9 mmol), NaBH₃CN (38 mg, 0.6 mmol) and acetic acid (0.034 mL, 0.6 mmol). The reaction mixture was stirred at 20 °C for 15 h, then hydrolyzed with 2M NaOH (10.0 mmol, 5 mL), and extracted with EtOAc (4 × 15 mL). The organic layers were washed with brine (15 mL), then dried over magnesium sulfate, and concentrate under vacuum (15 Torr). The residue was purified by column chromatography (silica gel, hexane/EtOAc: 20/1) to yield 32 mg of pure product **4** (82% yield). Physical and spectroscopic data follow: Colorless oil; $[\alpha]_D^{20} -12$ (*c* 1.03, CH₂Cl₂); *R_f* 0.18 (hexane); IR ν (film) 2926, 1602, 1498, 741 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 0.89 (3H, t, *J* = 9.9 Hz, CH₃), 1.20-1.45 (7H, m, CH₂), 1.53-1.64 (1H, m, CH₂), 1.83-1.92 (2H, m, CH₂), 2.64 (1H, dt, *J* = 16.2, 4.2 Hz, CH₂), 2.73-2.84 (1H, m, CH₂), 2.91 (3H, s, CH₃), 3.21 (1H, dq, *J* = 8.5, 4.2 Hz, CH), 6.51 (1H, d, *J* = 8.2 Hz, ArH), 6.57 (1H, td, *J* = 7.3, 1.0 Hz, ArH), 6.95 (1H, d, *J* = 7.3 Hz, ArH), 7.06 (1H, t, *J* = 7.7 Hz, ArH); ¹³C NMR (75 MHz, CDCl₃) δ 14.2 (CH₃), 22.8 (CH₂), 23.7 (CH₂), 24.6 (CH₂), 25.9 (CH₂), 31.4 (CH₂), 32.2 (CH₂), 38.1 (CH₃), 59.1 (CH), 110.5 (CH), 115.3 (CH), 122.0 (C), 127.2 (CH), 128.7 (CH), 145.5 (C); LRMS (EI) *m/z* 217 [M⁺ (⁸¹Br), 14%], 147 (11), 146 (100), 144 (8).

(3*S*,*R*_S)-*N*-(*tert*-Butanesulfinyl)-1-(2-bromophenyl)-5-(3,4-dimethoxyphenyl)pentan-3-amine (12).- To solution of aldimine (*R*)-**8** (632 mg, 2.0 mmol) in dry toluene (8 mL) was added dropwise a 0.5M solution of 2-(3,4-dimethoxyphenyl)ethylmagnesium bromide in THF (3.0 mmol, 6.0 mL) at -78 °C. The reaction was allowed to reach room temperature and after 4 h, it was cooled down to 0 °C and hydrolyzed

with water (5 mL), and extracted with EtOAc (4 × 15 mL). The organic layers were successively washed with water (15 mL), brine (10 mL) and then dried over magnesium sulfate and concentrate under vacuum (15 Torr). The residue was purified by column chromatography (silica gel, hexane/EtOAc: 1/1) to yield 452 mg of pure product **12** (47% yield, 86:14 dr). Physical and spectroscopic data follow: Yellow oil; $[\alpha]_D^{20}$ -30 (*c* 0.99, CH₂Cl₂); *R*_f 0.27 (hexane/EtOAc: 1/2); IR ν (film) 3262, 1515, 1261, 1235, 1024, 734 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 1.24 [9H, s, (CH₃)₃], 1.71-1.85 (2H, m, CH₂), 1.85-2.07 (3H, m, CH₂), 2.65-2.76 (2H, m, CH₂), 2.83-2.95 (1H, m, CH₂), 3.14 (1H, br d, *J* = 7.1 Hz, NH), 3.29-3.38 (1H, m, CH), 3.85 (3H, s, CH₃), 3.88 (3H, s, CH₃), 6.69-6.80 (3H, m, ArH), 7.03-7.09 (1H, m, ArH), 7.17-7.26 (2H, m, ArH), 7.52 (1H, dd, *J* = 8.0, 1.1 Hz, ArH); ¹³C NMR (75 MHz, CDCl₃) δ 22.9 (CH₃), 31.6 (CH₂), 32.6 (CH₂), 36.2 (CH₂), 38.2 (CH₂), 56.00 (CH₃), 56.04 (CH₃), 56.08 (C), 56.4 (CH), 111.4 (CH), 112.0 (CH), 120.4 (CH), 124.4 (C), 127.7 (CH), 127.9 (CH), 130.4 (CH), 133.0 (CH), 134.1 (C), 141.3 (C), 147.4 (C), 149.0 (C); LRMS (EI) *m/z* 425 [*M*⁺ - 58 (⁸¹Br), 33%], 423 (32), 376 (16), 374 (16), 261 (13), 259 (13), 197 (15), 195 (15), 192 (16), 171 (22), 169 (22), 165 (11), 164 (32), 152 (11), 151 (100), 132 (11), 117 (13), 107 (10), 91 (11), 77 (12).

(3S)-1-(2-Bromophenyl)-5-(3,4-dimethoxyphenyl)pentan-3-amine (13).- To solution of sulfinamide **12** (177 mg, 0.367 mmol) in THF (1.5 mL) was added dropwise a 6M solution of HCl (0.6 mmol, 0.10 mL) at 0 °C. The reaction mixture was stirred at 20 °C for 1 h. Then it was cooled down to 0 °C and basified with 2M NaOH (1.0 mmol, 0.5 mL). The reaction mixture was extracted with EtOAc (3 × 15 mL), dried over magnesium sulfate, filtered through a short pad of Celite, and concentrated under vacuum (15 Torr) to give 137 mg of pure product **13** (99%). Physical and spectroscopic data follow: Yellow oil; $[\alpha]_D^{20}$ +3.1 (*c* 1.09, CH₂Cl₂); *R*_f 0.38 (CH₂Cl₂/methanol: 9/1); IR ν (film) 2930, 1514, 1259, 1234, 1023, 748 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 1.54-1.70 (2H, m, CH₂), 1.70-1.92 (4H, m, CH₂), 2.60 (1H, ddd, *J* = 13.8, 10.0, 6.1 Hz, CH₂), 2.66-2.91 (4H, m, CH₂, NH₂), 3.85 (3H, s, CH₃), 3.86 (3H, s, CH₃), 6.71-6.75 (2H, m, ArH), 6.76-6.81 (1H, m, ArH), 7.00-7.08 (1H, m, ArH), 7.19-7.24 (2H, m, ArH), 7.52 (1H, d, *J* = 7.8 Hz, ArH); ¹³C NMR (75 MHz, CDCl₃) δ 32.2 (CH₂), 32.9 (CH₂), 38.3 (CH₂), 40.0 (CH₂), 50.8 (CH), 55.9 (CH₃), 56.0 (CH₃), 111.4 (CH), 111.8 (CH), 120.2 (CH), 124.4 (C), 127.6 (CH), 127.7 (CH), 130.4 (CH), 132.9 (CH), 134.9 (C), 141.6 (C), 147.3 (C), 148.9 (C); LRMS (EI) *m/z* 379 [*M*⁺ (⁸¹Br), 2%], 377 (2), 362 (20), 360 (19), 298 (41), 214 (25), 212 (26), 177 (22), 171 (19), 169 (20), 152 (33), 151 (100).

(S)-2-(3,4-Dimethoxyphenethyl)-1,2,3,4-tetrahydroquinoline (14).- To a solution of amine **13** (124 mg, 0.328 mmol) in dry toluene (5 mL) in a high pressure flask was successively added triphenylphosphine (19 mg, 0.073 mmol), Pd(OAc)₂ (5.4 mg, 0.024 mmol) and Cs₂CO₃ (214 mg, 0.656 mmol). The reaction mixture was stirred at 115 °C for 20 h. The mixture was left cooling to room temperature and hydrolyzed,

first with 3M HCl (3.0 mmol, 1 mL) and then with 2M NaOH (8.0 mmol, 4 mL), and extracted with EtOAc (4 × 15 mL). The organic layers were successively washed with brine (15 mL) and then dried over magnesium sulfate and concentrate under vacuum (15 Torr). The residue was purified by column chromatography (silica gel, hexane/EtOAc: 20/1) to yield 44 mg of pure product **14** (45% yield). Physical and spectroscopic data follow: Yellow oil; $[\alpha]_{\text{D}}^{20}$ -37 (*c* 1.00, CH₂Cl₂); *R*_f 0.41 (hexane/EtOAc: 3/1); IR ν (film) 3392, 2932, 1605, 1514, 1140, 1027, 745 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 1.59-1.74 (1H, m, CH₂), 1.76-1.86 (2H, m, CH₂), 1.99 (1H, dddd, *J* = 8.5, 5.5, 4.7, 2.4 Hz, CH₂), 2.62-2.89 (4H, m, CH₂), 3.30 (1H, dtd, *J* = 9.3, 6.3, 3.0 Hz, CH), 3.85 (3H, s, CH₃), 3.87 (3H, s, CH₃), 6.44 (1H, dd, *J* = 8.4, 1.1 Hz, ArH), 6.60 (1H, td, *J* = 7.4, 1.1 Hz, ArH), 6.71-6.83 (3H, m, ArH), 6.92-6.99 (2H, m, ArH); ¹³C NMR (75 MHz, CDCl₃) δ 26.3 (CH₂), 28.1 (CH₂), 31.9 (CH₂), 38.5 (CH₂), 51.3 (CH), 56.0 (CH₃), 56.1 (CH₃), 111.4 (CH), 111.7 (CH), 114.2 (CH), 117.1 (CH), 120.2 (CH), 121.4 (C), 126.8 (CH), 129.3 (CH), 134.6 (C), 144.6 (C), 147.4 (C), 149.0 (C); LRMS (EI) *m/z* 297 (M⁺, 38%), 298 (8), 152 (7), 133 (11), 132 (100), 117 (8).

(-)-Cuspareine [(S)-2-(3,4-Dimethoxyphenethyl)-1-methyl-1,2,3,4-tetrahydroquinoline (5)].- To a solution of tetrahydroquinoline **14** (30 mg, 0.10 mmol) in acetonitrile (1.2 mL) in a 10 mL round-bottom flask was successively added paraformaldehyde (27 mg, 0.9 mmol), NaBH₃CN (38 mg, 0.6 mmol) and acetic acid (0.034 mL, 0.6 mmol). The reaction mixture was stirred at 20 °C for 15 h, then hydrolyzed with 2M NaOH (10.0 mmol, 5 mL) and extracted with EtOAc (4 × 15 mL). The organic layers were washed with brine (15 mL) and then dried over magnesium sulfate and concentrate under vacuum (15 Torr). The residue was purified by column chromatography (silica gel, hexane/EtOAc: 20/1) to yield 25 mg of pure product **5** (80% yield). Physical and spectroscopic data follow: Colorless oil; $[\alpha]_{\text{D}}^{20}$ -19 (*c* 1.05, CH₂Cl₂); *R*_f 0.50 (hexane/EtOAc: 3/1); IR ν (film) 2932, 1498, 1260, 1233, 1028, 742 cm⁻¹; ¹H NMR (300 MHz, CDCl₃) δ 1.67-1.79 (1H, m, CH₂), 1.86-2.01 (3H, m, CH₂), 2.53 (ddd, *J* = 13.9, 10.1, 6.5 Hz, CH₂), 2.62-2.75 (2H, m, CH₂), 2.79-2.90 (1H, m, CH₂), 2.91 (3H, s, CH₃), 3.28 (1H, td, *J* = 8.4, 4.2 Hz, CH), 3.85 (3H, s, CH₃), 3.86 (3H, s, CH₃), 6.53 (1H, d, *J* = 8.2 Hz, ArH), 6.56-6.62 (1H, m, ArH), 6.69-6.75 (2H, m, ArH), 6.79 (1H, d, *J* = 8.0 Hz, ArH), 6.98 (1H, d, *J* = 7.3 Hz, ArH), 7.08 (1H, t, *J* = 7.7 Hz, ArH); ¹³C NMR (75 MHz, CDCl₃) δ 23.7 (CH₂), 24.5 (CH₂), 32.1 (CH₂), 33.2 (CH₂), 38.2 (CH₃), 55.99 (CH₃), 56.05 (CH₃), 58.5 (CH), 110.7 (CH), 111.4 (CH), 111.7 (CH), 115.5 (CH), 120.2 (CH), 121.8 (C), 127.2 (CH), 128.8 (CH), 134.8 (C), 145.4 (C), 147.3 (C), 149.0 (C); LRMS (EI) *m/z* 311 (M⁺, 25%), 147 (11), 146 (100).

ACKNOWLEDGEMENTS

This work was generously supported by the Spanish Ministerio de Ciencia e Innovación (Grant Nos. CTQ2007-65218 and CTQ2011-24165, and Consolider Ingenio 2010-CSD-2007-00006), the Generalitat Valenciana (Grant No. PROMETEO/2009/039 and FEDER) and the University of Alicante. J.A.S. thanks the Spanish Ministerio de Ciencia e Innovación for a predoctoral fellowship. We also thank MEDALCHEMY S.L. for a gift of chemicals.

REFERENCES

1. G. O. Kokwaro and G. Taylor, *Drug Chem. Toxicol.*, 1990, **13**, 347.
2. S. Omura and A. Nakagawa, *Tetrahedron Lett.*, 1981, **22**, 2199.
3. P. D. Leeson, R. W. Carling, K. W. Moore, A. M. Moseley, J. D. Smith, G. Stevenson, T. Chan, R. Baker, A. C. Foster, S. Grimwood, J. A. Kemp, G. R. Marshall, and K. Hoogsteen, *J. Med. Chem.*, 1992, **35**, 1954.
4. (a) I. Jacquemond-Collet, S. Hannedouche, N. Fabre, I. Fourasté, and C. Moulis, *Phytochemistry*, 1999, **51**, 1167; (b) I. Jacquemond-Collet, F. Benoit-Vical, M. A. Valentin, E. Stanislas, M. Mallié, and I. Fourasté, *Planta Med.*, 2002, **68**, 68.
5. P. J. Houghton, T. Z. Woldemariam, T. Watanabe, and M. Yates, *Planta Med.*, 1999, **65**, 250.
6. (a) W.-B. Wang, S.-M. Lu, P.-Y. Yang, X.-W. Han, and Y.-G. Zhou, *J. Am. Chem. Soc.*, 2003, **125**, 10536; (b) S.-M. Lu, Y.-Q. Wang, X.-W. Han, and Y.-G. Zhou, *Angew. Chem. Int. Ed.*, 2006, **45**, 2260; (c) W.-J. Tang, S.-F. Zhu, L.-J. Xu, Q.-L. Zhou, Q.-H. Fan, H.-F. Zhou, K. Lam, and A. S. C. Chan, *Chem. Commun.*, 2007, 613; (d) Z.-J. Wang, G.-J. Deng, Y. Li, Y.-M. He, W.-J. Tang, and Q.-H. Fan, *Org. Lett.*, 2007, **9**, 1243; (e) C. Deport, M. Buchotte, K. Abecassis, H. Tadaoka, T. Ayad, T. Ohshima, J.-P. Genet, K. Mashima, and V. Ratovelomanana-Vidal, *Synlett*, 2007, 2743; (f) O. Hara, T. Koshizawa, K. Makino, I. Kunimune, A. Namiki, and Y. Hamada, *Tetrahedron*, 2007, **63**, 6170; (g) D.-W. Wang, W. Zeng, and Y.-G. Zhou, *Tetrahedron: Asymmetry*, 2007, **18**, 1103; (h) M. Jahjah, M. Alame, S. Pellet-Rostaing, and M. Lemaire, *Tetrahedron: Asymmetry*, 2007, **18**, 2305; (i) X.-B. Wang, and Y.-G. Zhou, *J. Org. Chem.*, 2008, **73**, 5640; (j) F.-R. Gou, W. Li, X. Zhang, and Y.-M. Liang, *Adv. Synth. Catal.*, 2010, **352**, 2441; (k) T. Wang, L.-G. Zhuo, Z. Li, F. Chen, Z. Ding, Y. He, Q.-H. Fan, J. Xiang, Z.-X. Yu, and A. S. C. Chan, *J. Am. Chem. Soc.*, 2011, **133**, 9878.
7. G. Satyanarayana, D. Pflästerer, and G. Helmchen, *Eur. J. Org. Chem.*, 2011, 6877.
8. (a) F. Foubelo and M. Yus, *Tetrahedron: Asymmetry*, 2004, **15**, 3823; (b) J. C. González-Gómez, F. Foubelo, and M. Yus, *Synlett*, 2008, 2777; (c) M. Medjahdi, J. C. González-Gómez, F. Foubelo, and

- M. Yus, *Heterocycles*, 2008, **76**, 569; (d) M. Medjahdi, J. C. González-Gómez, F. Foubelo, and M. Yus, *J. Org. Chem.*, 2009, **74**, 7859; (e) H. K. Dema, F. Foubelo, and M. Yus, *Heterocycles*, 2010, **80**, 125; (f) J. C. González-Gómez, M. Medjahdi, F. Foubelo, and M. Yus, *J. Org. Chem.*, 2010, **75**, 6308; (g) H. K. Dema, F. Foubelo, and M. Yus, *Heterocycles*, 2011, **82**, 1411; (h) M. Medjahdi, J. C. González-Gómez, F. Foubelo, and M. Yus, *Eur. J. Org. Chem.*, 2011, 2230; (i) H. K. Dema, F. Foubelo, and M. Yus, *Jordan J. Chem.*, 2011, **6**, 247; (j) J. C. González-Gómez, F. Foubelo, and M. Yus, *Org. Synth.*, 2012, **89**, 88; (k) I. Bosque, J. C. González-Gómez, F. Foubelo, and M. Yus, *J. Org. Chem.*, 2012, **77**, 780 (correction: I. Bosque, J. C. González-Gómez, F. Foubelo, and M. Yus, *J. Org. Chem.*, 2012, **77**, 4190); (l) I. Bosque, J. C. González-Gómez, A. Guijarro, F. Foubelo, and M. Yus, *J. Org. Chem.*, 2012, **77**, 10340; (m) J. A. Sirvent, F. Foubelo, and M. Yus, *Chem. Commun.* 2012, **48**, 2543; (n) H. K. Dema, F. Foubelo, and M. Yus, *Helv. Chim. Acta*, 2012, **95**, 1790; (o) M. Medjahdi, J. C. González-Gómez, F. Foubelo, and M. Yus, *Heterocycles*, 2012, **86**, 727; (q) M. J. García-Muñoz, F. Foubelo, and M. Yus, *Eur. J. Org. Chem.*, 2013, 1287; (r) J. A. Sirvent, F. Foubelo, and M. Yus, *Eur. J. Org. Chem.*, 2013, 2461.
9. For reviews, see: (a) G.-Q. Lin, M.-H. Xu, Y.-W. Zhong, and X.-W. Sun, *Acc. Chem. Res.*, 2008, **41**, 831; (b) F. Ferreira, C. Botuha, F. Chemla, and A. Pérez-Luna, *Chem. Soc. Rev.*, 2009, **38**, 1162; (c) M. A. T. Robak, M. A. Herbage, and J. A. Ellman, *Chem. Rev.*, 2010, **110**, 3600.
10. (a) D. J. Weix and J. A. Ellman, *Org. Lett.*, 2003, **5**, 1317; (b) D. J. Weix and J. A. Ellman, *Org. Synth.*, 2005, **82**, 157.
11. (a) M. Wakayama and J. A. Ellman, *J. Org. Chem.*, 2009, **74**, 2646; (b) V. K. Aggarwal, N. Barbero, E. M. McGarrigle, G. Mickle, R. Navas, J. R. Suárez, M. G. Unthank, and M. Yar, *Tetrahedron Lett.*, 2009, **50**, 3482.
12. G. S. Lemen and J. P. Wolfe, *Org. Lett.*, 2011, **13**, 3218.
13. G. Liu, D. A. Cogan, and J. A. Ellman, *J. Am. Chem. Soc.*, 1997, **119**, 9913.
14. D. A. Cogan, G. Liu, and J. Ellman, *Tetrahedron*, 1999, **55**, 8883.
15. For similar *N*-arylation examples, see: (a) M. Liu, J. Li, X. Xiao, Y. Xie, and Y. Shi, *Chem. Commun.*, 2013, **49**, 1404; (b) T. Guo, R. Song, B.-H. Yuan, X.-Y. Chen, X.-W. Sun, and G.-Q. Lin, *Chem. Commun.*, 2013, **49**, 5402.

Please attach the original file
(Chemdraw, Chemwindows, ISIS Draw, etc)

**Enantioselective Synthesis of Tetrahydroquinolin
Alkaloids (-)-Angustureine and (-)-Cuspareine from
Chiral *tert*-Butanesulfinyl Imines**
Juan A. Sirvent, Francisco Foubelo,* and Miguel
Yus*

