Home

Search Collections Journals About Contact us My IOPscience

Energy deposition around swift proton and carbon ion tracks in biomaterials

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2017 J. Phys.: Conf. Ser. 875 112006

(http://iopscience.iop.org/1742-6596/875/12/112006)

View the table of contents for this issue, or go to the journal homepage for more

Download details:

IP Address: 193.145.230.254 This content was downloaded on 29/08/2017 at 08:32

Please note that terms and conditions apply.

You may also be interested in:

Biomaterials and therapeutic applications Angelo Ferraro

Optical research of biomaterials of Sorbulak O.V Esyrev, A A Kupchishin, A I Kupchishin et al.

A Monte Carlo track structure simulation code for the full-slowing-down carbon projectiles of energies 1 keV u-1--10 MeV u--1--in water T Liamsuwan and H Nikjoo

Study of energy deposition and stripper temperature for CYCIAE-100 Wei Su-Min, Zhang Tian-Jue, An Shi-Zhong et al.

Single electron ionization of ammonia and methane molecules by swift proton impact C A Tachino, J M Monti, O A Fojón et al.

Neutron doses due to beam losses in a novel concept of a proton therapy gantry V Talanov, D C Kiselev, D Meer et al.

Influence of surface electronic states on electron emission from metallic targets C D Archubi, M S Gravielle and V M Silkin

Monte Carlo Simulation on Energy Deposition of Low-EnergyElectrons in Liquid Water Tan Zhen-Yu, Xia Yue-Yuan, Zhao Ming-Wen et al. Numerical Simulation of Radial and Angular Distribution of -Ray's Energy Deposition in Scintillation Optical Fibre Tang Shibiao, Yin Zejie, Tang Yu et al.

Energy deposition around swift proton and carbon ion tracks in biomaterials

Rafael Garcia-Molina *¹, Maurizio Dapor ^{†2}, Pablo de Vera ^{‡3}, Isabel Abril ^{§4}

* Departamento de Física - CIOyN, Universidad de Murcia, E-30100 Murcia, Spain. [†] European Centre for Theoretical Studies in Nuclear Physics and Related Areas (ECT*), Bruno Kessler Foundation, and Trento Institute for Fundamental Physics and Applications (INFN-TIFPA), I-38123 Trento, Italy. [‡] School of Mathematics and Physics, Queen's University Belfast, BT7 1NN Belfast, Northern Ireland, UK. [§] Departament de Física Aplicada, Universitat d'Alacant, E-03080 Alacant, Spain

Synopsis Hadron therapy is a modern cancer treatment based on the interaction of proton or heavier ion beams with living tissue, whose purpose is the destruction of the malignant tumor cells producing minimal effects on the surrounding healthy tissue. To study the physical basis of the Relative Biological Effectiveness (RBE) of different projectiles, such as protons or carbon ions, we calculate the radial distribution of energy deposited by the secondary electrons generated in biomaterials by these ions at characteristic energies around the Bragg peak. This is done by means of the simulation code SEED, which follows in detail the motion and interactions of the secondary electrons as well as the subsequent electron cascade.

The interaction of energetic ions with biological targets lies in the basis of hadron therapy, which is a promising technique for oncological treatment due to its higher precision in energy delivery to the tumor cells, whereas sparing most of the healthy tissues surrounding the sensitive region [1].

The use of energetic ion beams for cancer treatment, suggested more than sixty years ago and being widely exploited nowadays, is based on the sharp and well-located dose maximum (the so-called Bragg peak) appearing at the end of their depth-dose curve. The advantages of ion beams as compared to other radiations lie mainly on their higher relative biological effectiveness (RBE), as well as in the precision with which the tumor can be irradiated, with minor side effects to the surrounding healthy regions.

A precise knowledge of the energy loss by swift ions in targets of biological interest (mainly liquid water, but also other constituents of living tissues, such as cell components...) is mandatory to suitably model the depth-dose profiles. This stage in the energy delivery by the ion beam to the target is successfully described by the SEICS code [2].

There are other important issues concerning the biodamage induced by ion bombardment. Among other questions, it is important to know the spatial distribution of the energy deposited around the ion's track by the secondary electrons generated by the projectile [3,4], as well as the subsequent electron cascade, which can propagate appreciable amounts of energy relatively far away from the ion's path.

Given the importance of low energy electrons in producing biodamage [5], it is essential to know in detail the spatial and energy distribution of electrons around the tracks of different types of projectiles (proton or carbon ions), in relevant biological targets for representative energies around the Bragg peak.

For this purpose, we use the SEED code [6], which provides a detailed description of the elastic and inelastic processes in condensed phase materials. The whole cascade of secondary electrons is followed, because any truncation or cut off underestimates the secondary electron emission yield. Also, at very low electron energies, trapping phenomena due to electron-polaron interactions and electronphonon interactions are considered.

By analyzing the radial distribution of energy deposited by the generated secondary electrons around the tracks of these projectiles in liquid water and other targets of biological interests, we hope to assess the physical basis of the higher RBE of carbon ion beams as compared to proton beams.

References

[1] M. Durante, J. S. Loeffler 2010 Nature Rev. Clin. Oncol. 7 37

[2] R. Garcia-Molina et al. 2011 Phys. Med. Biol. 56 6475

[3] P. de Vera et al. 2013 Phys. Rev. Lett. 110 148104

[4] P. de Vera et al. 2015 Phys. Rev. Lett. 114 018101

[5] B. Boudaïffa et al. 2000 Science 287 1658

[6] M. Dapor 2017 Transport of Energetic Electrons in Solids: Computer Simulation with Applications to Materials Analysis and Characterization, 2nd ed., Springer Tracts in Modern Physics, Springer



Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI. Published under licence by IOP Publishing Ltd

¹E-mail: rgm@um.es

²E-mail: dapor@ectstar.eu

³E-mail: p.devera@qub.ac.uk

⁴E-mail: ias@ua.es