

Salivary Immunoglobulin A responses to 6-minute walk test in elderly women

IVÁN RENTERÍA¹, PATRICIA CONCEPCIÓN GARCÍA SUAREZ¹, ERMILO CANTÓN MARTÍNEZ¹, PETER GRANDJEAN², ALBERTO JIMÉNEZ MALDONADO¹ 

¹Faculty of Sports, Autonomous University of Baja California, Campus Ensenada, Mexico

²Department of Health, Human Performance, and Recreation, Baylor University, United States of America

ABSTRACT

Objective: The aim of the present study was evaluate the effect of the 6-minute walk test (6MWT) on the secretory immunoglobulin A (slgA) kinetics in whole saliva of healthy elderly women. **Methods:** Nine healthy, elderly females (age 61 ± 6.7 years) performed the 6 MWT. Saliva samples and heart rate (HR) were measured in basal, immediately (IPE), and 30 minutes after the 6 MWT (30-Post Ex). Blood lactate [La] was measured at the same intervals. The ELISA assay was used to determine the slgA concentrations [slgA]. The Biuret assay was employed to determine the total proteins levels in saliva. **Results:** The exercise intensity reached by the participants was $\approx 84 \pm 16\%$ of maximum HR. Baseline [La] was 1.8 ± 0.9 mmol/L. [La] at IPE increased 3.4 mmol/L above baseline ($p = 0.02$). [slgA] increased by $233.3 \pm 109.3 \%$ at IPE and remained $211.2 \pm 100.1 \%$ greater at 30-Post Ex compared with basal ($p = 0.02$). The 6 MWT did not modify total proteins levels ($p > 0.05$). **Conclusion:** The 6 MWT, performed to submaximal intensity, appears to be enough of a stimulus to increases slgA levels in healthy elderly women independent of total proteins concentration in saliva. **Keywords:** 6-Minute walk test, Blood lactate, Elderly women, Secretory immunoglobulin A.

Cite this article as:

Rentería, I., García, P.C., Cantón, E., Grandjean, P., & Jiménez, A. (2019). Salivary Immunoglobulin A responses to 6-minute walk test in elderly women. *Journal of Human Sport and Exercise*, 14(1), 225-235. doi:<https://doi.org/10.14198/jhse.2019.141.19>

 **Corresponding author.** Facultad de Deportes Campus Ensenada, Universidad Autónoma de Baja California. Boulevard Zertuche s/n. Fraccionamiento Valle Dorado. Ensenada, Baja California. 22890. México. <https://orcid.org/0000-0003-4361-2560>

E-mail: jimenez.alberto86@uabc.edu.mx

Submitted for publication May 2018

Accepted for publication October 2018

Published *in press* October 2018

JOURNAL OF HUMAN SPORT & EXERCISE ISSN 1988-5202

© Faculty of Education. University of Alicante

doi:10.14198/jhse.2019.141.19

INTRODUCTION

The older population has become the main growth segment in the current society (Chang, Pan, Chen, Tsai, & Huang, 2012), and is projected to accelerate in the coming decades (WHO, 2015). It has been estimated that by the 2030, the people in the world aged over 60 years will be 1.4 billion, and by the 2050, the older population will reach almost 2.1 billion (WHO, 2015). The aging is a time-dependent functional decline process that affects most living organisms (Carmona & Michan, 2016; López-Otín, Blasco, Partridge, Serrano, & Kroemer, 2013). In agreement with this, the elderly people show a high risk to suffer cognitive problems (Uemura et al., 2012), development of cardiovascular diseases (de Souza et al., 2015; Shanmugasundaram, Rough, & Alpert, 2010) and immunosenescence (Prcina, Novak, Cigankova, & Kontsekova, 2018). The last condition is caused by changes in the functions in both adaptive and innate immune system (van der Geest et al., 2017).

Secretory IgA (sIgA) is a predominant immunoglobulin on the mucosal surface of humans and other mammals (Corthésy, 2013). In saliva, the plasma cells adjacent to ducts and acini of salivary glands are the principal source of sIgA (Bishop, Walker, Scanlon, Richards, & Rogers, 2006). Previously, authors indicated that people with salivary sIgA deficiency are susceptible to recurrent infections, principally at the upper respiratory tract (Teixeira, Martins, Martins, & Cunha, 2008). On the other hand, studies performed in young adults, demonstrated that the acute (single bout) exercise increases sIgA concentration [sIgA] (Allgrove, Gomes, Hough, & Gleeson, 2008; Sari-Sarraf, Reilly, Doran, & Atkinson, 2007), reduces [sIgA] (Baralic et al., 2010; Usui et al., 2011) or does not change [sIgA] (Li & Rush, 2009). The controversies could stem from differences in fitness levels, exercise modes, intensities and fidelity of sample collection.

The acute sIgA responses to single episodes of physical exercise in older individuals are largely unexplored (Rikli & Jones, 1998). In detail the sIgA secretion was elevated after just twenty minutes of low-intensity calisthenics in older women (Sakamoto et al., 2005). On the other hand, Hwang et al (Hwang, Park, & Lim, 2016) reported that following 3-months of Pilates training, the sIgA secretion rate increased transiently, but the authors did not find any effect on [sIgA] with a high-intensity cycle ergometry test (i.e., Åstrand Cycle Ergometry Test) in older women. The Åstrand Cycle Ergometry Test is a protocol employed to determine the VO_2Max (Åstrand & Ryhming, 1954). However, this test utilize expensive equipment. The exercise technicians or physicians, usually employ more accessible but validated test to determine the fitness level of the patients.

The 6-minute walk test (6 MWT) is an inexpensive and easy test commonly employed to determine the physical endurance in healthy people (Negreiros et al., 2017; Rikli & Jones, 1998; Someya, Mugii, & Oohata, 2015). Initially, Rikli & Jones indicated that during the 6MWT, subjects walked by 6 minutes as fast as they comfortably could, trying to cover the maximum distance possible (Rikli & Jones, 1998). However, others studies identified that if the tester, encourage verbally the participants, they get a better performance during the 6 MWT (Guimarães, Bellotti, Bacal, Mocelin, & Bocchi, 2002; Troosters et al., 2002). Concretely, with the encourage method, the participants walk almost to their maximum capacity (Guimarães et al., 2002; Guyatt et al., 1985). Consequently, the oxygen uptake is higher (Troosters et al., 2002). Regarding to these information, it is possible consider to the 6 MWT performed with the encouragement as a vigorous exercise. It is know that the high intensity exercise induces immunosuppression (Ihalainen, Schumann, Mero, & Ihalainen, 2016; Nieman, 1994). Particularly, the high intensity training reduces the [sIgA] in saliva (Ihalainen et al., 2016), increasing the risk to acquire recurrent upper respiratory tract infection (URTI) (Gleeson & Pyne, 2015; Teixeira et al., 2008). Thus, the 6 MWT performed with the encouragement method can result in a paradox situation. On one side, allows identify more accurately the fitness levels in the elderly population.

However, on the other side, could induce immunosuppression and make them prone to URTIs. Therefore, the aim of the present work was study the effect of the 6 MWT performed with the encouragement method on the salivary [sIgA] in healthy elderly women.

MATERIAL AND METHODS

The study was approved by the ethics committee of the Universidad Autónoma de Baja California, México. All experimental procedures were conducted in accordance with Helsinki Declaration.

Participants

Nine healthy elderly women (61 ± 6.7 years) of modest physical fitness living in Ensenada (Baja California, Mexico) were recruited. Participants were informed of the scope and procedure of the study and subsequently provided written consent. Participants completed the physical activity scale for elderly (PASE) to determine the physical activity in elderly people (Voorrips, Ravelli, Dongelmans, Deurenberg, & van Staveren, 1991). According to the PASE score, all participants could be classified as having moderate physical activity levels (Voorrips et al., 1991).

Study Design

The 6 MWT was performed after a 2-hr fasting period and between 9 and 11:00 AM. The blood lactate and heart rate (HR) were determined at baseline (just before exercise), immediately (IPE), and 30- minutes post-exercise (30 Post-Ex). Saliva and blood samples were collected with HR at each time point.

Anthropometry Measurements

Anthropometrics variables, such as body weight, height, and waist circumference (WC) were determined in all the subjects 24 hrs. after participants completed the PASE questionnaire. An electronic balance (RICE LAKE, Floor Level) was employed to determine the body weight. A stadiometer (Seca 213) was used to evaluate height. WC was determined following methodology previously reported (Howel, 2012). Body mass index (BMI) was calculated as body weight in kilograms divided by the square of body height in meters. All anthropometric measurements were performed between 10:00 AM and 12:00 PM, during which participants underwent the pre-exercise 2-hr fasting period.

6-minute Walk Test

The 6 MWT test, useful for estimating physical endurance in elderly people (Rikli & Jones, 1998), was performed following methodology previously reported (Guimarães et al., 2002; Rikli & Jones, 1998) with some modifications. In brief, one TANITA pedometer (PD-733F; Illinois: USA) was assigned to each participant. The pedometer was attached with a strap to the neck, and setting to display the steps. The pedometer was cleared of previous registers. The investigators made a mark on the ground, then, the participants performed 3 steps back from the mark. After this, the investigator requested participants to walk 10 steps from the mark previously made. One investigator evaluated the distance covered by 10 steps, with this, the stride length was determined. The pedometer was setting to zero. The participants were instructed to walk around a basketball field for 6 minutes, the aim was covered as much ground as possible. One tester walked alongside the participants encouraging to walk as fast as possible. This methodology encouraged the greatest effort in participants (Guimarães et al., 2002). Immediately after the 6 MWT, the tester registered the steps walked by the participant to determine the distance covered.

Cardiovascular and Metabolic Variables

Heart rate (HR) and blood pressure (BP) were measured after 5 minutes of seated rest and prior to completing the 6 MWT. BP was measured using the auscultatory method. HR was determined using the Polar FT1 heart rate monitor at the end of the 5-minute rest period and throughout exercise and recovery. The exercise intensity was determined with following formula: heart rate at 6 min/estimated maximal heart rate (220-age) (Someya et al., 2015). A hand-held lactate analyzer (Nova Biomedical) and test strips were employed to evaluate the lactate concentrations.

Saliva Collection

An unstimulated saliva sample was collected in each participant at the times indicated above. The saliva collection at baseline and 30 Post-Ex were performed following methodology previously reported (Akimoto et al., 2003; Baralic et al., 2010; Shimizu et al., 2007) with brief modifications. Participants sat quietly for 5 minutes, and then rinsed out their mouths with sterilized water three times. Participants remained in seated rest for another 5 minutes with slightly lowered head to allow the spontaneous saliva flow in the mouth. The participants expectorated three times for 2 minutes specifically at 0, 60 and 120 seconds into a falcon tube of 15 mL. During the saliva collection process, the participants were asked not to make any special effort to gather saliva in their mouths. But, they were asked to allow the saliva to flow naturally. Saliva collected IPE followed similar methodology, but without the prior rest period.

The saliva samples were immediately stored to -20°C for future analysis. Saliva flow was determined following previously-published methodology (Sakamoto et al., 2005; Teixeira et al., 2008). In brief, the total sample volume (mL) was divide by the time taken to produce the saliva sample (minutes) and reported as mL/min.

Saliva Analysis

slgA concentrations were determined using an enzyme-linked immunosorbent assay (ELISA) kit (1-1602; Salimetrics, LLC, Carlsbad, CA) according to the manufacturer's instructions. To avoid inter-assay variability, all the samples were assayed on the same microtiter plate. The samples were analyzed in duplicate and the mean concentrations were calculated. The limit of sensitivity for slgA was $2.5\ \mu\text{g/mL}$. Color was detected 10 min after the stop solution was added and the absorbance was read at 490 nm. (iMark microplate absorbance reader, BIO-RAD, Cat. 168-1130). The slgA secretion rate ($\mu\text{g/min}$) was determined by multiplying the absolute slgA concentration ($\mu\text{g/mL}$) with the saliva flow rate (SFR) (mL/min) (Baralic et al., 2010; Teixeira et al., 2008). Total proteins were determined by the Biuret protein assay (Spinreact). The bovine serum albumin was used as standard following the manufacturer's instructions.

Statistical Analysis

The data are presented as mean \pm standard deviation (SD). Groups of data were compared in a one-way, repeated-measures analysis of variance (ANOVA) followed by Tukey's post hoc test; a p value of ≤ 0.05 indicated statistical significance. All statistical analyses and figures were performed using the statistical software package GraphPad Prism 6.0 (CA, USA).


RESULTS

Participant baseline characteristics are displayed in the Table 1. In general, the women could be described as overweight, but with normal resting hemodynamics and having moderate physical activity levels. The mean VO_2Max for the participants was $32.21 \pm 2.87\ \text{mL/kg}\cdot\text{min}$ (Table 1).

Table 1. Baseline characteristics of the study subjects. (n = 9)

Variable	Mean	Standard Deviation
Age (yrs)	61	6.7
Body weight (kg)	67.7	7.1
Height (m)	1.56	0.05
Body Mass Index (kg/m ²)	27.5	2.2
Waist circumference (cm)	86.1	6.7
Resting HR (ppm)	73.6	8.7
Resting systolic BP (mmHg)	131.7	18.7
Resting diastolic BP (mmHg)	74.1	10.5
PASE Score (physical activity score)	12.2	4.7
VO ₂ Max estimated (mL/kg•min)	32.21	2.87

The mean distance covered during the 6 MWT was 590.8 ± 55.3 m – a rate of 1.64 m/sec. The mean exercise intensity was $84.1 \pm 15.9\%$ of HR max. Baseline HR (74 ± 9 bpm) increased to 134 ± 24 bpm IPE and then returned toward baseline by 30 Post-Ex (78 ± 11 bpm) ($p < 0.0001$) (Figure 1a). The Blood [La] increased 188% from baseline (1.8 ± 0.9 mmol/L) to 5.2 ± 3.1 mmol/L IPE ($p = 0.02$). Blood [La] remained 56% above baseline at 30 Post-Ex, but this difference was not significantly different (Figure 1b). There was a moderate but significant relationship ($r=0.69$, $p=0.03$) between the HR max and [La] at the end the 6 MWT (Figure 2).



*** $p < 0.00001$ vs Before exercise. ### $p < 0.0001$ vs IPE * $p = 0.02$ vs Before exercise. One way ANOVA. Tukey's pos hoc test

Figure 1. Changes in heart rate during the 6-minute walk test (6 MWT) in healthy elderly women. The heart rate was determined at basal state (Before exercise). Immediately finished the 6 MWT (IPE), and 30 minutes after finished the 6 MWT (30-Post Exe) (a). Effects of the 6-minute walk test (6 MWT) on the blood lactate levels in healthy elderly women. Blood lactate determined at baseline (Before exercise). Immediately finished the 6 MWT (IPE), and 30 minutes after finished the 6 MWT (30-Post Exe) (b)





Figure 2. Correlation analysis of the blood lactate levels vs heart rate at finished the 6-minute walk test

The 6 MWT increased the salivary [sIgA] from $41.1 \pm 20.9 \mu\text{g/mL}$ at baseline to $89.1 \pm 39.6 \mu\text{g/mL}$ IPE ($p = 0.028$), at 30 Post-Ex, the mean [sIgA] was $78.9 \pm 41.5 \mu\text{g/mL}$ (Figure 3a). The Figure 3b shows that 6 MWT increased significantly the relative [sIgA] at both IPE and 30 Post-Ex with respect to basal state ($F(2, 21) = 5.57, p = 0.01$). Baseline SFR ($0.35 \pm 0.15 \text{ mL/min}$) remained unaltered IPE ($0.22 \pm 0.12 \text{ mL/min}$) or after exercise ($0.28 \pm 0.17 \text{ mL/min}$) ($p = 0.20$) (Figure 3c). In the same sense, the sIgA secretion rate did not change with the exercise (baseline = $13.7 \pm 8.0 \mu\text{g/min}$; IPE = $20.9 \pm 14.8 \mu\text{g/min}$; 30 Post-Ex = $19.9 \pm 16.5 \mu\text{g/min}$) (Figure 3d). Total [protein] in saliva did not change with the 6 MWT (baseline = $5.15 \pm 1.64 \text{ mg/mL}$; IPE = $7.65 \pm 2.86 \text{ mg/mL}$; 30 Post-Ex = $5.94 \pm 1.85 \text{ mg/mL}$) (Figure 4).



* $p < 0.05$ vs Before exercise. One-way ANOVA. Tukey's pos hoc test

Figure 3. Effect of the 6-minute walk test (6 MWT) on the absolute sIgA concentration in saliva (a), relative sIgA concentration in saliva (b), saliva flow rate (c), and salivary sIgA secretion rate (d). Sample determined at basal state (Before exercise). Immediately finished the 6 MWT (IPE), and 30 minutes after finished the 6 MWT (30-Post Exe)




Figure4. Effect of the 6 MWT on the total saliva protein concentration. Sample determined at basal state (Before exercise). Immediately finished the 6 MWT (IPE), and 30 minutes after finished the 6 MWT (30-Post Exe)

DISCUSSION

The present study was designed to characterize the salivary sIgA response to the 6 MWT in healthy elderly women. Results of the current study, demonstrate that [sIgA] are elevated after a 6MWT, performed at greater than 80% of maximal HR, in moderately-active older women. The higher post-exercise [sIgA] was not accompanied by changes in SFR, sIgA secretion rate or total salivary proteins. Our results may be interpreted to indicate that a brief and moderate-vigorous field exercise testing in elderly women does not induce immunosuppression at the upper respiratory tracts (indicated by [sIgA]).

The mean 6 MWT distance covered by the present cohort is similar to that previously reported in older physically active women (Janaudis-Ferreira, Sundelin, & Wadell, 2010). Furthermore, the exercise intensity reached by the participants in the current study was ~ 84%. This data is in agreement with others studies in which the encouragement method was utilized during the 6MWT (Jenkins et al., 2009). The current study demonstrates an immune-simulative effect of a single episode (6 minutes) of physical exercise to submaximal effort in older individuals with moderate physical activity level. In concordance with our data, a recent study reported a higher absolute sIgA concentration in older women after a graded exercise test performed to >80 % HR (Hwang et al., 2016). However, in that study, the length of the exercise session was not reported. Besides, in the Hwang' work just reported the absolute [sIgA]. Others authors also reported a higher absolute sIgA concentration after a single exercise session to low intensity (Sakamoto et al., 2009). However, more than absolute concentrations, authors indicated that the relative [sIgA] are preferable data to determine the risk of URTI (Neville, Gleeson, & Folland, 2008). For this reason, in the present study, we determine the impact of the 6 MWT on the relative [sIgA]. Our data indicate that the 6MWT performed to submaximal effort, increases immediately the relative [sIgA], this effect is still present after 30 minutes of finished the 6MWT. To our best knowledge, this is a novel data. Currently, the molecular mechanism to explain the higher sIgA after the exercise is unknown. Therefore, further studies are necessary to identify the underlying molecular mechanism that explain the changes in salivary [sIgA] during the physical exercise in older population. The sIgA secretion rate was not modify by the 6MWT. Other reports indicated a higher sIgA secretion rate after a single session (Hwang et al., 2016; Sakamoto et al., 2005). In these studies, a higher SFR after the acute exercise also was reported. Contrary, in the current study, the field exercise testing did not change the SFR. Considering that SFR is a primary component for increasing the sIgA secretion rate during the exercise in elderly subjects (Sloan, Engels, Fahlman, Yarandi, & Davis, 2013), We suggest that the no effect of the 6 MWT on the SFR in the participants, could explain partially, the lack effect on the sIgA secretion rate.

In addition to the immunological results, the present work found a significant relationship between the blood lactate and the maximal heart rate during the 6 MWT. Together, the [La] and HR responses indicate that the 6 MWT was performed at sub-maximal effort. Previous authors reported that the blood lactate and the HR max during the 1-Mile walk test have a direct relationship in healthy elderly adults (Bazzano, Cunningham, Cama, & Falconio, 1998). These background and the data of the current study, suggest that the metabolic variable (lactate) and heart rate are good indicators to control the intensity during the field exercise testing at least in apparently healthy elderly people.

CONCLUSION

The results of the present data showed that, the 6 MWT performed to submaximal effort increased [sIgA] in elderly healthy women. Neither changes on the SFR, sIgA secretion rates, and the unaltered concentrations of salivary total protein measured through the exercise, indicate specific effect of the field exercise testing on the immunoglobulin. The mechanisms from moderate-to-high-intensity exercise that increase [sIgA] and potentially enhance immune function remain to be determined. Additionally, it is worth indicate that the encouragement method affect the performance during the 6MWT (Guyatt et al., 1985), this situation open the possibility to study the impact of the 6 MWT performed without the encouragement on immune response in healthy elderly women.

ACKNOWLEDGMENTS

We greatly appreciate the time and effort provided by the elderly subjects. We thank members of the SNTE Ensenada, participants of the community program for older adults “Envejecimiento Activo” (Facultad de Deportes, Universidad Autónoma de Baja California Campus Ensenada) for their cooperation during the current study. We also thank our students for their skillful assistance during the 6 minute walk test.

COMPETING INTERESTS

The authors declare that they have no competing interests.

REFERENCES

- Akimoto, T., Kumai, Y., Akama, T., Hayashi, E., Murakami, H., Soma, R., ... Kono, I. (2003). Effects of 12 months of exercise training on salivary secretory IgA levels in elderly subjects. *British Journal of Sports Medicine*, 37(February 2009), 76–79. <https://doi.org/10.1136/bjism.37.1.76>
- Allgrove, J. E., Gomes, E., Hough, J., & Gleeson, M. (2008). Effects of exercise intensity on salivary antimicrobial proteins and markers of stress in active men. *Journal of Sports Sciences*, 26(6), 653–661. <https://doi.org/10.1080/02640410701716790>
- Astrand, P., & Ryhming, I. (1954). A nomogram for calculation of aerobic capacity (physical fitness) from pulse rate during sub-maximal work. *J Appl Physiol*, 7(2), 218–221. <https://doi.org/10.1152/jappl.1954.7.2.218>
- Baralic, I., Miletic, I., Djordjevic, B., Terzic, T., Radojevic-Skodric, S., & Nikolic, G. (2010). Effect of sensory stimulation on salivary IgA secretion rate in karate players. *Biology of Sport*, 27(4), 273–278. <https://doi.org/10.5604/20831862.927493>
- Bazzano, C., Cunningham, L. N., Cama, G., & Falconio, T. (1998). The relationship of lactate to 1-Mile walk performance in women aged 60 to 70 years. *Journal of Aging & Physical Activity*, 6, 285–289. <https://doi.org/10.1123/japa.6.3.285>

- Bishop, N. C., Walker, G. J., Scanlon, G. A., Richards, S., & Rogers, E. (2006). Salivary IgA responses to prolonged intensive exercise following caffeine ingestion. *Medicine and Science in Sports and Exercise*, 38(3), 513–519. <https://doi.org/10.1249/01.mss.0000187412.47477.ee>
- Carmona, J. J., & Michan, S. (2016). Biology of Healthy Aging and Longevity. *Revista de Investigacion Clinica; Organo Del Hospital de Enfermedades de La Nutricion*, 68(1), 7–16.
- Chang, Y.-K., Pan, C.-Y., Chen, F.-T., Tsai, C.-L., & Huang, C.-C. (2012). Effect of Resistance-Exercise Training on Cognitive Function in Healthy Older Adults: A Review. *Journal of Aging and Physical Activity*, 20(4), 497–517. <https://doi.org/10.1123/japa.20.4.497>
- Corthésy, B. (2013). Role of secretory IgA in infection and maintenance of homeostasis. *Autoimmunity Reviews*, 12(6), 661–665. <https://doi.org/10.1016/j.autrev.2012.10.012>
- de Souza, J. D., Ribeiro, A. Q., Martinho, K. O., Franco, F. S., Martins, M. V., Rodrigues, M. G., ... Tinôco, A. L. A. (2015). Lipid profile and associated factors among elderly people, attended at the Family Health Strategy, Viçosa/MG. *Nutricion Hospitalaria*, 32(2), 771–778. <https://doi.org/10.3305/nh.2015.32.2.8875>
- Gleeson, M., & Pyne, D. B. (2015). Respiratory inflammation and infections in high-performance athletes. *Immunology and Cell Biology*, 94(2), 124–131. <https://doi.org/10.1038/icb.2015.100>
- Guimarães, G. V., Bellotti, G., Bacal, F., Mocelin, A., & Bocchi, E. A. (2002). Can the cardiopulmonary 6-minute walk test reproduce the usual activities of patients with heart failure? *Arquivos Brasileiros de Cardiologia*, 78(6), 553–560. <https://doi.org/10.1590/S0066-782X2002000600003>
- Guyatt, G. H., Sullivan, M. J., Thompson, P. J., Fallen, E. L., Pugsley, S. O., Taylor, D. W., & Berman, L. B. (1985). The 6-minute walk: A new measure of exercise capacity in patients with chronic heart failure. *Can Med Assoc*, 132(9), 919–923. <https://doi.org/10.1016/j.physio.2011.11.003>
- Howel, D. (2012). Waist Circumference and Abdominal Obesity among Older Adults: Patterns, Prevalence and Trends. *PLoS ONE*, 7(10). <https://doi.org/10.1371/journal.pone.0048528>
- Hwang, Y., Park, J., & Lim, K. (2016). Effects of pilates exercise on salivary Secretory immunoglobulin a levels in older women. *Journal of Aging and Physical Activity*, 24(3), 399–406. <https://doi.org/10.1123/japa.2015-0005>
- Ihalainen, J. K., Schumann, M., Mero, A. A., & Ihalainen, J. K. (2016). Mucosal immunity and upper respiratory tract symptoms in recreational endurance runners. *Physiology, Applied*, 41(1), 96–102.
- Janaudis-Ferreira, T., Sundelin, G., & Wadell, K. (2010). Comparison of the 6-minute walk distance test performed on a non-motorised treadmill and in a corridor in healthy elderly subjects. *Physiotherapy*, 96(3), 234–239. <https://doi.org/10.1016/j.physio.2009.11.015>
- Jenkins, S., Cecins, N., Camarri, B., Williams, C., Thompson, P., & Eastwood, P. (2009). Regression equations to predict 6-minute walk distance in middle-aged and elderly adults. *Physiotherapy Theory and Practice*, 25(7), 516–522. <https://doi.org/10.3109/09593980802664711>
- Li, T.-L., & Rush, B. (2009). The Effects of Prolonged Strenuous Exercise on Salivary Secretion of IgA Subclasses in Men. *International Journal*, 1(3), 69–74.
- López-Otín, C., Blasco, M. A., Partridge, L., Serrano, M., & Kroemer, G. (2013). The hallmarks of aging. *Cell*, 153(6). <https://doi.org/10.1016/j.cell.2013.05.039>
- Negreiros, A., Padula, R. S., Andrea Bretas Bernardes, R., de Moraes, M. V., Pires, R. S., & Chiavegato, L. D. (2017). Predictive validity analysis of six reference equations for the 6-minute walk test in healthy Brazilian men: a cross-sectional study. *Brazilian Journal of Physical Therapy*, 21(5), 350–356. <https://doi.org/10.1016/j.bjpt.2017.06.003>
- Neville, V., Gleeson, M., & Folland, J. P. (2008). Salivary IgA as a risk factor for upper respiratory infections in elite professional athletes. *Medicine and Science in Sports and Exercise*, 40(7), 1228–1236. <https://doi.org/10.1249/MSS.0b013e31816be9c3>

- Nieman, D. C. (1994). Exercise, upper respiratory tract infections and the immune system. *Medicine Science Sport Exercise*, 26 (2), 128-139. <https://doi.org/10.1249/00005768-199402000-00002>
- Prcina, M., Novak, M., Cigankova, V., & Kontsekova, E. (2018). Immunosenescence - the role in the immunotherapy of older population. *Bratislavské Lekárske Listy*, 119(4), 217–220. https://doi.org/10.4149/BLL_2018_040
- Rikli, R. E., & Jones, C. J. (1998). The Reliability and Validity of a 6-Minute Walk Test as a Measure of Physical Endurance in Older Adults. *Journal of Aging & Physical Activity*. <https://doi.org/10.1097/00005768-199805001-00421>
- Sakamoto, Y., Ueki, S., Kasai, T., Takato, J., Shimanuki, H., Honda, H., ... Haga, H. (2009). Effect of exercise, aging and functional capacity on acute secretory immunoglobulin A response in elderly people over 75 years of age. *Geriatrics and Gerontology International*, 9(1), 81–88. <https://doi.org/10.1111/j.1447-0594.2008.00502.x>
- Sakamoto, Y., Ueki, S., Shimanuki, H., Kasai, T., Takato, J., Ozaki, H., ... Haga, H. (2005). Effects of low-intensity physical exercise on acute changes in resting saliva secretory IgA levels in the elderly. *Geriatrics and Gerontology International*, 5(3), 202–206. <https://doi.org/10.1111/j.1447-0594.2005.00297.x>
- Sari-Sarraf, V., Reilly, T., Doran, D. A., & Atkinson, G. (2007). The effects of single and repeated bouts of soccer-specific exercise on salivary IgA. *Archives of Oral Biology*, 52(6), 526–532. <https://doi.org/10.1016/j.archoralbio.2006.11.016>
- Shanmugasundaram, M., Rough, S. J., & Alpert, J. S. (2010). Dyslipidemia in the elderly: Should it be treated? *Clinical Cardiology*, 33(1), 4–9. <https://doi.org/10.1002/clc.20702>
- Shimizu, K., Kimura, F., Akimoto, T., Akama, T., Kuno, S., & Kono, I. (2007). Effect of free-living daily physical activity on salivary secretory IgA in elderly. *Medicine and Science in Sports and Exercise*, 39(4), 593–598. <https://doi.org/10.1249/mss.0b013e318031306d>
- Sloan, C. A., Engels, H. J., Fahlman, M. M., Yarandi, H. E., & Davis, J. E. (2013). Effects of exercise on S-IGA and URS in postmenopausal women. *International Journal of Sports Medicine*, 34(1), 81–86. <https://doi.org/10.1055/s-0032-1314817>
- Someya, F., Mugii, N., & Oohata, S. (2015). Cardiac hemodynamic response to the 6-minute walk test in young adults and the elderly. *BMC Res Notes*, 8, 355. <https://doi.org/10.1186/s13104-015-1331-5>
- Teixeira, A. M., Martins, R., Martins, M., & Cunha, M. D. R. (2008). Changes in Functional Fitness, Mood States and Salivary IgA Levels after Exercise Training for 19 Weeks in Elderly Subjects. *International Journal of Applied Sports Sciences*, 20(2), 16–26.
- Troosters, T., Vilaro, J., Rabinovich, R., Casas, A., Barberà, J. A., Rodriguez-Roisin, R., & Roca, J. (2002). Physiological responses to the 6-min walk test in patients with chronic obstructive pulmonary disease. *European Respiratory Journal*, 20(3), 564–569. <https://doi.org/10.1183/09031936.02.02092001>
- Uemura, K., Doi, T., Shimada, H., Makizako, H., Yoshida, D., Tsutsumimoto, K., ... Suzuki, T. (2012). Effects of Exercise Intervention on Vascular Risk Factors in Older Adults with Mild Cognitive Impairment: A Randomized Controlled Trial. *Dementia and Geriatric Cognitive Disorders Extra*, 2(1), 445–455. <https://doi.org/10.1159/000343486>
- Usui, T., Yoshikawa, T., Ueda, S., Katsura, Y., Orita, K., & Fujimoto, S. (2011). Effects of Acute Prolonged Strenuous Exercise on the Salivary Stress Markers and Inflammatory Cytokines. *Jpn J Phys Fitness Sports Med*, 60(3), 295–304. <https://doi.org/10.7600/jspfsm.60.295>
- van der Geest, K. S. M., Wang, Q., Eijsvogels, T. M. H., Koenen, H. J. P., Joosten, I., Brouwer, E., ... Boots, A. M. H. (2017). Changes in peripheral immune cell numbers and functions in octogenarian

- walkers - an acute exercise study. *Immunity and Ageing*, 14(1), 1–13.
<https://doi.org/10.1186/s12979-017-0087-2>
- Voorrips, L. E., Ravelli, A. C., Dongelmans, P. C., Deurenberg, P., & van Staveren, W. A. (1991). A physical activity questionnaire for the elderly 108. *Med.Sci Sports Exerc.*, 23(0195–9131 (Print)), 974–979. <https://doi.org/10.1249/00005768-199108000-00015>
- WHO. (2015). World population, ageing. United Nations, Department of Economic and Social Affairs, Population Division (2015). World Population Ageing, United Nat((ST/ESA/SER.A/390), 164. <https://doi.org/ST/ESA/SER.A/390>

