

Noname manuscript No.
(will be inserted by the editor)

Corrigendum: Almost periodic functions in terms of Bohr's equivalence relation

J.M. Sepulcre · T. Vidal

Received: date / Accepted: date

The authors wish to draw the attention to a mistake which appears in the proof of Proposition 3 of the above quoted paper [4]. Specifically, after label (17), the statement “we can also construct a vector $\mathbf{x}_0 \in \mathbb{R}^{\sharp G_A}$ such that, by taking into account Remarks 1 and 2, verifies

$$a_j = a_{1,j} e^{i\langle \mathbf{r}_j, \mathbf{x}_0 \rangle}, \quad j = 1, 2, \dots, \quad \text{with } \lambda_j \in \Lambda$$

is correct when it is possible to obtain an integral basis for the set of exponents or frequencies Λ , but it is not correct in the general case. We next define the concept of *integral basis*.

Let $G_A = \{g_1, g_2, \dots, g_k, \dots\}$ be a basis of the \mathbb{Q} -vector space generated by a set $\Lambda = \{\lambda_1, \lambda_2, \dots\}$ of exponents, (by abuse of notation, it is said that G_A is a basis for Λ), which implies that G_A is linearly independent over the rational numbers and each λ_j is expressible as a finite linear combination of terms of G_A , say

$$\lambda_j = \sum_{k=1}^{i_j} r_{j,k} g_k, \quad \text{for some } r_{j,k} \in \mathbb{Q}, \quad i_j \in \mathbb{N}. \quad (2)$$

We will say that G_A is an *integral basis* for Λ when $r_{j,k} \in \mathbb{Z}$ for any j, k .

By taking into account this situation, in order to maintain the validity of Proposition 3 (not only for the case when it is possible to obtain an integral basis for the set of exponents) and some other subsequent results of our paper, the equivalence relation which is inspired by that of Bohr is now revised to adapt correctly the situation in the general case. In this way, Definition 3 of the above quoted paper is now modified in the following terms.

J.M. Sepulcre · T. Vidal
Department of Mathematics
University of Alicante
03080-Alicante, Spain

Definition 3' Given $\Lambda = \{\lambda_1, \lambda_2, \dots, \lambda_j, \dots\}$ a set of exponents, consider $A_1(p)$ and $A_2(p)$ two exponential sums in the class \mathcal{S}_Λ , say $A_1(p) = \sum_{j \geq 1} a_j e^{\lambda_j p}$ and $A_2(p) = \sum_{j \geq 1} b_j e^{\lambda_j p}$. We will say that A_1 is equivalent to A_2 if for each integer value $n \geq 1$, with $n \leq \#\Lambda$, it is satisfied $a_n^* \sim b_n^*$, where $a_n^*, b_n^* : \{\lambda_1, \lambda_2, \dots, \lambda_n\} \rightarrow \mathbb{C}$ are the functions given by $a_n^*(\lambda_j) := a_j$ y $b_n^*(\lambda_j) := b_j$, $j = 1, 2, \dots, n$ and \sim is in Definition 1.

It is clear that the relation defined in the foregoing definition is an equivalence relation. That is, it is reflective, symmetric and transitive. By abuse of notation, we will use \sim for both equivalence relations introduced in definitions 1 and 3'.

Analogously, Definition 5 must be rewritten in the following terms.

Definition 5' Given $\Lambda = \{\lambda_1, \lambda_2, \dots, \lambda_j, \dots\}$ a set of exponents, let f_1 and f_2 denote two equivalence classes of $B(\mathbb{R}, \mathbb{C}) / \simeq$ (resp. $B(U, \mathbb{C}) / \simeq$) whose associated Fourier series (resp. Dirichlet series) are given by

$$\sum_{j \geq 1} a_j e^{i\lambda_j t} \text{ and } \sum_{j \geq 1} b_j e^{i\lambda_j t}, \quad a_j, b_j \in \mathbb{C}, \quad \lambda_j \in \Lambda.$$

$$\text{(resp. } \sum_{j \geq 1} a_j e^{\lambda_j s} \text{ and } \sum_{j \geq 1} b_j e^{\lambda_j s}, \quad a_j, b_j \in \mathbb{C}, \quad \lambda_j \in \Lambda.$$

We will say that f_1 is equivalent to f_2 if for each integer value $n \geq 1$, with $n \leq \#\Lambda$, it is satisfied $a_n^* \sim b_n^*$, where $a_n^*, b_n^* : \{\lambda_1, \dots, \lambda_n\} \rightarrow \mathbb{C}$ are the functions given by $a_n^*(\lambda_j) := a_j$ and $b_n^*(\lambda_j) := b_j$, $j = 1, 2, \dots, n$, and \sim is in Definition 1.

Under the modification of Definition 3, Proposition 1 must now be rewritten in the following way.

Proposition 1' Given $\Lambda = \{\lambda_1, \lambda_2, \dots, \lambda_j, \dots\}$ a set of exponents, consider $A_1(p)$ and $A_2(p)$ two exponential sums in the class \mathcal{S}_Λ , say $A_1(p) = \sum_{j \geq 1} a_j e^{\lambda_j p}$ and $A_2(p) = \sum_{j \geq 1} b_j e^{\lambda_j p}$. Fixed a basis G_Λ for Λ , for each $j \geq 1$ let $\mathbf{r}_j \in \mathbb{R}^{\#\Lambda}$ be the vector of rational components verifying (2). Then $A_1 \sim A_2$ if and only if for any integer value $n \geq 1$, with $n \leq \#\Lambda$, there exists $\mathbf{x}_n = (x_{n,1}, x_{n,2}, \dots, x_{n,k}, \dots) \in \mathbb{R}^{\#\Lambda}$ such that $b_j = a_j e^{\langle \mathbf{r}_j, \mathbf{x}_n \rangle}$ for $j = 1, 2, \dots, n$.

Furthermore, if G_Λ is an integral basis for Λ then $A_1 \sim A_2$ if and only if there exists $\mathbf{x}_0 = (x_{0,1}, x_{0,2}, \dots, x_{0,k}, \dots) \in \mathbb{R}^{\#\Lambda}$ such that $b_j = a_j e^{\langle \mathbf{r}_j, \mathbf{x}_0 \rangle}$ for every $j \geq 1$.

Proof For each integer value $n \geq 1$, let V_n be the \mathbb{Q} -vector space generated by $\{\lambda_1, \dots, \lambda_n\}$, V the \mathbb{Q} -vector space generated by Λ , and $G_\Lambda = \{g_1, g_2, \dots, g_k, \dots\}$ a basis of V . If $A_1 \sim A_2$, by Definition 3' for any integer value $n \geq 1$, with $n \leq \#\Lambda$, there exists a \mathbb{Q} -linear map $\psi_n : V_n \rightarrow \mathbb{R}$ such that

$$b_j = a_j e^{i\psi_n(\lambda_j)}, \quad j = 1, 2, \dots, n.$$

Hence $b_j = a_j e^{i \sum_{k=1}^{i_j} r_{j,k} \psi_n(g_k)}$, $j = 1, 2, \dots, n$ or, equivalently,

$$b_j = a_j e^{i \langle \mathbf{r}_j, \mathbf{x}_n \rangle}, \quad j = 1, 2, \dots, n,$$

with $\mathbf{x}_n := (\psi_n(g_1), \psi_n(g_2), \dots)$. Conversely, suppose the existence, for any integer value $n \geq 1$ with $n \leq \sharp \Lambda$, of a vector $\mathbf{x}_n = (x_{n,1}, x_{n,2}, \dots, x_{n,k}, \dots) \in \mathbb{R}^{\sharp G_\Lambda}$ such that $b_j = a_j e^{\langle \mathbf{r}_j, \mathbf{x}_n \rangle i}$, $j = 1, 2, \dots, n$. Thus a \mathbb{Q} -linear map $\psi_n : V_n \rightarrow \mathbb{R}$ can be defined from $\psi_n(g_k) := x_{n,k}$, $k \geq 1$. Therefore

$$\psi_n(\lambda_j) = \sum_{k=1}^{i_j} r_{j,k} \psi(g_k) = \langle \mathbf{r}_j, \mathbf{x}_n \rangle, \quad j = 1, 2, \dots, n,$$

and the result follows.

Now, suppose that G_Λ is an integral basis for Λ and $A_1 \sim A_2$. Thus, by above, for each fixed integer value $n \geq 1$, let $\mathbf{x}_n = (x_{n,1}, x_{n,2}, \dots) \in \mathbb{R}^{\sharp G_\Lambda}$ be a vector such that

$$b_j = a_j e^{i \langle \mathbf{r}_j, \mathbf{x}_n \rangle}, \quad j = 1, 2, \dots, n.$$

Since each component of \mathbf{r}_j is an integer number, without loss of generality, we can take $\mathbf{x}_n \in [0, 2\pi)^{\sharp G_\Lambda}$ as the unique vector in $[0, 2\pi)^{\sharp G_\Lambda}$ satisfying the above equalities, where we assume $x_{n,k} = 0$ for any k such that $r_{j,k} = 0$ for $j = 1, \dots, n$. Therefore, under this assumption, if $m > n$ then $x_{m,k} = x_{n,k}$ for any k so that $x_{n,k} \neq 0$. In this way, we can construct a vector $\mathbf{x}_0 = (x_{0,1}, x_{0,2}, \dots, x_{0,k}, \dots) \in [0, 2\pi)^{\sharp G_\Lambda}$ such that $b_j = a_j e^{\langle \mathbf{r}_j, \mathbf{x}_0 \rangle i}$ for every $j \geq 1$. Indeed, if $r_{1,k} \neq 0$ then the component $x_{0,k}$ is chosen as $x_{1,k}$, and if $r_{1,k} = 0$ then each component $x_{0,k}$ is defined as $x_{n+1,k}$ where $r_{j,k} = 0$ for $j = 1, \dots, n$ and $r_{n+1,k} \neq 0$. Conversely, if there exists $\mathbf{x}_0 = (x_{0,1}, x_{0,2}, \dots, x_{0,k}, \dots) \in \mathbb{R}^{\sharp G_\Lambda}$ such that $b_j = a_j e^{\langle \mathbf{r}_j, \mathbf{x}_0 \rangle i}$ for every $j \geq 1$, then it is clear that $A_1 \sim A_2$ under Definition 3'. \square

From Proposition 1', it is worth noting that old Definition 3 and Definition 3' are equivalent in the case that it is possible to obtain an integral basis for the set of exponents Λ . Consequently, all the results of the quoted above paper which can be formulated in terms of an integral basis are also valid under old Definition 3' (in particular, the mistake above does not concern the results on the finite exponential sums in Section 3 and the Riemann zeta function in Section 5). Moreover, in this case, the set of all exponential sums $A(p)$ in an equivalence class \mathcal{G} in $\mathcal{S}_\Lambda / \sim$ can be determined by a function $E_{\mathcal{G}} : \mathbb{R}^{\sharp G_\Lambda} \rightarrow \mathcal{S}_\Lambda$ of the form

$$E_{\mathcal{G}}(\mathbf{x}) := \sum_{j \geq 1} a_j e^{\langle \mathbf{r}_j, \mathbf{x} \rangle i} e^{\lambda_j p}, \quad \mathbf{x} = (x_1, x_2, \dots, x_k, \dots) \in \mathbb{R}^{\sharp G_\Lambda},$$

where $a_1, a_2, \dots, a_j, \dots$ are the coefficients of an exponential sum in \mathcal{G} and the \mathbf{r}_j 's are the vectors associated with a prefixed integral basis G_Λ for Λ .

However, both definitions 3 and 3' are not equivalent in the general case. For example, consider the set of exponents $\Lambda_0 = \{\lambda_1, \lambda_2, \dots, \lambda_j, \dots\}$ given by

$$\lambda_j = 2j - 1 + \frac{1}{2(2j-1)} \in \mathbb{Q}, \quad j = 1, 2, \dots,$$

then $G_{\Lambda_0} = \{1\}$ is a basis for Λ_0 but it is not an integral basis for Λ_0 . In fact, it is not possible to get an integral basis for Λ_0 . Now, consider the exponential sums $B_1(p) = \sum_{j \geq 1} e^{-\lambda_j p}$ and $B_2(p) = -B_1(p) = \sum_{j \geq 1} -e^{-\lambda_j p}$. Then $B_1(p)$ and $B_2(p)$ are equivalent according to Definition 3'. Indeed, take $a_j = 1$ and $b_j = -1$ for each $j = 1, 2, \dots$, $G_{\Lambda_0} = \{1\}$, $r_j = \lambda_j$ for each $j = 1, 2, \dots$ and $x_n = 2\pi \prod_{k=1}^n (2k-1)$, $n = 1, 2, \dots$. Fixed $n \geq 1$, then it is clear that $b_j = a_j e^{\langle r_j, x_n \rangle}$ for every $j = 1, 2, \dots, n$, i.e.

$$-1 = e^{i(2j-1 + \frac{1}{2(2j-1)}) 2\pi \prod_{k=1}^n (2k-1)}, \quad j = 1, 2, \dots, n.$$

Nevertheless, as it is deduced from Bohr's paper [2], $B_1(p)$ and $B_2(p)$ are not equivalent according to old Definition 3.

Now, *mutatis mutandi*, the proofs of Lemma 1, Lemma 2 and Proposition 3 can be rewritten. Specifically, the modifications are as follows.

- In the proof of Lemma 1: "taking $\mathbf{x}_0 = \tau \mathbf{g}$ " must be changed by "taking $\mathbf{x}_n = \tau \mathbf{g}$ for any integer value $n \geq 1$ ".
- In the proof of Lemma 2: "there exists $\mathbf{x}_0 \in \mathbb{R}^{\sharp \Lambda}$ such that $b_j = a_j e^{\langle r_j, \mathbf{x}_0 \rangle}$ for each $j \geq 1$ " must be changed by "for any integer value $n \geq 1$, with $n \leq \sharp \Lambda$, there exists $\mathbf{x}_n \in \mathbb{R}^{\sharp \Lambda}$ such that $b_j = a_j e^{\langle r_j, \mathbf{x}_n \rangle}$ for each $j = 1, \dots, n$ ".
- In the proof of Lemma 2: "define the sequence of trigonometric polynomials

$$Q_k(t) := \sum_{j \geq 1} p_{j,k} a_j e^{\langle r_j, \mathbf{x}_0 \rangle} e^{i \lambda_j t}, \quad k = 1, 2, \dots$$

must be changed by "define the sequence of trigonometric polynomials

$$Q_k(t) := \sum_{j \geq 1} p_{j,k} b_j e^{i \lambda_j t}, \quad k = 1, 2, \dots,$$

where, fixed k , we can take $b_j = a_j e^{\langle r_j, \mathbf{x}_{n_k} \rangle}$ with n_k the greatest integer value j such that $p_{j,k} \neq 0$."

- In the proof of Lemma 2: "However, note that

$$M\{|Q_{k_1}(t) - Q_{k_2}(t)|^2\} = \sum_{j \geq 1} (p_{j,k_1} - p_{j,k_2})^2 |e^{\langle r_j, \mathbf{x}_0 \rangle}|^2 |a_j|^2 =$$

$$M\{|P_{k_1}(t) - P_{k_2}(t)|^2\},$$

must be changed by "However, note that

$$M\{|Q_{k_1}(t) - Q_{k_2}(t)|^2\} = \sum_{j \geq 1} (p_{j,k_1} - p_{j,k_2})^2 |e^{\langle r_j, \mathbf{x}_{n_{k_2}} \rangle}|^2 |a_j|^2 =$$

$$M\{|P_{k_1}(t) - P_{k_2}(t)|^2\},$$

where we suppose that $n_{k_2} \geq n_{k_1}$."

- In the proof of Lemma 2: “we have $f_2(t) \in AP(\mathbb{R}, \mathbb{C})$. Moreover, $\{Q_k(t)\}_{k \geq 1}$ also converges formally to the series $\sum_{j \geq 1} a_j e^{i \langle \mathbf{r}_j, \mathbf{x}_0 \rangle} e^{i \lambda_j t}$, which, by [1, p. 21], represents the Fourier series of $f_2(t)$. Finally, by taking into account Definition 5 (in terms of Proposition 1) we have $f_1 \sim f_2$.” must be changed by “we have $f_2(t) \in AP(\mathbb{R}, \mathbb{C})$ and, by [1, p. 21], $\sum_{j \geq 1} b_j e^{i \lambda_j t}$ represents its Fourier series. Finally, by taking into account Definition 5 (in terms of Proposition 1') we have $f_1 \sim f_2$.”
- In the proof of Proposition 3: “Since $f_1 \sim f_l$ for each $l = 1, 2, \dots$, we deduce from Proposition 1 that there exists $\mathbf{x}_l = (x_{l,1}, x_{l,2}, \dots) \in \mathbb{R}^{\sharp G_A}$ such that

$$a_{l,j} = a_{1,j} e^{i \langle \mathbf{r}_j, \mathbf{x}_l \rangle}, \quad j = 1, 2, \dots, \quad \text{with } \lambda_j \in A.” \quad (15)$$

must be changed by “Since $f_1 \sim f_l$ for each $l = 1, 2, \dots$, we deduce from Proposition 1' that for any integer value $n \geq 1$, with $n \leq \sharp A$, there exists $\mathbf{x}_{l,n} = (x_{l,n,1}, x_{l,n,2}, \dots) \in \mathbb{R}^{\sharp G_A}$ such that

$$a_{l,j} = a_{1,j} e^{i \langle \mathbf{r}_j, \mathbf{x}_{l,n} \rangle}, \quad j = 1, 2, \dots, n \quad \text{with } \lambda_j \in A.” \quad (15)$$

- In the proof of Proposition 3: “and, since (16) is satisfied for any $k = 1, 2, \dots$, we can also construct a vector $\mathbf{x}_0 \in \mathbb{R}^{\sharp G_A}$ such that, by taking into account remarks 1 and 2, verifies

$$a_j = a_{1,j} e^{i \langle \mathbf{r}_j, \mathbf{x}_0 \rangle}, \quad j = 1, 2, \dots, \quad \text{with } \lambda_j \in A.”$$

must be changed by “and, since (16) is satisfied for any $k = 1, 2, \dots$, we can construct, for any integer value $n \geq 1$ with $n \leq \sharp A$, a vector $\mathbf{x}_{0,n} \in \mathbb{R}^{\sharp G_A}$ such that, by taking into account remarks 1 and 2, verifies

$$a_j = a_{1,j} e^{i \langle \mathbf{r}_j, \mathbf{x}_{0,n} \rangle}, \quad j = 1, 2, \dots, n \quad \text{with } \lambda_j \in A.”$$

The rest of the article remains unchanged.

Finally, the authors wish to thank Mattia Righetti for his quote to Bohr's example given by the set of exponents A_0 considered above [2, 3].

References

1. Besicovitch, A.S.: Almost periodic functions, Dover, New York, 1954.
2. Bohr, H.: Zur Theorie der allgemeinen Dirichletschen Reihen. (German) Math. Ann. **79**, 136-156 (1918).
3. Righetti, M.: On Bohr's equivalence theorem. J. Math. Anal. Appl., corrigendum *ibid.* 449, 939-940 (2017).
4. Sepulcre, J.M., Vidal, T.: Almost periodic functions in terms of Bohr's equivalence relation. Ramanujan J., **46** (1), 245–267 (2018).